

A BLUEPRINT FOR THE FORMALIZATION OF CARLESON'S THEOREM ON CONVERGENCE OF FOURIER SERIES

LARS BECKER, MARÍA INÉS DE FRUTOS-FERNÁNDEZ, LEO DIEDERING,
FLORIS VAN DOORN, SÉBASTIEN GOUËZEL, ASGAR JAMNESHAN,
EVGENIA KARUNUS, EDWARD VAN DE MEENT, PIETRO MONTICONE,
JASPER MULDER-SOHN, JIM PORTEGIES, JORIS ROOS, MICHAEL ROTHGANG,
RAJULA SRIVASTAVA, JAMES SUNDSTROM, JEREMY TAN, AND CHRISTOPH THIELE

ABSTRACT. This paper is the blueprint underlying the Lean formalization of the proof of Carleson's classical result [Car66] asserting almost everywhere convergence of Fourier series of continuous functions. We break up the proof into two steps, a reduction of the classical result to a new theorem that appears in the sibling communication [BvDJ⁺24] and a proof of this new theorem, which is also detailed as blueprint in this paper. An early version of this blueprint was used to initiate the Lean formalization. During the formalization, many contributors elaborated the blueprint with minor corrections, modifications and extensions. The final version is presented here as a guide through the accompanying Lean code.

CONTENTS

1. Introduction	2
1.1. Statement of the metric space Carleson theorems	5
2. Proof of Metric Space Carleson, overview	8
2.1. Auxiliary lemmas	14
3. Proof of Metric Space Carleson	16
4. Proof of Finitary Carleson	21
4.1. Proof of Grid Existence Lemma	22
4.2. Proof of Tile Structure Lemma	28
5. Proof of discrete Carleson	31
5.1. Organisation of the tiles	32
5.2. Proof of the Exceptional Sets Lemma	34
5.3. Auxiliary lemmas	39
5.4. Proof of the Forest Union Lemma	42
5.5. Proof of the Forest Complement Lemma	47
6. Proof of the Antichain Operator Proposition	50
6.1. The density arguments	50
6.2. Proof of the Tile Correlation Lemma	54
6.3. Proof of the Antichain Tile Count Lemma	57
7. Proof of the Forest Operator Proposition	62
7.1. The pointwise tree estimate	62

Date: June 10, 2026.

7.2.	An auxiliary L^2 tree estimate	68
7.3.	The quantitative L^2 tree estimate	73
7.4.	Almost orthogonality of separated trees	76
7.5.	Proof of the Tiles with large separation Lemma	79
7.6.	Proof of The Remaining Tiles Lemma	89
7.7.	Forests	93
8.	Proof of the Hölder cancellative condition	97
9.	Proof of Vitali covering and Hardy–Littlewood	100
10.	Two-sided Metric Space Carleson	104
10.1.	Proof of Cotlar’s Inequality	106
10.2.	Calderón-Zygmund Decomposition	112
11.	Proof of The Classical Carleson Theorem	122
11.1.	The classical Carleson theorem	123
11.2.	Smooth functions.	128
11.3.	The truncated Hilbert transform	128
11.4.	The proof of the van der Corput Lemma	133
11.5.	Partial sums as orthogonal projections	134
11.6.	The error bound	135
11.7.	Carleson on the real line	138
	References	141

1. INTRODUCTION

Trigonometric series represent functions as possibly infinite linear combinations of pure frequencies. They gained particular prominence through the work of J. Fourier, who used them in his analytical theory of heat [Fou22], see also [Fou13], thereby establishing them as a tool for solving partial differential equations. Fourier also made the groundbreaking claim that a wide range of functions could be represented using trigonometric series. This sparked the interest of many mathematicians, including Dirichlet, who gave some rigorous conditions for convergence of Fourier series, as trigonometric series are now called. Dirichlet also opened a branch of analytic number theory partially inspired by the ideas of Fourier. Nowadays, Fourier analysis plays an important role in many areas of mathematics.

With Euler’s formula to represent pure frequencies in mind, a trigonometric polynomial can be expressed as

$$S_N(x) := \sum_{n=-N}^N c_n e^{inx} . \quad (1.0.1)$$

The Fourier series is then defined as the limit f of such a sequence S_N as N tends to ∞ . Fourier’s vision to represent rather general functions raises two fundamental questions. The first question is to identify the appropriate choice of coefficients c_n to use to represent a given f . The second question addresses the convergence of S_N to f .

The first question has a fairly canonical and standard answer, provided by the Fourier integral formula:

$$c_n := \hat{f}_n := \frac{1}{2\pi} \int_0^{2\pi} f(x)e^{-inx} dx, \tag{1.0.2}$$

where the precise interpretation of the integral depends on the chosen theory of integration. For a continuous function f , Riemann's notion of the integral suffices. If f is integrable in the Lebesgue sense, $f \in L^1(0, 2\pi)$, the Lebesgue integral is appropriate. More generally, if f is a distribution in the sense of Schwartz, supported in $[0, 2\pi]$ the integral can be understood as an evaluation against the periodic test function e^{-inx} . In each case, the more general definition reduces to the simpler one within the respective more restrictive domain. Hence, the Fourier coefficients given by (1.0.2) serve as a universal choice. This choice is unique in several respects, in particular if one is to exactly reproduce a trigonometric polynomial f in the form (1.0.1).

The second question of convergence bifurcates into the question of pointwise convergence of the series (1.0.1) (with coefficients given by (1.0.2)) for a given x on the one hand and convergence of the functions S_N to the function f in a suitable function space with corresponding topology on the other hand. There are at least as many function spaces for the question of convergence as there are different definitions of the integral elaborated earlier. There are some very canonical answers to the convergence question in function spaces, albeit not known at the time of Fourier and Dirichlet. One example is convergence in the Hilbert space sense for f in $L^2(0, 2\pi)$, as discovered in the first decade of the twentieth century as a consequence of the rapid development of Lebesgue integration theory. Another canonical example is convergence in the sense of distributions for general distributional f , as discovered a few decades after Lebesgue integration. For some other natural spaces, such as $L^1(0, 2\pi)$, there is no guarantee of convergence in the norm of that space even if f is in the space.

In contrast to these examples of function spaces with a very natural theory of convergence of Fourier series in the topology of the function space, there are no similarly elegant solutions to the characterization of pointwise convergence. In particular, the space of functions f such that the sequence $S_N(x)$ converges for every x does not have a good characterization in terms of f itself. Similarly, the space of all functions f such that the sequence of coefficients \hat{f}_n is absolutely summable has also no good characterization.

When the Fourier integral is defined in the Lebesgue sense and $f \in L^1(0, 2\pi)$, then the function f itself is meaningful not everywhere but only pointwise almost everywhere in the Lebesgue sense. The question of pointwise convergence to f for all x then becomes meaningless, and instead one asks for almost everywhere convergence. Such convergence was conjectured by N. Luzin [Luz13] for the space $L^2(0, 2\pi)$, see also the collected works [Luz53]. Kolmogorov's example [Kol23] of an L^1 function whose Fourier series diverges at almost every point seemed to provide some evidence against Luzin's conjecture. Indeed, in the 1960s, L. Carleson tried to construct a counterexample to Luzin's conjecture. In his own recollection, his efforts led him to

better and better understand how such a potential counterexample should look like. In the end, the counterexample had to satisfy so many properties that the properties started to contradict each other, and Carleson realized that a counterexample could not exist. Thus, he had proved Luzin's conjecture [Car66]. In particular, he had proven the more elementary statement

Theorem 1.0.1 (classical Carleson). *Let f be a 2π -periodic complex-valued continuous function on \mathbb{R} . Then for almost all $x \in \mathbb{R}$ we have*

$$\lim_{N \rightarrow \infty} S_N f(x) = f(x), \quad (1.0.3)$$

where $S_N f$ is the N -th partial Fourier sum of f defined in (1.0.1) with coefficients (1.0.2).

Here, almost every x means in the Lebesgue sense, i.e., for every $\epsilon > 0$ the set of x where convergence fails can be covered by a sequence of intervals such that the sum of the lengths of these intervals is less than ϵ . While Carleson had proven the more general Luzin conjecture for functions in $L^2[0, 2\pi]$, even the more elementary statement for continuous functions was not known before Carleson's work. Moreover, until now, the elementary statement has not seen any substantially easier proof than those generalizing to L^2 , partially because there is no readily usable criterion on the level of Fourier coefficients to distinguish between continuous functions and L^2 functions.

Shortly after Carleson's breakthrough, Hunt [Hun68] proved the analogous result for L^p functions with $p > 1$ and for functions in $L^1 \log(L)^2$. Billard [Bil67] adapted Carleson's arguments to prove almost everywhere convergence of Walsh-Fourier series for functions in L^2 .

In [Fef73], C. Fefferman gave an alternative proof of Carleson's theorem via an a priori bound for Carleson's operator, the maximally modulated singular integral

$$Tf(x) := \sup_N \int_{-\pi}^{\pi} e^{iNy} f(x-y) \frac{1}{y} dy. \quad (1.0.4)$$

In [LT00], a duality between the approaches by Carleson and Fefferman was pointed out and a more symmetric and self-dual proof was presented.

Various strengthenings of Fefferman's estimates for Carleson's operator have appeared since, such as bounds for a higher dimensional Carleson operator in the isotropic [Sjö71], [PT03] and anisotropic [Roo19] setting. The supremum norm in the variable N in (1.0.4) was strengthened to maximal multiplier norms in [DLT⁺08] with applications to return times theorems in ergodic theory. It was strengthened to variation norms V^r with $r > 2$ in [OST⁺12] with applications to nonlinear analysis.

Stein and Wainger [SW01] proposed to study variants of (1.0.4) replacing the linear modulation phase Ny by polynomial phases in y and proved a result for polynomials without a linear term. This was generalized by Lie to general polynomial Carleson operators, first for a supremum over all quadratic polynomials [Lie09] and then for a supremum over all polynomials in [Lie20]. The polynomial Carleson operator was further generalized to higher dimensions and Hölder regular kernels in

[Zor21]. The development of polynomial Carleson operators has led to further generalisations, for example to almost everywhere convergence of Malmquist-Takenaka series [Mna22] and maximally modulated singular Radon transforms [Ram21], [Bec24].

Surveys to Carleson's theorem can be found in [Lac04], [Dem15].

In this blueprint, we prove Theorem 1.0.1 as a corollary to a further generalization of the polynomial Carleson operator towards doubling metric measure spaces, Theorem 1.1.1 below, which is new and appears in the sibling communication [BvDJ⁺24]. Theorem 1.1.1 is an axiomatic approach to Carleson type theorems on doubling metric measure spaces. This axiomatic approach is suitable for formalization and a good route towards the classical theorem.

Early drafts of the sibling communication [BvDJ⁺24] existed in summer 2023. Based on this, the first draft of the present blueprint was written in the first half of 2024, containing a much more detailed proof, which involved increasing the size by a factor of four, and adding the derivation of Carleson's classical result. In June 2024, the fourth author launched a public website to post the blueprint, using the open-source software `leanblueprint` developed by Patrick Massot, calling for contributions to formalize the proof. The goal was to formalize the blueprint in the Lean proof assistant [dMU21], building on top of its mathematical library `Mathlib` [mat20]. The work was split up into about 180 tasks, to be claimed by individual contributors. Most tasks were to formalize the proof of a single lemma from the blueprint, and some were to develop basic theory or refactor existing code. The contributors adapted the blueprint to fix some gaps found during the formalization and gave feedback that led to discussions about the proof. This even resulted in a few changes to the general setup and the main theorems. All of the gaps found required only fairly localized changes to the blueprint, indicating that the initial blueprint was already of high quality. The formalization was completed in July 2025. It is attached to this arXiv posting, and the latest version can be found on Github.

Everyone that completed a substantial amount of tasks is included as a coauthor of the blueprint. The authors acknowledge contributions in the form of small formalization additions, pointing out corrections to the blueprint, or supplying ideas to the Lean efforts by the following people: Michel Alexis, Bolton Bailey, Julian Berman, Joachim Breitner, Martin Dvořák, Georges Gonthier, Aaron Hill, Austin Letson, Bhavik Mehta, Eric Paul, Clara Torres, Dennis Tsar, Andrew Yang, Ruben van de Velde.

Acknowledgement. L.B., M.I.d.F.F., L.D., F.v.D., M.R., R.S., and C.T. were funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2047/1 – 390685813. L.B., R.S., and C.T. were also supported by SFB 1060. A.J. is funded by the TÜBITAK (Scientific and Technological Research Council of Türkiye) under Grant Number 123F122. J.R. was supported in part by NSF grant DMS-2154835 and a HIM fellowship for the Fall 2024 trimester program in Bonn.

1.1. Statement of the metric space Carleson theorems. Let

$$a \geq 4 \tag{1.1.1}$$

be a natural number that as it gets larger will allow for more general applications of Theorem 1.1.1 but will worsen the constants in the estimates.

A doubling metric measure space (X, ρ, μ, a) is a complete and locally compact metric space (X, ρ) equipped with a non-zero locally finite Borel measure μ that satisfies the doubling condition that for all $x \in X$ and all $R > 0$ we have

$$\mu(B(x, 2R)) \leq 2^a \mu(B(x, R)), \quad (1.1.2)$$

where we have denoted by $B(x, R)$ the open ball of radius R centered at x :

$$B(x, R) := \{y \in X : \rho(x, y) < R\}. \quad (1.1.3)$$

In a doubling metric measure space the closed balls are compact and μ is positive on all non-empty open sets.

A collection Θ of real valued continuous functions on the doubling metric measure space (X, ρ, μ, a) is called compatible, if there is a point $o \in X$ where all the functions are equal to 0, and if there exists for each ball $B \subset X$ a metric d_B on Θ , such that the following five properties (1.1.4), (1.1.5), (1.1.6), (1.1.7), and (1.1.8) are satisfied. For every ball $B \subset X$

$$\sup_{x, y \in B} |\vartheta(x) - \vartheta(y) - \theta(x) + \theta(y)| \leq d_B(\vartheta, \theta). \quad (1.1.4)$$

For any two balls $B_1 = B(x_1, R)$, $B_2 = B(x_2, 2R)$ in X with $x_1 \in B_2$ and any $\vartheta, \theta \in \Theta$,

$$d_{B_2}(\vartheta, \theta) \leq 2^a d_{B_1}(\vartheta, \theta). \quad (1.1.5)$$

For any two balls B_1, B_2 in X with $B_1 \subset B_2$ and any $\vartheta, \theta \in \Theta$

$$d_{B_1}(\vartheta, \theta) \leq d_{B_2}(\vartheta, \theta) \quad (1.1.6)$$

and for any two balls $B_1 = B(x_1, R)$, $B_2 = B(x_2, 2^a R)$ with $B_1 \subset B_2$, and $\vartheta, \theta \in \Theta$,

$$2d_{B_1}(\vartheta, \theta) \leq d_{B_2}(\vartheta, \theta). \quad (1.1.7)$$

For every ball B in X and every d_B -ball \tilde{B} of radius $2R$ in Θ , there is a collection \mathcal{B} of at most 2^a many d_B -balls of radius R covering \tilde{B} , that is,

$$\tilde{B} \subset \bigcup \mathcal{B}. \quad (1.1.8)$$

Further, a compatible collection Θ is called cancellative, if for any ball B in X of radius R , any Lipschitz function $\varphi : X \rightarrow \mathbb{C}$ supported on B , and any $\vartheta, \theta \in \Theta$ we have

$$\left| \int_B e(\vartheta(x) - \theta(x)) \varphi(x) d\mu(x) \right| \leq 2^a \mu(B) \|\varphi\|_{\text{Lip}(B)} (1 + d_B(\vartheta, \theta))^{-\frac{1}{a}}, \quad (1.1.9)$$

where $\|\cdot\|_{\text{Lip}(B)}$ denotes the inhomogeneous Lipschitz norm on B :

$$\|\varphi\|_{\text{Lip}(B)} = \sup_{x \in B} |\varphi(x)| + R \sup_{x, y \in B, x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)}.$$

A one-sided Calderón–Zygmund kernel K on the doubling metric measure space (X, ρ, μ, a) is a measurable function

$$K : X \times X \rightarrow \mathbb{C} \quad (1.1.10)$$

such that for all $x, y', y \in X$ with $x \neq y$, we have

$$|K(x, y)| \leq \frac{2^{a^3}}{V(x, y)} \quad (1.1.11)$$

and if $2\rho(y, y') \leq \rho(x, y)$, then

$$|K(x, y) - K(x, y')| \leq \left(\frac{\rho(y, y')}{\rho(x, y)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x, y)}, \quad (1.1.12)$$

where

$$V(x, y) := \mu(B(x, \rho(x, y))).$$

Define the maximally truncated non-tangential singular integral T_* associated with K by

$$T_*f(x) := \sup_{R_1 < R_2} \sup_{\rho(x, x') < R_1} \left| \int_{R_1 < \rho(x', y) < R_2} K(x', y) f(y) d\mu(y) \right|. \quad (1.1.13)$$

We define the generalized Carleson operator T by

$$Tf(x) := \sup_{\vartheta \in \Theta} \sup_{0 < R_1 < R_2} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right|, \quad (1.1.14)$$

where $e(r) = e^{ir}$.

Our first main result is the following restricted weak type estimate for T in the range $1 < q \leq 2$, which by interpolation techniques recovers L^q estimates for the open range $1 < q < 2$.

Theorem 1.1.1 (metric space Carleson). *For all integers $a \geq 4$ and real numbers $1 < q \leq 2$ the following holds. Let (X, ρ, μ, a) be a doubling metric measure space. Let Θ be a cancellative compatible collection of functions and let K be a one-sided Calderón–Zygmund kernel on (X, ρ, μ, a) . Assume that for every bounded measurable function g on X supported on a set of finite measure we have*

$$\|T_*g\|_2 \leq 2^{a^3} \|g\|_2, \quad (1.1.15)$$

where T_* is defined in (1.1.13). Then for all Borel sets F and G in X and all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$, we have, with T defined in (1.1.14),

$$\left| \int_G Tf d\mu \right| \leq \frac{2^{443a^3}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (1.1.16)$$

In some applications, such as the Walsh-case of Carleson's theorem, the kernel K naturally depends also on the modulation functions ϑ . The fact that we don't assume Hölder-continuity of the kernel K in the first argument allows us to also capture this situation.

We now state another Theorem with a slightly weaker replacement of the assumption (1.1.15). It implies the Walsh-case of Carleson's theorem. For a Borel function $Q : X \rightarrow \Theta$, and $\vartheta \in \Theta$ and $x \in X$ define

$$R_Q(\vartheta, x) = \sup\{r > 0 : d_{B(x, r)}(\vartheta, Q(x)) < 1\} \quad (1.1.17)$$

and define further

$$T_Q^\vartheta f(x) := \sup_{R_1 < R_2} \sup_{\rho(x, x') < R_1} \left| \int_{R_1 < \rho(x', y) < \min\{R_2, R_Q(\vartheta, x')\}} K(x', y) f(y) \, d\mu(y) \right| \quad (1.1.18)$$

Define further the linearized generalized Carleson operator T_Q by

$$T_Q f(x) := \sup_{0 < R_1 < R_2} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(Q(x)(y)) \, d\mu(y) \right|, \quad (1.1.19)$$

where again $e(r) = e^{ir}$.

Theorem 1.1.2 (linearised metric Carleson). *For all integers $a \geq 4$ and real numbers $1 < q \leq 2$ the following holds. Let (X, ρ, μ, a) be a doubling metric measure space. Let Θ be a cancellative compatible collection of functions. Let $Q : X \rightarrow \Theta$ be a Borel function with finite range. Let K be a one-sided Calderón–Zygmund kernel on (X, ρ, μ, a) . Assume that for every $\vartheta \in \Theta$ and every bounded measurable function g on X supported on a set of finite measure we have*

$$\|T_Q^\vartheta g\|_2 \leq 2^{a^3} \|g\|_2, \quad (1.1.20)$$

where T_Q^ϑ is defined in (1.1.18). Then for all Borel sets F and G in X and all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$, we have, with T_Q defined in (1.1.19),

$$\left| \int_G T_Q f \, d\mu \right| \leq \frac{2^{443a^3}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (1.1.21)$$

Remark 1.1.3. The value of the constant factor 2^{443a^3} in Theorems 1.1.1 and 1.1.2 is by no means sharp. One source of non-sharpness is our choice to write for readability most constants in the form 2^{na^3} for some explicit constant n .

An a posteriori byproduct of the Lean formalization of this document is that Theorems 1.1.1 and 1.1.2 remain true with 2^{44a^3} instead of 2^{443a^3} . This is obtained by changing the parameter D introduced in (2.0.1) from 2^{100a^2} to 2^{7a^2} , checking that exactly the same proof goes through, tracking the constants, and getting 2^{44a^3} in the end. This value is again by no means sharp. In the formalization we define the parameter D as 2^{ca^2} and the constants in this blueprint are obtained by setting $c = 100$.

2. PROOF OF METRIC SPACE CARLESON, OVERVIEW

This section organizes the proof of Theorem 1.1.1 into sections 3, 4, 5, 6, 7, 8, and 9. These sections are mutually independent except for referring to the statements formulated in the present section. Section 3 proves the main Theorem 1.1.1, while sections 4, 5, 6, 7, 8, and 9 each prove one proposition that is stated in the present section. The present section also introduces all definitions used across these sections.

Section 2.1 proves some auxiliary lemmas that are used in more than one of the sections 3-9.

Let a, q be given as in Theorem 1.1.1.

Define

$$D := 2^{100a^2}, \quad (2.0.1)$$

$$\kappa := 2^{-10a}, \quad (2.0.2)$$

and

$$Z := 2^{12a}. \quad (2.0.3)$$

Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ be the unique compactly supported, piece-wise linear, continuous function with corners precisely at $\frac{1}{4D}$, $\frac{1}{2D}$, $\frac{1}{4}$ and $\frac{1}{2}$ which satisfies

$$\sum_{s \in \mathbb{Z}} \psi(D^{-s}x) = 1 \quad (2.0.4)$$

for all $x > 0$. This function vanishes outside $[\frac{1}{4D}, \frac{1}{2}]$, is constant one on $[\frac{1}{2D}, \frac{1}{4}]$, and is Lipschitz with constant $4D$.

Let a doubling metric measure space (X, ρ, μ, a) be given. Let a cancellative compatible collection Θ of functions on X be given. Let $o \in X$ be a point such that $\vartheta(o) = 0$ for all $\vartheta \in \Theta$.

Let a Borel measurable function $Q : X \rightarrow \Theta$ with finite range be given. Let a one-sided Calderón–Zygmund kernel K on X be given so that for every $\vartheta \in \Theta$ the operator T_Q^ϑ defined in (1.1.18) satisfies (1.1.20).

For $s \in \mathbb{Z}$, we define

$$K_s(x, y) := K(x, y)\psi(D^{-s}\rho(x, y)), \quad (2.0.5)$$

so that for each $x, y \in X$ with $x \neq y$ we have

$$K(x, y) = \sum_{s \in \mathbb{Z}} K_s(x, y).$$

In Section 3, we prove Theorem 1.1.1 and Theorem 1.1.2 from the finitary version, Proposition 2.0.1 below. Recall that a function from a measure space to a finite set is measurable if the pre-image of each of the elements in the range is measurable.

Proposition 2.0.1 (finitary Carleson). *Let $\sigma_1, \sigma_2 : X \rightarrow \mathbb{Z}$ be measurable functions with finite range and $\sigma_1 \leq \sigma_2$. Let F, G be bounded Borel sets in X . Then there is a Borel set G' in X with $2\mu(G') \leq \mu(G)$ such that for all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$.*

$$\begin{aligned} \int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y)) \, d\mu(y) \right| d\mu(x) \\ \leq \frac{2^{442a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \end{aligned} \quad (2.0.6)$$

Let measurable functions $\sigma_1 \leq \sigma_2 : X \rightarrow \mathbb{Z}$ with finite range be given. Let bounded Borel sets F, G in X be given. Let S be the smallest integer such that the ranges of σ_1 and σ_2 are contained in $[-S, S]$ and F and G are contained in the ball $B(o, \frac{1}{4}D^S)$.

In Section 4, we prove Proposition 2.0.1 using a bound for a dyadic model formulated in Proposition 2.0.2 below.

A grid structure (\mathcal{D}, c, s) on X consists of a finite collection \mathcal{D} of pairs (I, k) of Borel sets in X and integers $k \in [-S, S]$, the projection $s : \mathcal{D} \rightarrow [-S, S]$, $(I, k) \mapsto k$ to the second component which is assumed to be surjective and called scale function, and a function $c : \mathcal{D} \rightarrow X$ called center function such that the five properties (2.0.7),

(2.0.8), (2.0.9), (2.0.10), and (2.0.11) hold. We call the elements of \mathcal{D} dyadic cubes. By abuse of notation, we will usually write just I for the cube (I, k) , and we will write $I \subset J$ to mean that for two cubes $(I, k), (J, l) \in \mathcal{D}$ we have $I \subset J$ and $k \leq l$.

For each dyadic cube I and each $-S \leq k < s(I)$ we have

$$I \subset \bigcup_{J \in \mathcal{D}: s(J)=k} J. \quad (2.0.7)$$

Any two non-disjoint dyadic cubes I, J with $s(I) \leq s(J)$ satisfy

$$I \subset J. \quad (2.0.8)$$

There exists a $I_0 \in \mathcal{D}$ with $s(I_0) = S$ and $c(I_0) = o$ and for all cubes $J \in \mathcal{D}$, we have

$$J \subset I_0. \quad (2.0.9)$$

For any dyadic cube I ,

$$c(I) \in B(c(I), \frac{1}{4}D^{s(I)}) \subset I \subset B(c(I), 4D^{s(I)}). \quad (2.0.10)$$

For any dyadic cube I and any t with $tD^{s(I)} \geq D^{-S}$,

$$\mu(\{x \in I : \rho(x, X \setminus I) \leq tD^{s(I)}\}) \leq 2t^\kappa \mu(I). \quad (2.0.11)$$

A tile structure $(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q}, c, s)$ for a given grid structure (\mathcal{D}, c, s) is a finite set \mathfrak{P} of elements called tiles with five maps

$$\begin{aligned} \mathcal{J}: \mathfrak{P} &\rightarrow \mathcal{D} \\ \Omega: \mathfrak{P} &\rightarrow \mathcal{P}(\Theta) \\ \mathcal{Q}: \mathfrak{P} &\rightarrow \mathcal{Q}(X) \\ c: \mathfrak{P} &\rightarrow X \\ s: \mathfrak{P} &\rightarrow \mathbb{Z} \end{aligned}$$

with \mathcal{J} surjective and $\mathcal{P}(\Theta)$ denoting the power set of Θ such that the five properties (2.0.13), (2.0.14), (2.0.15), (2.0.18), and (2.0.19) hold. For each dyadic cube I , the restriction of the map Ω to the set

$$\mathfrak{P}(I) = \{\mathfrak{p} : \mathcal{J}(\mathfrak{p}) = I\} \quad (2.0.12)$$

is injective and we have the disjoint covering property (we use the union symbol with dot on top to denote a disjoint union)

$$\mathcal{Q}(X) \subset \dot{\bigcup}_{\mathfrak{p} \in \mathfrak{P}(I)} \Omega(\mathfrak{p}). \quad (2.0.13)$$

For any tiles $\mathfrak{p}, \mathfrak{q}$ with $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{q})$ and $\Omega(\mathfrak{p}) \cap \Omega(\mathfrak{q}) \neq \emptyset$ we have

$$\Omega(\mathfrak{q}) \subset \Omega(\mathfrak{p}). \quad (2.0.14)$$

For each tile \mathfrak{p} ,

$$\mathcal{Q}(\mathfrak{p}) \in B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.2) \subset \Omega(\mathfrak{p}) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 1), \quad (2.0.15)$$

where

$$B_{\mathfrak{p}}(\vartheta, R) := \{\theta \in \Theta : d_{\mathfrak{p}}(\vartheta, \theta) < R\}, \quad (2.0.16)$$

and

$$d_{\mathfrak{p}} := d_{B(c(\mathfrak{p}), \frac{1}{4}D^{s(\mathfrak{p})})}. \quad (2.0.17)$$

We have for each tile \mathfrak{p}

$$c(\mathfrak{p}) = c(\mathcal{J}(\mathfrak{p})), \quad (2.0.18)$$

$$s(\mathfrak{p}) = s(\mathcal{J}(\mathfrak{p})). \quad (2.0.19)$$

Proposition 2.0.2 (discrete Carleson). *Let (\mathcal{D}, c, s) be a grid structure and*

$$(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q}, c, s)$$

a tile structure for this grid structure. Define for $\mathfrak{p} \in \mathfrak{P}$

$$E(\mathfrak{p}) = \{x \in \mathcal{J}(\mathfrak{p}) : Q(x) \in \Omega(\mathfrak{p}), \sigma_1(x) \leq s(\mathfrak{p}) \leq \sigma_2(x)\} \quad (2.0.20)$$

and

$$T_{\mathfrak{p}}f(x) = \mathbf{1}_{E(\mathfrak{p})}(x) \int K_{s(\mathfrak{p})}(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (2.0.21)$$

Then there exists a Borel set G' with $2\mu(G') \leq \mu(G)$ such that for all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{P}} T_{\mathfrak{p}}f(x) \right| d\mu(x) \leq \frac{2^{442a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (2.0.22)$$

The proof of Proposition 2.0.2 is done in Section 5 by a reduction to two further propositions that we state below.

Fix a grid structure (\mathcal{D}, c, s) and a tile structure $(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q}, c, s)$ for this grid structure.

We define the relation

$$\mathfrak{p} \leq \mathfrak{p}' \quad (2.0.23)$$

on $\mathfrak{P} \times \mathfrak{P}$ meaning $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p}')$ and $\Omega(\mathfrak{p}') \subset \Omega(\mathfrak{p})$. We further define for $\lambda, \lambda' > 0$ the relation

$$\lambda \mathfrak{p} \lesssim \lambda' \mathfrak{p}' \quad (2.0.24)$$

on $\mathfrak{P} \times \mathfrak{P}$ meaning $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p}')$ and

$$B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \lambda') \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \lambda). \quad (2.0.25)$$

Define for a tile \mathfrak{p} and $\lambda > 0$

$$E_1(\mathfrak{p}) := \{x \in \mathcal{J}(\mathfrak{p}) \cap G : Q(x) \in \Omega(\mathfrak{p})\}, \quad (2.0.26)$$

$$E_2(\lambda, \mathfrak{p}) := \{x \in \mathcal{J}(\mathfrak{p}) \cap G : Q(x) \in B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \lambda)\}. \quad (2.0.27)$$

Given a subset \mathfrak{P}' of \mathfrak{P} , we define $\mathfrak{P}(\mathfrak{P}')$ to be the set of all $\mathfrak{p} \in \mathfrak{P}$ such that there exist $\mathfrak{p}' \in \mathfrak{P}'$ with $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p}')$. Define the densities

$$\text{dens}_1(\mathfrak{P}') := \sup_{\mathfrak{p}' \in \mathfrak{P}'} \sup_{\lambda \geq 2} \lambda^{-a} \sup_{\mathfrak{p} \in \mathfrak{P}(\mathfrak{P}'), \lambda \mathfrak{p}' \lesssim \mathfrak{p}} \frac{\mu(E_2(\lambda, \mathfrak{p}))}{\mu(\mathcal{J}(\mathfrak{p}))}, \quad (2.0.28)$$

$$\text{dens}_2(\mathfrak{P}') := \sup_{\mathfrak{p}' \in \mathfrak{P}'} \sup_{r \geq 4D^s(\mathfrak{p})} \frac{\mu(F \cap B(c(\mathfrak{p}), r))}{\mu(B(c(\mathfrak{p}), r))}. \quad (2.0.29)$$

An antichain is a subset \mathfrak{A} of \mathfrak{P} such that for any distinct $\mathfrak{p}, \mathfrak{q} \in \mathfrak{A}$ we do not have $\mathfrak{p} \leq \mathfrak{q}$.

The following proposition is proved in Section 6.

Proposition 2.0.3 (antichain operator). *For any antichain \mathfrak{A} and for all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ and all $g : X \rightarrow \mathbb{C}$ with $|g| \leq \mathbf{1}_G$*

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x) \right| \quad (2.0.30)$$

$$\leq \frac{2^{117a^3}}{q-1} \text{dens}_1(\mathfrak{A})^{\frac{q-1}{8a^4}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \quad (2.0.31)$$

Let $n \geq 0$. An n -forest is a pair $(\mathfrak{U}, \mathfrak{T})$ where \mathfrak{U} is a subset of \mathfrak{P} and \mathfrak{T} is a map assigning to each $\mathfrak{u} \in \mathfrak{U}$ a nonempty set $\mathfrak{T}(\mathfrak{u}) \subset \mathfrak{P}$ called tree such that the following properties (2.0.32), (2.0.33), (2.0.34), (2.0.35), (2.0.36), and (2.0.37) hold.

For each $\mathfrak{u} \in \mathfrak{U}$ and each $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$ we have $\mathcal{J}(\mathfrak{p}) \neq \mathcal{J}(\mathfrak{u})$ and

$$4\mathfrak{p} \lesssim \mathfrak{u}. \quad (2.0.32)$$

For each $\mathfrak{u} \in \mathfrak{U}$ and each $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{T}(\mathfrak{u})$ and $\mathfrak{p}' \in \mathfrak{P}$ we have

$$\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{T}(\mathfrak{u}), \mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}'' \implies \mathfrak{p}' \in \mathfrak{T}(\mathfrak{u}). \quad (2.0.33)$$

We have

$$\left\| \sum_{\mathfrak{u} \in \mathfrak{U}} \mathbf{1}_{\mathcal{J}(\mathfrak{u})} \right\|_{\infty} \leq 2^n. \quad (2.0.34)$$

We have for every $\mathfrak{u} \in \mathfrak{U}$

$$\text{dens}_1(\mathfrak{T}(\mathfrak{u})) \leq 2^{4a+1-n}. \quad (2.0.35)$$

We have for $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}$ with $\mathfrak{u} \neq \mathfrak{u}'$ and $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u}')$ with $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{u})$ that

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{u})) > 2^{Z(n+1)}. \quad (2.0.36)$$

We have for every $\mathfrak{u} \in \mathfrak{U}$ and $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$ that

$$B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})}) \subset \mathcal{J}(\mathfrak{u}). \quad (2.0.37)$$

The following proposition is proved in Section 7.

Proposition 2.0.4 (forest operator). *For any $n \geq 0$ and any n -forest $(\mathfrak{U}, \mathfrak{T})$ we have for all $f, g : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ and $|g| \leq \mathbf{1}_G$*

$$\begin{aligned} & \left| \int \overline{g(x)} \sum_{\mathfrak{u} \in \mathfrak{U}} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}} f(x) d\mu(x) \right| \\ & \leq 2^{440a^3} 2^{-\frac{q-1}{q}n} \text{dens}_2 \left(\bigcup_{\mathfrak{u} \in \mathfrak{U}} \mathfrak{T}(\mathfrak{u}) \right)^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \end{aligned}$$

Theorem 1.1.1 is formulated at the level of generality for general kernels satisfying the mere Hölder regularity condition (1.1.12). On the other hand, the cancellative condition (1.1.9) is a testing condition against more regular, namely Lipschitz functions. To bridge the gap, we follow [Zor21] to observe a variant of (1.1.9) that we formulate in the following proposition proved in Section 8.

Define

$$\tau := \frac{1}{a}. \quad (2.0.38)$$

Define for any open ball B of radius R in X the L^∞ -normalized τ -Hölder norm by

$$\|\varphi\|_{C^\tau(B)} = \sup_{x \in B} |\varphi(x)| + R^\tau \sup_{x, y \in B, x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)^\tau}. \quad (2.0.39)$$

Proposition 2.0.5 (Holder van der Corput). *Let $z \in X$ and $R > 0$ and set $B = B(z, R)$. Let $\varphi : X \rightarrow \mathbb{C}$ be supported on B and satisfy $\|\varphi\|_{C^\tau(B(z, 2R))} < \infty$. Let $\vartheta, \theta \in \Theta$. Then*

$$\left| \int e(\vartheta(x) - \theta(x)) \varphi(x) dx \right| \leq 2^{7a} \mu(B) \|\varphi\|_{C^\tau(B(z, 2R))} (1 + d_B(\vartheta, \theta))^{-\frac{1}{2a^2 + a^3}}. \quad (2.0.40)$$

We further formulate a classical Vitali covering result and maximal function estimate that we need throughout several sections. This following proposition will typically be applied to the absolute value of a complex valued function and be proved in Section 9. By a ball B we mean a set $B(x, r)$ with $x \in X$ and $r > 0$ as defined in (1.1.3). For a finite collection \mathcal{B} of balls in X and $1 \leq p < \infty$ define the measurable function $M_{\mathcal{B}, p} u$ on X by

$$M_{\mathcal{B}, p} u(x) := \left(\sup_{B \in \mathcal{B}} \frac{\mathbf{1}_B(x)}{\mu(B)} \int_B |u(y)|^p d\mu(y) \right)^{\frac{1}{p}}. \quad (2.0.41)$$

Define further $M_{\mathcal{B}} := M_{\mathcal{B}, 1}$.

Proposition 2.0.6 (Hardy–Littlewood). *Let \mathcal{B} be a finite collection of balls in X . If for some $\lambda > 0$ and some measurable function $u : X \rightarrow [0, \infty)$ we have*

$$\int_B u(x) d\mu(x) \geq \lambda \mu(B) \quad (2.0.42)$$

for each $B \in \mathcal{B}$, then

$$\lambda \mu\left(\bigcup \mathcal{B}\right) \leq 2^{2a} \int_X u(x) d\mu(x). \quad (2.0.43)$$

For every measurable function v and $1 \leq p_1 < p_2$ we have

$$\|M_{\mathcal{B}, p_1} v\|_{p_2} \leq 2^{2a} \frac{p_2}{p_2 - p_1} \|v\|_{p_2}. \quad (2.0.44)$$

Moreover, given any measurable bounded function $w : X \rightarrow \mathbb{C}$ there exists a measurable function $Mw : X \rightarrow [0, \infty)$ such that the following (2.0.45) and (2.0.46) hold. For each ball $B \subset X$ and each $x \in B$

$$\frac{1}{\mu(B)} \int_B |w(y)| d\mu(y) \leq Mw(x) \quad (2.0.45)$$

and for all $1 \leq p_1 < p_2 \leq \infty$

$$\|M(w^{p_1})^{\frac{1}{p_1}}\|_{p_2} \leq 2^{4a} \frac{p_2}{p_2 - p_1} \|w\|_{p_2}. \quad (2.0.46)$$

This completes the overview of the proof of Theorem 1.1.1.

2.1. Auxiliary lemmas. We close this section by recording some auxiliary lemmas about the objects defined in Section 2, which will be used in multiple sections to follow.

First, we record an estimate for the metrical entropy numbers of balls in the space Θ equipped with any of the metrics d_B , following from the doubling property (1.1.8).

Lemma 2.1.1 (ball metric entropy). *Let $B' \subset X$ be a ball. Let $r > 0$, $\vartheta \in \Theta$ and $k \in \mathbb{N}$. Suppose that $\mathcal{Z} \subset B_{B'}(\vartheta, r2^k)$ satisfies that $\{B_{B'}(z, r) \mid z \in \mathcal{Z}\}$ is a collection of pairwise disjoint sets. Then*

$$|\mathcal{Z}| \leq 2^{ka}.$$

Proof. By applying property (1.1.8) k times, we obtain a collection $\mathcal{Z}' \subset \Theta$ with $|\mathcal{Z}'| = 2^{ka}$ and

$$B_{B'}(\vartheta, r2^k) \subset \bigcup_{z' \in \mathcal{Z}'} B_{B'}(z', \frac{r}{2}).$$

Then each $z \in \mathcal{Z}$ is contained in one of the balls $B(z', \frac{r}{2})$, but by the separation assumption no such ball contains more than one element of \mathcal{Z} . Thus $|\mathcal{Z}| \leq |\mathcal{Z}'| = 2^{ka}$. \square

The next lemma concerns monotonicity of the metrics $d_{B(c(I), \frac{1}{4}D^{s(I)})}$ with respect to inclusion of cubes I in a grid.

Lemma 2.1.2 (monotone cube metrics). *Let (\mathcal{D}, c, s) be a grid structure. Denote for cubes $I \in \mathcal{D}$*

$$I^\circ := B(c(I), \frac{1}{4}D^{s(I)}).$$

Let $I, J \in \mathcal{D}$ with $I \subset J$. Then for all $\vartheta, \theta \in \Theta$ we have

$$d_{I^\circ}(\vartheta, \theta) \leq d_{J^\circ}(\vartheta, \theta),$$

and if $I \neq J$ then we have

$$d_{I^\circ}(\vartheta, \theta) \leq 2^{-95a} d_{J^\circ}(\vartheta, \theta).$$

Proof. If $s(I) \geq s(J)$ then (2.0.8) and the assumption $I \subset J$ imply $I = J$. Then the lemma holds by reflexivity.

If $s(J) \geq s(I) + 1$, then using the monotonicity property (1.1.6), (2.0.1) and (1.1.7), we get

$$d_{I^\circ}(\vartheta, \theta) \leq d_{B(c(I), 4D^{s(I)})}(\vartheta, \theta) \leq 2^{-100a} d_{B(c(I), 4D^{s(J)})}(\vartheta, \theta). \quad (2.1.1)$$

Using (2.0.10), together with the inclusion $I \subset J$, we obtain

$$c(I) \in I \subset J \subset B(c(J), 4D^{s(J)})$$

and consequently by the triangle inequality

$$B(c(I), 4D^{s(J)}) \subset B(c(J), 8D^{s(J)}).$$

Using this together with the monotonicity property (1.1.6) and (1.1.5) in (2.1.1), we obtain

$$\begin{aligned} d_{I^s}(\vartheta, \theta) &\leq 2^{-100a} d_{B(c(J), 8D^s(J))}(\vartheta, \theta) \\ &\leq 2^{-100a+5a} d_{B(c(J), \frac{1}{4}D^s(J))}(\vartheta, \theta) \\ &= 2^{-95a} d_{J^s}(\vartheta, \theta). \end{aligned}$$

This proves the second inequality claimed in the Lemma, from which the first follows since $a \geq 4$ and hence $2^{-95a} \leq 1$. \square

We also record the following basic estimates for the kernels K_s .

Lemma 2.1.3 (kernel summand). *Let $-S \leq s \leq S$ and $x, y, y' \in X$. If $K_s(x, y) \neq 0$, then we have*

$$\frac{1}{4}D^{s-1} \leq \rho(x, y) \leq \frac{1}{2}D^s. \quad (2.1.2)$$

We have

$$|K_s(x, y)| \leq \frac{2^{102a^3}}{\mu(B(x, D^s))} \quad (2.1.3)$$

and

$$|K_s(x, y) - K_s(x, y')| \leq \frac{2^{127a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}}. \quad (2.1.4)$$

Proof. By Definition (2.0.5), the function K_s is the product of K with a function which is supported in the set of all x, y satisfying (2.1.2). This proves (2.1.2).

Using (1.1.11) and the lower bound in (2.1.2) we obtain

$$|K_s(x, y)| \leq |K(x, y)| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4}D^{s-1}))} \quad (2.1.5)$$

Using $D = 2^{100a^2}$ and the doubling property (1.1.2) $2 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{2a+101a^3}}{\mu(B(x, D^s))}. \quad (2.1.6)$$

Using $a \geq 4$ proves (2.1.3).

To prove (2.1.4) when $2\rho(y, y') > \rho(x, y)$, use the lower bound in (2.1.2) and the inequality $2\rho(y, y') > \frac{1}{4}D^{s-1}$. Then (2.1.4) follows from the triangle inequality, (2.1.3) and $a \geq 4$.

If $2\rho(y, y') \leq \rho(x, y)$, we rewrite $|K_s(x, y) - K_s(x, y')|$ as

$$|(K(x, y) - K(x, y'))\psi(D^{-s}\rho(x, y)) + K(x, y)(\psi(D^{-s}\rho(x, y)) - \psi(D^{-s}\rho(x, y')))|. \quad (2.1.7)$$

An upper bound for $|K(x, y) - K(x, y')|$ is obtained similarly to the proof of (2.1.3), using (1.1.12) and the lower bound in (2.1.2)

$$|K(x, y) - K(x, y')| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4}D^{s-1}))} \left(\frac{\rho(y, y')}{\frac{1}{4}D^{s-1}} \right)^{\frac{1}{a}}. \quad (2.1.8)$$

As above, this is estimated by

$$\leq \frac{4D2^{2a+101a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}} = \frac{2^{2+2a+100a^2+101a^3}}{\mu(B(x, D^s))} \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}}. \quad (2.1.9)$$

We have the trivial bound $|\psi(D^{-s}\rho(x, y))| \leq 1$, and (2.1.6) provides a bound for $|K(x, y)|$. Finally, we show that

$$|\psi(D^{-s}\rho(x, y)) - \psi(D^{-s}\rho(x, y'))| \leq 4D \left(\frac{\rho(y, y')}{D^s} \right)^{\frac{1}{a}} \quad (2.1.10)$$

by considering separately the cases $\rho(y, y')/D^s \geq 1$ and $\rho(y, y')/D^s < 1$. In the former case, the inequality is trivial; in the latter case, it follows from the fact that ψ is Lipschitz with constant $4D$.

Combining the above bounds and using $a \geq 4$ proves (2.1.4) when $2\rho(y, y') \leq \rho(x, y)$. \square

3. PROOF OF METRIC SPACE CARLESON

In this section we prove Theorem 1.1.1 and Theorem 1.1.2.

Let (X, ρ, μ, a) be a doubling metric measure space and Θ a cancellative, compatible collection of functions on X . Let K be a one-sided Calderon-Zygmund kernel.

We begin by proving some continuity properties of the integrand in (1.1.14).

Lemma 3.0.1 (int continuous). *Let f be a measurable function with $|f| \leq 1$. Then the function*

$$G : X \times \Theta \times (0, \infty) \times (0, \infty) \rightarrow \mathbb{C}$$

$$G(x, \vartheta, R_1, R_2) := \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) \, d\mu(y)$$

is continuous in ϑ for fixed x, R_1, R_2 . It is right-continuous in R_1 for fixed x, ϑ, R_2 and left-continuous in R_2 for fixed x, ϑ, R_1 . Finally, it is measurable in x and bounded for fixed ϑ, R_1, R_2 .

Proof. Measurability in x follows from joint measurability of

$$K(x, y) \mathbf{1}_{B(x, R_2) \setminus B(x, R_1)}(y)$$

in x and y and (part of the proof of) Fubini's theorem.

(Partial) continuity in R_1 and R_2 is also a standard lemma in measure theory. It follows for example by splitting the integrand as $F_1 - F_{-1} + iF_i - iF_{-i}$ for positive, disjointly supported functions F_{\pm} and applying the monotone convergence theorem to each summand.

For continuity in ϑ note that

$$\begin{aligned} & |G(x, \vartheta, R_1, R_2) - G(x, \vartheta', R_1, R_2)| \\ &= \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) (e(\vartheta(y)) - e(\vartheta'(y))) \, d\mu(y) \right| \\ &\leq \int_{R_1 < \rho(x, y) < R_2} |K(x, y)| |f(y)| |e(\vartheta(y)) - e(\vartheta'(y)) - e(\vartheta(o)) + e(\vartheta'(o)) - 1| \, d\mu(y). \end{aligned}$$

By 1-Lipschitz continuity of e^{ix} , the property (1.1.4) of the metrics d , the kernel upper bound (1.1.11) and $|f| \leq 1$ this is

$$\leq \mu(B(x, R_2)) \sup_{R_1 < \rho(x, y) < R_2} \frac{2^{a^3}}{V(x, y)} d_{B(x, \rho(o, x) + R_2)}(\vartheta, \vartheta').$$

If $R_1 < \rho(x, y) < R_2$ then there exists n with $B(x, R_2) \subset B(x, 2^n \rho(x, y))$ and $2^n \leq 2R_2/R_1$. Hence, by the doubling property (1.1.2),

$$V(x, y) = \mu(B(x, \rho(x, y))) \geq 2^{-an} \mu(B(x, R_2)) \geq (2R_2/R_1)^{-a} \mu(B(x, R_2)).$$

Hence

$$|G(x, \vartheta, R_1, R_2) - G(x, \vartheta', R_1, R_2)| \leq 2^{a^3} \left(\frac{2R_2}{R_1} \right)^a d_{B(x, \rho(o, x) + R_2)}(\vartheta, \vartheta').$$

Since the topology on Θ is the one induced by any of the metrics d_B , continuity follows.

Finally, for boundedness as a function of x note that we also have by a similar computation using $|e(\vartheta)| = 1$

$$|G(x, \vartheta, R_1, R_2)| \leq 2^{a^3} \left(\frac{2R_2}{R_1} \right)^a.$$

□

We now prove Theorem 1.1.1 using Theorem 1.1.2.

Proof of Theorem 1.1.1. Let Borel sets F, G in X be given. Let a Borel function $f : X \rightarrow \mathbb{C}$ with $f \leq \mathbf{1}_F$ be given.

Let $\Theta' \subset \Theta$ be a countable dense set. By Lemma 3.0.1 we have

$$\begin{aligned} & \sup_{\vartheta \in \Theta'} \sup_{0 < R_1 < R_2} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right| \\ &= \sup_{\vartheta \in \Theta'} \sup_{0 < R_1 < R_2, R_i \in \mathbb{Q}} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right|. \end{aligned}$$

Consider an enumeration of Θ' and let Θ_n be the finite set consisting of the first $n + 1$ functions in the enumeration. Then by the monotone convergence theorem

$$\begin{aligned} & \left| \int_G T f d\mu \right| \\ &= \int_G \sup_{\vartheta \in \Theta'} \sup_{0 < R_1 < R_2, R_i \in \mathbb{Q}} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right| d\mu(x) \\ &= \lim_{n \rightarrow \infty} \int_G \sup_{\vartheta \in \Theta_n} \sup_{0 < R_1 < R_2, R_i \in \mathbb{Q}} \left| \int_{R_1 < \rho(x, y) < R_2} K(x, y) f(y) e(\vartheta(y)) d\mu(y) \right| d\mu(x). \end{aligned}$$

For each n , let $Q_n(x)$ be the measurable function specifying the maximizer in the supremum in ϑ in the previous line. It is possible to construct such a measurable function of x , because the function inside the supremum is measurable in x , being a countable supremum of measurable functions, and because Θ_n is finite. (For example

one may fix some order on the finite set Θ_n and pick the smallest maximizer with respect to that order. This is a measurable choice function.) Then the previous display becomes

$$\begin{aligned} &\leq \lim_{n \rightarrow \infty} \int_G \sup_{0 < R_1 < R_2} \left| \int_{R_1 < \rho(x,y) < R_2} K(x,y) f(y) e(Q_n(x)(y)) \, d\mu(y) \right| \, d\mu(x). \\ &= \lim_{n \rightarrow \infty} \int_G T_{Q_n} f \, d\mu. \end{aligned}$$

It remains to verify the assumptions of Theorem 1.1.2, which when applied here completes the proof.

The assumptions of Theorem 1.1.1 and Theorem 1.1.2 are the same, with the exception of the assumption (1.1.20). The assumption (1.1.20) however is weaker than the assumption (1.1.15) of Theorem 1.1.1. Indeed, setting for fixed x, ϑ the outer radius R_2 in (1.1.13) to $\min\{R'_2, R_Q(\vartheta, x')\}$, where R'_2 is the outer radius in (1.1.18), shows that (1.1.18) is smaller than or equal to (1.1.13). Thus we can apply Theorem 1.1.2, which completes the proof. \square

We continue with the proof of Theorem 1.1.2, via a series of reductions to simpler lemmas.

Let a measurable function Q with finite range be given.

Proof of Theorem 1.1.2. Let Borel sets F, G in X with finite measure be given. Let a Borel function $f : X \rightarrow \mathbb{C}$ with $f \leq \mathbf{1}_F$ be given.

For each $0 < R_1, R_2, R$, we define $T_{R_1, R_2, R} f$ as in (3.0.2). By Lemma 3.0.1, $T_{R_1, R_2, R} f$ is measurable and bounded, and we clearly have for each $x \in X$

$$T_Q f(x) = \lim_{n \rightarrow \infty} \sup_{2^{-n} < R_1 < R_2 < 2^n} T_{R_1, R_2, 2^n} f(x).$$

For each x and all f , the functions $\sup_{2^{-n} < R_1 < R_2 < 2^n} T_{R_1, R_2, 2^n} f(x)$ are measurable by Lemma 3.0.1 and form an increasing sequence in n . By the monotone convergence theorem, the claimed estimate (1.1.21) then follows from Lemma 3.0.2. \square

Lemma 3.0.2 (R truncation).

Let F, G be Borel sets in X . Let $f : X \rightarrow \mathbb{C}$ be a Borel function with $|f| \leq \mathbf{1}_F$. Then for all $R \in 2^{\mathbb{N}}$ we have

$$\int \mathbf{1}_G \sup_{1/R < R_1 < R_2 < R} |T_{R_1, R_2, R} f(x)| \, d\mu(x) \leq \frac{2^{443a^3}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \quad (3.0.1)$$

where

$$T_{R_1, R_2, R} f(x) = \mathbf{1}_{B(o, R)}(x) \int_{R_1 < \rho(x,y) < R_2} K(x,y) f(y) e(Q(x)(y)) \, d\mu(y). \quad (3.0.2)$$

Proof. Let F, G, f as in the lemma be given. Let $R \in 2^{\mathbb{N}}$ be given. By replacing G with $G \cap B(o, R)$ if necessary, a replacement that does not change the conclusion of the Lemma 3.0.2, it suffices to show (3.0.1) under the assumption that G is contained in $B(o, R)$ and thus bounded. We make this assumption. For every $x \in G$ and $R_2 < R$, the domain of integration in (3.0.2) is contained in $B(o, 2R)$. By

replacing F with $F \cap B(o, 2R)$ if necessary, and correspondingly restricting f to $B(o, 2R)$, it suffices to show (3.0.1) under the assumption that F is contained in $B(o, 2R)$ and thus bounded. We make this assumption.

Using the definition (2.0.5) of K_s and the partition of unity (2.0.4), we observe that for fixed $R_1 < R_2$ we have

$$K(x, y)\mathbf{1}_{R_1 < \rho(x, y) < R_2} = \sum_{s=s_1-2}^{s_2+2} K_s(x, y)\mathbf{1}_{R_1 < \rho(x, y) < R_2}, \quad (3.0.3)$$

where $s_1 = \lceil \log_D 2R_1 \rceil + 3$ and $s_2 = \lceil \log_D 4R_2 \rceil - 2$. To obtain the identity (3.0.3), we have used that on the set where $R_1 < \rho(x, y) < R_2$ the kernels K_s vanish unless s is inside the interval of summation in (3.0.3). Similarly, we observe

$$\sum_{s=s_1}^{s_2} K_s(x, y)\mathbf{1}_{R_1 < \rho(x, y) < R_2} = \sum_{s=s_1}^{s_2} K_s(x, y), \quad (3.0.4)$$

because on the support of K_s with $s_1 \leq s \leq s_2$ we have necessarily $R_1 < \rho(x, y) < R_2$. We thus express (3.0.2) as the sum of

$$T_{s_1, s_2} f(x) := \sum_{s_1 \leq s \leq s_2} \int K_s(x, y) f(y) e(Q(x)(y)) d\mu(y) \quad (3.0.5)$$

and

$$\sum_{s=s_1-2, s_1-1, s_2+1, s_2+2} \int_{R_1 < \rho(x, y) < R_2} K_s(x, y) f(y) e(Q(x)(y)) d\mu(y), \quad (3.0.6)$$

We apply the triangle inequality and estimate (3.0.1) separately with $T_{R_1, R_2, R}$ replaced by (3.0.5) and by (3.0.6). To handle the case (3.0.5), we employ Lemma 3.0.3. Here, we utilize the fact that if $\frac{1}{R} \leq R_1 \leq R_2 \leq R$, then s_1 and s_2 as in (3.0.5) are in an interval $[-S, S]$ for some sufficiently large S depending on R . To handle the case (3.0.6), we use the triangle inequality and the properties (2.1.2), (2.1.3) of K_s and $|f| \leq \mathbf{1}_F$ to obtain for arbitrary s the inequality

$$\begin{aligned} & \left| \int_{R_1 < \rho(x, y) < R_2} K_s(x, y) f(y) e(Q(x)(y)) d\mu(y) \right| \\ & \leq \frac{2^{102a^3}}{\mu(B(x, D^s))} \int_{B(x, D^s)} \mathbf{1}_F(y) d\mu(y) \leq 2^{102a^3} M \mathbf{1}_F(x), \end{aligned} \quad (3.0.7)$$

where $M \mathbf{1}_F$ is as defined in Proposition 2.0.6. Now the left-hand side of (3.0.1), with $T_{R_1, R_2, R}$ replaced by (3.0.6), can be estimated using Hölder's inequality and Proposition 2.0.6 by

$$2^{102a^3+2} \int \mathbf{1}_G(x) M \mathbf{1}_F(x) d\mu(x) \leq \frac{2^{102a^3+4a+3} q}{q-1} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}.$$

Summing the contributions from (3.0.5) and (3.0.6) completes the proof. \square

Lemma 3.0.3 (S truncation). *Let F, G be bounded Borel sets in X . Let $f : X \rightarrow \mathbb{C}$ be a Borel function with $|f| \leq 1_F$. Then for all $S \in \mathbb{Z}$ we have*

$$\int \mathbf{1}_G(x) \sup_{-S \leq s_1 \leq s_2 \leq S} |T_{s_1, s_2} f(x)| d\mu(x) \leq \frac{2^{442a^3+2}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \quad (3.0.8)$$

where

$$T_{s_1, s_2} f(x) = \sum_{s_1 \leq s \leq s_2} \int_X K_s(x, y) f(y) e(Q(x)(y)) d\mu(y). \quad (3.0.9)$$

Proof of Lemma 3.0.3. We reduce Lemma 3.0.3 to Lemma 3.0.4. For each x , let $\sigma_1(x)$ be the minimal element $s' \in [-S, S]$ such that

$$\max_{s' \leq s_2 \leq S} |T_{s', s_2} f(x)| = \max_{-S \leq s_1 \leq s_2 \leq S} |T_{s_1, s_2} f(x)| =: T_{1, x}.$$

Similarly, let $\sigma_2(x)$ be the minimal element $s'' \in [-S, S]$ such that

$$|T_{\sigma_2(x), s''} f(x)| = T_{1, x}.$$

With these choices, and noting that with the definition of $T_{2, \sigma_1, \sigma_2}$ from (3.0.11)

$$T_{\sigma_1(x), \sigma_2(x)} f(x) = T_{2, \sigma_1, \sigma_2} f(x),$$

we conclude that the left-hand side of (3.0.8) and (3.0.10) are equal. Therefore, Lemma 3.0.3 follows from Lemma 3.0.4. \square

Lemma 3.0.4 (linearized truncation). *Let $\sigma_1, \sigma_2 : X \rightarrow \mathbb{Z}$ be measurable functions with finite range and $\sigma_1 \leq \sigma_2$. Then we have*

$$\int \mathbf{1}_G(x) |T_{2, \sigma_1, \sigma_2} f(x)| d\mu(x) \leq \frac{2^{442a^3+2}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}, \quad (3.0.10)$$

with

$$T_{2, \sigma_1, \sigma_2} f(x) = \sum_{\sigma_1(x) \leq s \leq \sigma_2(x)} \int_X K_s(x, y) f(y) e(Q(x)(y)) d\mu(y). \quad (3.0.11)$$

Proof of Lemma 3.0.4. Fix σ_1, σ_2 and Q as in the lemma. Applying Proposition 2.0.1 recursively, we obtain a sequence of sets G_n with $G_0 = G$ and, for each $n \geq 0$, $G_{n+1} \subset G_n$, $\mu(G_n) \leq 2^{-n} \mu(G)$ and

$$\begin{aligned} & \int \mathbf{1}_{G_n \setminus G_{n+1}}(x) |T_{2, \sigma_1, \sigma_2} f(x)| d\mu(x) \\ & \leq \frac{2^{442a^3}}{(q-1)^5} \mu(G_n)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \end{aligned} \quad (3.0.12)$$

Adding the first n of these inequalities, we obtain by bounding a geometric series

$$\int \mathbf{1}_{G \setminus G_n}(x) |T_{2, \sigma_1, \sigma_2} f(x)| d\mu(x) \leq \frac{2^{442a^3+2}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (3.0.13)$$

As the integrand is non-negative and non-decreasing in n , we obtain by the monotone convergence theorem

$$\int \mathbf{1}_G(x) |T_{2, \sigma_1, \sigma_2} f(x)| d\mu(x) \leq \frac{2^{442a^3+2}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (3.0.14)$$

This completes the proof of Lemma 3.0.4 and thus Theorem 1.1.1. \square

4. PROOF OF FINITARY CARLESON

To prove Proposition 2.0.1, we already fixed in Section 2 measurable functions σ_1, σ_2, Q and Borel sets F, G . We have also defined S to be the smallest integer such that the ranges of σ_1 and σ_2 are contained in $[-S, S]$ and F and G are contained in the ball $B(o, \frac{1}{4}D^S)$.

The proof of the next lemma is done in Section 4.1, following the construction of dyadic cubes in [Chr90, §3].

Lemma 4.0.1 (grid existence). *There exists a grid structure (\mathcal{D}, c, s) .*

The next lemma, which we prove in Section 4.2, should be compared with the construction in [Zor21, Lemma 2.12].

Lemma 4.0.2 (tile structure). *For a given grid structure (\mathcal{D}, c, s) , there exists a tile structure $(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q}, c, s)$.*

Choose a grid structure (\mathcal{D}, c, s) with Lemma 4.0.1 and a tile structure for this grid structure $(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q}, c, s)$ with Lemma 4.0.2. Applying Proposition 2.0.2, we obtain a Borel set G' in X with $2\mu(G') \leq \mu(G)$ such that for all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have (2.0.22).

Lemma 4.0.3 (tile sum operator). *We have for all $x \in G \setminus G'$*

$$\sum_{\mathfrak{p} \in \mathfrak{P}} T_{\mathfrak{p}} f(x) = \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (4.0.1)$$

Proof. Fix $x \in G \setminus G'$. Sorting the tiles \mathfrak{p} on the left-hand-side of (4.0.1) by the value $s(\mathfrak{p}) \in [-S, S]$, it suffices to prove for every $-S \leq s \leq S$ that

$$\sum_{\mathfrak{p} \in \mathfrak{P}: s(\mathfrak{p})=s} T_{\mathfrak{p}} f(x) = 0 \quad (4.0.2)$$

if $s \notin [\sigma_1(x), \sigma_2(x)]$ and

$$\sum_{\mathfrak{p} \in \mathfrak{P}: s(\mathfrak{p})=s} T_{\mathfrak{p}} f(x) = \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y). \quad (4.0.3)$$

if $s \in [\sigma_1(x), \sigma_2(x)]$. If $s \notin [\sigma_1(x), \sigma_2(x)]$, then by definition of $E(\mathfrak{p})$ we have $x \notin E(\mathfrak{p})$ for any \mathfrak{p} with $s(\mathfrak{p}) = s$ and thus $T_{\mathfrak{p}} f(x) = 0$. This proves (4.0.2).

Now assume $s \in [\sigma_1(x), \sigma_2(x)]$. By (2.0.7), (2.0.9), (2.0.10), the fact that $c(I_0) = o$ and $G \subset B(o, \frac{1}{4}D^S)$, there is at least one $I \in \mathcal{D}$ with $s(I) = s$ and $x \in I$. By (2.0.8), this I is unique. By (2.0.13), there is precisely one $\mathfrak{p} \in \mathfrak{P}(I)$ such that $Q(x) \in \Omega(\mathfrak{p})$. Hence there is precisely one $\mathfrak{p} \in \mathfrak{P}$ with $s(\mathfrak{p}) = s$ such that $x \in E(\mathfrak{p})$. For this \mathfrak{p} , the value $T_{\mathfrak{p}} f(x)$ by its definition in (2.0.21) equals the right-hand side of (4.0.3). This proves the lemma. \square

We use this to prove Proposition 2.0.1.

Proof of Proposition 2.0.1. We now estimate with Lemma 4.0.3 and Proposition 2.0.2

$$\int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y)) d\mu(y) \right| d\mu(x) \quad (4.0.4)$$

$$= \int_{G \setminus G'} \left| \sum_{s=\sigma_1(x)}^{\sigma_2(x)} \int K_s(x, y) f(y) e(Q(x)(y) - Q(x)(x)) d\mu(y) \right| d\mu(x) \quad (4.0.5)$$

$$= \int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{P}} T_{\mathfrak{p}} f(x) \right| d\mu(x) \leq \frac{2^{442a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (4.0.6)$$

This proves (2.0.6) for the chosen set G' and arbitrary f and thus completes the proof of Proposition 2.0.1. \square

4.1. Proof of Grid Existence Lemma. We begin with the construction of the centers of the dyadic cubes.

Lemma 4.1.1 (counting balls). *Let $-S \leq k \leq S$. Consider $Y \subset X$ such that for any $y \in Y$, we have*

$$y \in B(o, 4D^S - D^k), \quad (4.1.1)$$

furthermore, for any $y' \in Y$ with $y \neq y'$, we have

$$B(y, D^k) \cap B(y', D^k) = \emptyset. \quad (4.1.2)$$

Then the cardinality of Y is bounded by

$$|Y| \leq 2^{3a+200Sa^3}. \quad (4.1.3)$$

Proof. Let k and Y be given. By applying the doubling property (1.1.2) inductively, we have for each integer $j \geq 0$

$$\mu(B(y, 2^j D^k)) \leq 2^{aj} \mu(B(y, D^k)). \quad (4.1.4)$$

Since X is the union of the balls $B(y, 2^j D^k)$ and μ is not zero, at least one of the balls $B(y, 2^j D^k)$ has positive measure, thus $B(y, D^k)$ has positive measure.

Applying (4.1.4) for $j' = \ln_2(8D^{2S}) = 3 + 2S \cdot 100a^2$ by (2.0.1), using $-S \leq k \leq S$, $y \in B(o, 4D^S)$, and the triangle inequality, we have

$$B(o, 4D^S) \subset B(y, 8D^S) \subset B(y, 2^{j'} D^k). \quad (4.1.5)$$

Using the disjointedness of the balls in (4.1.2), (4.1.1), and the triangle inequality for ρ , we obtain

$$|Y| \mu(B(o, 4D^S)) \leq 2^{j'a} \sum_{y \in Y} \mu(B(y, D^k)) \quad (4.1.6)$$

$$\leq 2^{j'a} \mu\left(\bigcup_{y \in Y} B(y, D^k)\right) \leq 2^{j'a} \mu(o, 4D^S). \quad (4.1.7)$$

As $\mu(o, 4D^S)$ is not zero, the lemma follows. \square

For each $-S \leq k \leq S$, let Y_k be a set of maximal cardinality in X such that $Y = Y_k$ satisfies the properties (4.1.1) and (4.1.2) and such that $o \in Y_k$. By the upper bound of Lemma 4.1.1, such a set exists.

For each $-S \leq k \leq S$, choose an enumeration of the points in the finite set Y_k and thus a total order $<$ on Y_k .

Lemma 4.1.2 (cover big ball). *For each $-S \leq k \leq S$, the ball $B(o, 4D^S - D^k)$ is contained in the union of the balls $B(y, 2D^k)$ with $y \in Y_k$.*

Proof. Let x be any point of $B(o, 4D^S - D^k)$. By maximality of $|Y_k|$, the ball $B(x, D^k)$ intersects one of the balls $B(y, D^k)$ with $y \in Y_k$. By the triangle inequality, $x \in B(y, 2D^k)$. \square

Define the set

$$\mathcal{C} := \{(y, k) : -S \leq k \leq S, y \in Y_k\} \quad (4.1.8)$$

We totally order the set \mathcal{C} lexicographically by setting $(y, k) < (y', k')$ if $k < k'$ or both $k = k'$ and $y < y'$. In what follows, we define recursively in the sense of this order a function

$$(I_1, I_2, I_3) : \mathcal{C} \rightarrow \mathcal{P}(X) \times \mathcal{P}(X) \times \mathcal{P}(X). \quad (4.1.9)$$

Assume the sets $I_j(y', k')$ have already been defined for $j = 1, 2, 3$ if $k' < k$ and if $k = k'$ and $y' < y$.

If $k = -S$, define for $j \in \{1, 2\}$ the set $I_j(y, k)$ to be $B(y, jD^{-S})$. If $-S < k$, define for $j \in \{1, 2\}$ and $y \in Y_k$ the set $I_j(y, k)$ to be

$$\bigcup \{I_3(y', k-1) : y' \in Y_{k-1} \cap B(y, jD^k)\}. \quad (4.1.10)$$

Define for $-S \leq k \leq S$ and $y \in Y_k$

$$I_3(y, k) := I_1(y, k) \cup \left[I_2(y, k) \setminus \left[X_k \cup \bigcup \{I_3(y', k) : y' \in Y_k, y' < y\} \right] \right] \quad (4.1.11)$$

with

$$X_k := \bigcup \{I_1(y', k) : y' \in Y_k\}. \quad (4.1.12)$$

Lemma 4.1.3 (basic grid structure). *For each $-S \leq k \leq S$ and $1 \leq j \leq 3$ the following holds.*

If $j \neq 2$ and for some $x \in X$ and $y_1, y_2 \in Y_k$ we have

$$x \in I_j(y_1, k) \cap I_j(y_2, k), \quad (4.1.13)$$

then $y_1 = y_2$.

If $j \neq 1$, then

$$B(o, 4D^S - 2D^k) \subset \bigcup_{y \in Y_k} I_j(y, k). \quad (4.1.14)$$

We have for each $y \in Y_k$,

$$B(y, \frac{1}{2}D^k) \subset I_3(y, k) \subset B(y, 4D^k). \quad (4.1.15)$$

Proof. We prove these statements simultaneously by induction on the ordered set of pairs (y, k) . Let $-S \leq k \leq S$.

We first consider (4.1.13) for $j = 1$. If $k = -S$, disjointedness of the sets $I_1(y, -S)$ follows by definition of I_1 and Y_k . If $k > -S$, assume x is in $I_1(y_m, k)$ for $m = 1, 2$. Then, for $m = 1, 2$, there is $z_m \in Y_{k-1} \cap B(y_m, D^k)$ with $x \in I_3(z_m, k-1)$. Using (4.1.13) inductively for $j = 3$, we conclude $z_1 = z_2$. This implies that the balls $B(y_1, D^k)$ and $B(y_2, D^k)$ intersect. By construction of Y_k , this implies $y_1 = y_2$. This proves (4.1.13) for $j = 1$.

We next consider (4.1.13) for $j = 3$. Assume x is in $I_3(y_m, k)$ for $m = 1, 2$ and $y_m \in Y_k$. If x is in X_k , then by definition (4.1.11), $x \in I_1(y_m, k)$ for $m = 1, 2$. As we have already shown (4.1.13) for $j = 1$, we conclude $y_1 = y_2$. This completes the proof in case $x \in X_k$, and we may assume x is not in X_k . By definition (4.1.11), x is not in $I_3(z, k)$ for any z with $z < y_1$ or $z < y_2$. Hence, neither $y_1 < y_2$ nor $y_2 < y_1$, and by totality of the order of Y_k , we have $y_1 = y_2$. This completes the proof of (4.1.13) for $j = 3$.

We show (4.1.14) for $j = 2$. In case $k = -S$, this follows from Lemma 4.1.2. Assume $k > -S$. Let x be a point of $B(o, 4D^S - 2D^k)$. By induction, there is $y' \in Y_{k-1}$ such that $x \in I_3(y', k-1)$. Using the inductive statement (4.1.15), we obtain $x \in B(y', 4D^{k-1})$. As $D > 4$, by applying the triangle inequality with the points, o , x , and y' , we obtain that $y' \in B(o, 4D^S - D^k)$. By Lemma 4.1.2, y' is in $B(y, 2D^k)$ for some $y \in Y_k$. It follows that $x \in I_2(y, k)$. This proves (4.1.14) for $j = 2$.

We show (4.1.14) for $j = 3$. Let $x \in B(o, 4D^S - 2D^k)$. In case $x \in X_k$, then by definition of X_k we have $x \in I_1(y, k)$ for some $y \in Y_k$ and thus $x \in I_3(y, k)$. We may thus assume $x \notin X_k$. As we have already seen (4.1.14) for $j = 2$, there is $y \in Y_k$ such that $x \in I_2(y, k)$. We may assume this y is minimal with respect to the order in Y_k . Then $x \in I_3(y, k)$. This proves (4.1.14) for $j = 3$.

Next, we show the first inclusion in (4.1.15). Let $x \in B(y, \frac{1}{2}D^k)$. As $I_1(y, k) \subset I_3(y, k)$, it suffices to show $x \in I_1(y, k)$. If $k = -S$, this follows immediately from the assumption on x and the definition of I_1 . Assume $k > -S$. By the inductive statement (4.1.14) and $D > 4$, there is a $y' \in Y_{k-1}$ such that $x \in I_3(y', k-1)$. By the inductive statement (4.1.15), we conclude $x \in B(y', 4D^{k-1})$. By the triangle inequality with points x , y , y' , and $D \geq 8$, we have $y' \in B(y, D^k)$. It follows by definition (4.1.10) that $I_3(y', k-1) \subset I_1(y, k)$, and thus $x \in I_3(y, k)$. This proves the first inclusion in (4.1.15).

We show the second inclusion in (4.1.15). Let $x \in I_3(y, k)$. As $I_1(y, k) \subset I_2(y, k)$ directly from the definition (4.1.10), it follows by definition (4.1.11) that $x \in I_2(y, k)$. By definition (4.1.10), there is $y' \in Y_{k-1} \cap B(y, 2D^k)$ with $x \in I_3(y', k-1)$. By induction, $x \in B(y', 4D^{k-1})$. By the triangle inequality applied to the points x, y', y and $D > 4$, we conclude $x \in B(y, 4D^k)$. This shows the second inclusion in (4.1.15) and completes the proof of the lemma. \square

Lemma 4.1.4 (cover by cubes). *Let $-S \leq l \leq k \leq S$ and $y \in Y_k$. We have*

$$I_3(y, k) \subset \bigcup_{y' \in Y_l} I_3(y', l). \quad (4.1.16)$$

Proof. Let $-S \leq l \leq k \leq S$ and $y \in Y_k$. If $l = k$, the inclusion (4.1.16) is true from the definition of set union. We may then assume inductively that $k > l$ and the statement of the lemma is true if k is replaced by $k - 1$. Let $x \in I_3(y, k)$. By definition (4.1.11), $x \in I_j(y, k)$ for some $j \in \{1, 2\}$. By (4.1.10), $x \in I_3(w, k - 1)$ for some $w \in Y_{k-1}$. We conclude (4.1.16) by induction. \square

Lemma 4.1.5 (dyadic property). *Let $-S \leq l \leq k \leq S$ and $y \in Y_k$ and $y' \in Y_l$ with $I_3(y', l) \cap I_3(y, k) \neq \emptyset$. Then*

$$I_3(y', l) \subset I_3(y, k). \quad (4.1.17)$$

Proof. Let l, k, y, y' be as in the lemma. Pick $x \in I_3(y', l) \cap I_3(y, k)$. Assume first $l = k$. By (4.1.13) of Lemma 4.1.3, we conclude $y' = y$, and thus (4.1.17). Now assume $l < k$. By induction, we may assume that the statement of the lemma is proven for $k - 1$ in place of k .

By Lemma 4.1.4, there is a $y'' \in Y_{k-1}$ such that $x \in I_3(y'', k - 1)$. By induction, we have $I_3(y', l) \subset I_3(y'', k - 1)$. It remains to prove

$$I_3(y'', k - 1) \subset I_3(y, k). \quad (4.1.18)$$

We make a case distinction and assume first $x \in X_k$. By Definition (4.1.11), we have $x \in I_1(y, k)$. By Definition (4.1.10), there is a $v \in Y_{k-1} \cap B(y, D^k)$ with $x \in I_3(v, k - 1)$. By (4.1.13) of Lemma 4.1.3, we have $v = y''$. By Definition (4.1.10), we then have $I_3(y'', k - 1) \subset I_1(y, k)$. Then (4.1.18) follows by Definition (4.1.11) in the given case.

Assume now the case $x \notin X_k$. By (4.1.11), we have $x \in I_2(y, k)$. Moreover, for any $u < y$ in Y_k , we have $x \notin I_3(u, k)$. Let $u < y$. By transitivity of the order in Y_k , we conclude $x \notin I_2(u, k)$. By (4.1.10) and the disjointedness property of Lemma 4.1.3, we have $I_3(y'', k - 1) \cap I_2(u, k) = \emptyset$. Similarly, $I_3(y'', k - 1) \cap I_1(u, k) = \emptyset$. Hence $I_3(y'', k - 1) \cap I_3(u, k) = \emptyset$. As $u < y$ was arbitrary, we conclude with (4.1.11) the claim in the given case. This completes the proof of (4.1.18), and thus also (4.1.17). \square

For $-S \leq k' \leq k \leq S$ and $y' \in Y_{k'}$, $y \in Y_k$ write $(y', k' | y, k)$ if $I_3(y', k') \subset I_3(y, k)$ and

$$\inf_{x \in X \setminus I_3(y, k)} \rho(y', x) < 6D^{k'}. \quad (4.1.19)$$

Lemma 4.1.6 (transitive boundary). *Assume $-S \leq k'' < k' < k \leq S$ and $y'' \in Y_{k''}$, $y' \in Y_{k'}$, $y \in Y_k$. Assume there is $x \in X$ such that*

$$x \in I_3(y'', k'') \cap I_3(y', k') \cap I_3(y, k). \quad (4.1.20)$$

If $(y'', k'' | y, k)$, then also $(y'', k'' | y', k')$ and $(y', k' | y, k)$

Proof. As $x \in I_3(y'', k'') \cap I_3(y', k')$ and $k'' < k'$, we have by Lemma 4.1.5 that $I_3(y'', k'') \subset I_3(y', k')$. Similarly, $I_3(y', k') \subset I_3(y, k)$. Pick $x' \in X \setminus I_3(y, k)$ such that

$$\rho(y'', x') < 6D^{k''}, \quad (4.1.21)$$

which exists as $(y'', k'' | y, k)$. As $x' \in X \setminus I_3(y', k')$ as well, we conclude $(y'', k'' | y', k')$. By the triangle inequality, we have

$$\rho(y', x') \leq \rho(y', x) + \rho(x, y'') + \rho(y'', x') \quad (4.1.22)$$

Using the choice of x and (4.1.15) as well as (4.1.21), we estimate this by

$$< 4D^{k'} + 4D^{k''} + 6D^{k''} \leq 6D^{k'}, \quad (4.1.23)$$

where we have used $D > 5$ and $k'' < k'$. We conclude $(y', k'|y, k)$. \square

Lemma 4.1.7 (small boundary). *Let $K = 2^{4a+1}$. For each $-S + K \leq k \leq S$ and $y \in Y_k$ we have*

$$\sum_{z \in Y_{k-K}: (z, k-K|y, k)} \mu(I_3(z, k-K)) \leq \frac{1}{2} \mu(I_3(y, k)). \quad (4.1.24)$$

Proof. Let K be as in the lemma. Let $-S + K \leq k \leq S$ and $y \in Y_k$.

Pick k' so that $k - K \leq k' \leq k$. For each $y'' \in Y_{k-K}$ with $(y'', k - K|y, k)$, by Lemma 4.1.4 and Lemma 4.1.5, there is a unique $y' \in Y_{k'}$ such that

$$I_3(y'', k - K) \subset I_3(y', k') \subset I_3(y, k). \quad (4.1.25)$$

Using Lemma 4.1.6, $(y', k'|y, k)$.

We conclude using the disjointedness property of Lemma 4.1.3 that

$$\sum_{y'': (y'', k-K|y, k)} \mu(I_3(y'', k - K)) \leq \sum_{y': (y', k'|y, k)} \mu(I_3(y', k')). \quad (4.1.26)$$

Adding over $k - K < k' \leq k$, and using

$$\mu(I_3(y', k')) \leq 2^{4a} \mu(B(y', \frac{1}{4}D^{k'}))$$

from the doubling property (1.1.2) and (4.1.15) gives

$$K \sum_{y'': (y'', k-K|y, k)} \mu(I_3(y'', k - K)) \quad (4.1.27)$$

$$\leq 2^{4a} \sum_{k-K < k' \leq k} \left[\sum_{y': (y', k'|y, k)} \mu(B(y', \frac{1}{4}D^{k'})) \right] \quad (4.1.28)$$

Each ball $B(y', \frac{1}{4}D^{k'})$ occurring in (4.1.28) is contained in $I_3(y', k')$ by (4.1.15) and in turn contained in $I_3(y, k)$ by (4.1.25). Assume for the moment all these balls are pairwise disjoint. Then by additivity of the measure,

$$K \sum_{y'': (y'', k-K|y, k)} \mu(I_3(y'', k - K)) \leq 2^{4a} \mu(I_3(y, k)) \quad (4.1.29)$$

which by $K = 2^{4a+1}$ implies (4.1.24).

It thus remains to prove that the balls occurring in (4.1.28) are pairwise disjoint. Let (u, l) and (u', l') be two parameter pairs occurring in the sum of (4.1.28) and let $B(u, \frac{1}{4}D^l)$ and $B(u', \frac{1}{4}D^{l'})$ be the corresponding balls. If $l = l'$, then the balls are equal or disjoint by (4.1.15) and (4.1.13) of Lemma 4.1.3. Assume then without loss of generality that $l' < l$. Towards a contradiction, assume that

$$B(u, \frac{1}{4}D^l) \cap B(u', \frac{1}{4}D^{l'}) \neq \emptyset \quad (4.1.30)$$

As $(u', l'|y, k)$, there is a point x in $X \setminus I_3(y, k)$ with $\rho(x, u') < 6D^{l'}$. Using $D > 25$, we conclude from the triangle inequality and (4.1.30) that $x \in B(u, \frac{1}{2}D^l)$. However,

$B(u, \frac{1}{2}D^l) \subset I_3(u, l)$, and $I_3(u, l) \subset I_3(y, k)$, a contradiction to $x \notin I_3(y, k)$. This proves the lemma. \square

Lemma 4.1.8 (smaller boundary). *Let $K = 2^{4a+1}$ and let $n \geq 0$ be an integer. Then for each $-S + nK \leq k \leq S$ we have*

$$\sum_{y' \in Y_{k-nK}: (y', k-nK|y, k)} \mu(I_3(y', k-nK)) \leq 2^{-n} \mu(I_3(y, k)). \quad (4.1.31)$$

Proof. We prove this by induction on n . If $n = 0$, both sides of (4.1.31) are equal to $\mu(I_3(y, k))$ by (4.1.13). If $n = 1$, this follows from Lemma 4.1.7.

Assume $n > 1$ and (4.1.31) has been proven for $n - 1$. We write (4.1.31)

$$\sum_{y'' \in Y_{k-nK}: (y'', k-nK|y, k)} \mu(I_3(y'', k-nK)) \quad (4.1.32)$$

$$= \sum_{y' \in Y_{k-K}: (y', k-K|y, k)} \left[\sum_{y'' \in Y_{k-nK}: (y'', k-nK|y', k-K)} \mu(I_3(y'', k-nK)) \right] \quad (4.1.33)$$

Applying the induction hypothesis, this is bounded by

$$= \sum_{y' \in Y_{k-K}: (y', k-K|y, k)} 2^{1-n} \mu(I_3(y', k-K)) \quad (4.1.34)$$

Applying (4.1.24) gives (4.1.31), and proves the lemma. \square

Lemma 4.1.9 (boundary measure). *For each $-S \leq k \leq S$ and $y \in Y_k$ and $0 < t < 1$ with $tD^k \geq D^{-S}$ we have*

$$\mu(\{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\}) \leq 2t^\kappa \mu(I_3(y, k)). \quad (4.1.35)$$

Proof. Let $x \in I_3(y, k)$ with $\rho(x, X \setminus I_3(y, k)) \leq tD^k$. Let $K = 2^{4a+1}$ as in Lemma 4.1.8. Let n be the largest integer such that $D^{nK} \leq \frac{1}{t}$, so that $tD^k \leq D^{k-nK}$ and

$$D^{nK} > \frac{1}{tD^k}. \quad (4.1.36)$$

Let $k' = k - nK$, by the assumption $tD^k \geq D^{-S}$, we have $k' \geq -S$. By (4.1.16), there exists $y' \in Y_{k'}$ with $x \in I_3(y', k')$. By the squeezing property (4.1.15) and the assumption on x , we have

$$\rho(y', X \setminus I_3(y, k)) \leq \rho(x, y') + \rho(x, X \setminus I_3(y, k)) \leq 4D^{k'} + tD^k.$$

By the assumption on n and the definition of k' , this is

$$\leq 4D^{k'} + D^{k-nK} < 6D^{k'}.$$

Together with (4.1.17) thus $(y', k'|y, k)$. We have shown that

$$\begin{aligned} & \{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\} \\ & \subset \bigcup_{y' \in Y_{k-nK}: (y', k-nK|y, k)} I_3(y', k-nK). \end{aligned}$$

Using monotonicity and additivity of the measure and Lemma 4.1.8, we obtain

$$\mu(\{x \in I_3(y, k) : \rho(x, X \setminus I_3(y, k)) \leq tD^k\}) \leq 2^{-n} \mu(I_3(y, k)).$$

By (4.1.36) and the definition (2.0.1) of D , this is bounded by

$$2t^{1/(100a^2K)}\mu(I_3(y, k)),$$

which completes the proof by the definition (2.0.2) of κ . \square

Let $\tilde{\mathcal{D}}$ be the set of all $I_3(y, k)$ with $k \in [-S, S]$ and $y \in Y_k$. Define

$$s(I_3(y, k)) := k \tag{4.1.37}$$

$$c(I_3(y, k)) := y. \tag{4.1.38}$$

We define \mathcal{D} to be the set of all $I \in \tilde{\mathcal{D}}$ such that $I \subset I_3(o, S)$.

Proof of Lemma 4.0.1. We first show that $(\tilde{\mathcal{D}}, c, s)$ satisfies properties (2.0.7), (2.0.8), (2.0.10) and (2.0.11). Property (2.0.10) follows from (4.1.15), while (2.0.11) follows from Lemma 4.1.9.

Let $x \in B(o, D^S)$. We show properties (2.0.7) and (2.0.8) for $(\tilde{\mathcal{D}}, c, s)$ and x .

We first show (2.0.7) for $(\tilde{\mathcal{D}}, c, s)$ by contradiction. Then there is an I violating the conclusion of (2.0.7). Pick such $I = I_3(y, l)$ such that l is minimal. By assumption, we have $-S \leq k < l$; in particular $-S < l$. By definition, $I_3(y, l)$ is contained in $I_1(y, l) \cup I_2(y, l)$, which is contained in the union of $I_3(y', l-1)$ with $y' \in Y_{l-1}$. By minimality of l , each such $I_3(y', l-1)$ is contained in the union of all $I_3(z, k)$ with $z \in Y_k$. This proves (2.0.7).

We now show (2.0.8) for $(\tilde{\mathcal{D}}, c, s)$. Assume to get a contradiction that there are non-disjoint $I, J \in \tilde{\mathcal{D}}$ with $s(I) \leq s(J)$ and $I \not\subset J$. We may assume the existence of such I and J with minimal $s(J) - s(I)$. Let $k = s(I)$. Assume first $s(J) = k$. Let $I = I_3(y_1, k)$ and $J = I_3(y_2, k)$ with $y_1, y_2 \in Y_k$. If $y_1 = y_2$, then $I = J$, a contradiction to $I \not\subset J$. If $y_1 \neq y_2$, then $I \cap J = \emptyset$ by (4.1.13), a contradiction to the non-disjointness of I, J . Assume now $s(J) > k$. Choose $y \in I \cap J$. By property (2.0.7), there is $K \in \tilde{\mathcal{D}}$ with $s(K) = s(J) - 1$ and $y \in K$. By construction of J , and pairwise disjointness of all $I_3(w, s(J) - 1)$ that we have already seen, we have $K \subset J$. By minimality of $s(J)$, we have $I \subset K$. This proves $I \subset J$ and thus (2.0.8).

Now note that properties (2.0.8), (2.0.10) and (2.0.11) immediately carry over to (\mathcal{D}, c, s) by restriction. (2.0.9) is true for (\mathcal{D}, c, s) by definition, and (2.0.7) follows from (2.0.7) and (2.0.8) for $(\tilde{\mathcal{D}}, c, s)$. \square

4.2. Proof of Tile Structure Lemma. Choose a grid structure (\mathcal{D}, c, s) with Lemma 4.0.1. Let $I \in \mathcal{D}$. Suppose that

$$\mathcal{Z} \subset Q(X) \tag{4.2.1}$$

is such that for any $\vartheta, \theta \in \mathcal{Z}$ with $\vartheta \neq \theta$ we have

$$B_{I^\circ}(\vartheta, 0.3) \cap B_{I^\circ}(\theta, 0.3) \cap Q(X) = \emptyset. \tag{4.2.2}$$

Since $Q(X)$ is finite, there exists a set \mathcal{Z} satisfying both (4.2.1) and (4.2.2) of maximal cardinality among all such sets. We pick for each $I \in \mathcal{D}$ such a set $\mathcal{Z}(I)$.

Lemma 4.2.1 (frequency ball cover). *For each $I \in \mathcal{D}$, we have*

$$Q(X) \subset \bigcup_{z \in \mathcal{Z}(I)} B_{I^\circ}(z, 0.7). \tag{4.2.3}$$

Proof. Let $\theta \in \bigcup_{\vartheta \in Q(X)} B_{I^\circ}(\vartheta, 1)$. By maximality of $\mathcal{Z}(I)$, there must be a point $z \in \mathcal{Z}(I)$ such that $B_{I^\circ}(z, 0.3) \cap B_{I^\circ}(\theta, 0.3) \neq \emptyset$. Else, $\mathcal{Z}(I) \cup \{\theta\}$ would be a set of larger cardinality than $\mathcal{Z}(I)$ satisfying (4.2.1) and (4.2.2). Fix such z , and fix a point $z_1 \in B_{I^\circ}(z, 0.3) \cap B_{I^\circ}(\theta, 0.3)$. By the triangle inequality, we deduce that

$$d_{I^\circ}(z, \theta) \leq d_{I^\circ}(z, z_1) + d_{I^\circ}(\theta, z_1) < 0.3 + 0.3 = 0.6,$$

and hence $\theta \in B_{I^\circ}(z, 0.7)$. \square

We define

$$\begin{aligned} \mathfrak{P} &= \{(I, z) : I \in \mathcal{D}, z \in \mathcal{Z}(I)\}, \\ \mathcal{J}((I, z)) &= I \quad \text{and} \quad \mathcal{Q}((I, z)) = z. \end{aligned}$$

We further set

$$s(\mathfrak{p}) = s(\mathcal{J}(\mathfrak{p})), \quad c(\mathfrak{p}) = c(\mathcal{J}(\mathfrak{p})).$$

Then (2.0.18), (2.0.19) hold by definition.

It remains to construct the map Ω , and verify properties (2.0.13), (2.0.14) and (2.0.15). We first construct an auxiliary map Ω_1 . For each $I \in \mathcal{D}$, we pick an enumeration of the finite set $\mathcal{Z}(I)$

$$\mathcal{Z}(I) = \{z_1, \dots, z_M\}.$$

We define $\Omega_1 : \mathfrak{P} \mapsto \mathcal{P}(\Theta)$ as below. Set

$$\Omega_1((I, z_1)) = B_{I^\circ}(z_1, 0.7) \setminus \bigcup_{z \in \mathcal{Z}(I) \setminus \{z_1\}} B_{I^\circ}(z, 0.3)$$

and then define iteratively

$$\Omega_1((I, z_k)) = B_{I^\circ}(z_k, 0.7) \setminus \bigcup_{z \in \mathcal{Z}(I) \setminus \{z_k\}} B_{I^\circ}(z, 0.3) \setminus \bigcup_{i=1}^{k-1} \Omega_1((I, z_i)). \quad (4.2.4)$$

Lemma 4.2.2 (disjoint frequency cubes). *For each $I \in \mathcal{D}$, and $\mathfrak{p}_1, \mathfrak{p}_2 \in \mathfrak{P}(I)$, if*

$$\Omega_1(\mathfrak{p}_1) \cap \Omega_1(\mathfrak{p}_2) \neq \emptyset,$$

then $\mathfrak{p}_1 = \mathfrak{p}_2$.

Proof. By the definition of the map \mathcal{J} , we have

$$\mathfrak{P}(I) = \{(I, z) : z \in \mathcal{Z}(I)\}.$$

By (4.2.4), the set $\Omega_1((I, z_k))$ is disjoint from each $\Omega_1((I, z_i))$ with $i < k$. Thus the sets $\Omega_1(\mathfrak{p})$, $\mathfrak{p} \in \mathfrak{P}(I)$ are pairwise disjoint. \square

Lemma 4.2.3 (frequency cube cover). *For each $I \in \mathcal{D}$, it holds that*

$$\bigcup_{z \in \mathcal{Z}(I)} B_{I^\circ}(z, 0.7) \subset \bigcup_{\mathfrak{p} \in \mathfrak{P}(I)} \Omega_1(\mathfrak{p}). \quad (4.2.5)$$

For every $\mathfrak{p} \in \mathfrak{P}$, it holds that

$$B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.3) \subset \Omega_1(\mathfrak{p}) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 0.7). \quad (4.2.6)$$

Proof. For (4.2.6) let $\mathbf{p} = (I, z)$. The second inclusion in (4.2.6) then follows from (4.2.4) and the equality $B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.7) = B_{I^c}(z, 0.7)$, which is true by definition. For the first inclusion in (4.2.6) let $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.3)$. Let k be such that $z = z_k$ in the enumeration we chose above. It follows immediately from (4.2.4) and (4.2.2) that $\vartheta \notin \Omega_1((I, z_i))$ for all $i < k$. Thus, again from (4.2.4), we have $\vartheta \in \Omega_1((I, z_k))$.

To show (4.2.5) let $\vartheta \in \bigcup_{z \in \mathcal{Z}(I)} B_{I^c}(z, 0.7)$. If there exists $z \in \mathcal{Z}(I)$ with $\vartheta \in B_{I^c}(z, 0.3)$, then

$$z \in \Omega_1((I, z)) \subset \bigcup_{\mathbf{p} \in \mathfrak{P}(I)} \Omega_1(\mathbf{p})$$

by the first inclusion in (4.2.6).

Now suppose that there exists no $z \in \mathcal{Z}(I)$ with $\vartheta \in B_{I^c}(z, 0.3)$. Let k be minimal such that $\vartheta \in B_{I^c}(z_k, 0.7)$. Since $\Omega_1((I, z_i)) \subset B_{I^c}(z_i, 0.7)$ for each i by (4.2.4), we have that $\vartheta \notin \Omega_1((I, z_i))$ for all $i < k$. Hence $\vartheta \in \Omega_1((I, z_k))$, again by (4.2.4). \square

Now we are ready to define the function Ω . We define for all $\mathbf{p} \in \mathfrak{P}(I_0)$

$$\Omega(\mathbf{p}) = \Omega_1(\mathbf{p}). \quad (4.2.7)$$

For all other cubes $I \in \mathcal{D}$, $I \neq I_0$, there exists, by (2.0.8) and (2.0.9), $J \in \mathcal{D}$ with $I \subset J$ and $I \neq J$. On such I we define Ω by recursion. We can pick an inclusion minimal $J \in \mathcal{D}$ among the finitely many cubes such that $I \subset J$ and $I \neq J$. This J is unique: Suppose that J' is another inclusion minimal cube with $I \subset J'$ and $I \neq J'$. Without loss of generality, we have that $s(J) \leq s(J')$. By (2.0.8), it follows that $J \subset J'$. Since J' is minimal with respect to inclusion, it follows that $J = J'$. Then we define

$$\Omega(\mathbf{p}) = \bigcup_{z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})} \Omega((J, z)) \cup B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2). \quad (4.2.8)$$

We now verify that $(\mathfrak{P}, J, \Omega, \mathcal{Q}, c, s)$ forms a tile structure.

Proof of Lemma 4.0.2. First, we prove (2.0.15). If $I = I_0$, then (2.0.15) holds for all $\mathbf{p} \in \mathfrak{P}(I)$ by (4.2.7) and (4.2.6). Now suppose that I is not maximal in \mathcal{D} with respect to set inclusion. Then we may assume by induction that for all $J \in \mathcal{D}$ with $I \subset J$ and all $\mathbf{p}' \in \mathfrak{P}(J)$, (2.0.15) holds. Let J be the unique minimal cube in \mathcal{D} with $I \subsetneq J$.

Suppose that $\vartheta \in \Omega(\mathbf{p})$. If $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$, then since

$$B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 1),$$

we conclude that $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.7)$. If not, by (4.2.8), there exists $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ with $\vartheta \in \Omega(J, z)$. Using the triangle inequality and (4.2.6), we obtain

$$d_{I^c}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq d_{I^c}(\mathcal{Q}(\mathbf{p}), z) + d_{I^c}(z, \vartheta) \leq 0.7 + d_{I^c}(z, \vartheta).$$

By Lemma 2.1.2 and the induction hypothesis, this is estimated by

$$\leq 0.7 + 2^{-95a} d_{J^c}(z, \vartheta) \leq 0.7 + 2^{-95a} \cdot 1 < 1.$$

This shows the second inclusion in (2.0.15). The first inclusion is immediate from (4.2.8).

Next, we show (2.0.13). Let $I \in \mathcal{D}$.

If $I = I_0$, then disjointedness of the sets $\Omega(\mathbf{p})$ for $\mathbf{p} \in \mathfrak{P}(I)$ follows from the definition (4.2.7) and Lemma 4.2.2. To obtain the inclusion in (2.0.13) one combines the inclusions (4.2.3) and (4.2.5) of Lemma 4.2.3 with (4.2.7).

Now we turn to the case where there exists $J \in \mathcal{D}$ with $I \subset J$ and $I \neq J$. In this case we use induction: It suffices to show (2.0.13) under the assumption that it holds for all cubes $J \in \mathcal{D}$ with $I \subset J$. As shown before definition (4.2.8), we may choose the unique inclusion minimal such J . To show disjointedness of the sets $\Omega(\mathbf{p})$, $\mathbf{p} \in \mathfrak{P}(I)$ we pick two tiles $\mathbf{p}, \mathbf{p}' \in \mathfrak{P}(I)$ and $\vartheta \in \Omega(\mathbf{p}) \cap \Omega(\mathbf{p}')$. Then we are by (4.2.8) in one of the following four cases.

1. There exist $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ such that $\vartheta \in \Omega(J, z)$, and there exists $z' \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$ such that $\vartheta \in \Omega(J, z')$. By the induction hypothesis, that (2.0.13) holds for J , we must have $z = z'$. By Lemma 4.2.2, we must then have $\mathbf{p} = \mathbf{p}'$.

2. There exists $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p})$ such that $\vartheta \in \Omega(J, z)$, and $\vartheta \in B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 0.2)$. Using the triangle inequality, Lemma 2.1.2 and (2.0.15), we obtain

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), z) \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) + d_{\mathbf{p}'}(z, \vartheta) \leq 0.2 + 2^{-95a} \cdot 1 < 0.3.$$

Thus $z \in \Omega_1(\mathbf{p}')$ by (4.2.6). By Lemma 4.2.2, it follows that $\mathbf{p} = \mathbf{p}'$.

3. There exists $z' \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$ such that $\vartheta \in \Omega(J, z')$, and $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$. This case is the same as case 2., after swapping \mathbf{p} and \mathbf{p}' .

4. We have $\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2) \cap B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), 0.2)$. In this case it follows that $\mathbf{p} = \mathbf{p}'$ since the sets $B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 0.2)$ are pairwise disjoint by the inclusion (4.2.6) and Lemma 4.2.2.

To show the inclusion in (2.0.13), let $\vartheta \in \mathcal{Q}(X)$. By the induction hypothesis, there exists $\mathbf{p} \in \mathfrak{P}(J)$ such that $\vartheta \in \Omega(\mathbf{p})$. By definition of the set \mathfrak{P} , we have $\mathbf{p} = (J, z)$ for some $z \in \mathcal{Z}(J)$. Thus, by (4.2.3), there exists $z' \in \mathcal{Z}(I)$ with $z \in B_{J'}(z', 0.7)$. Then by Lemma (4.2.3) there exists $\mathbf{p}' \in \mathfrak{P}(I)$ with $z \in \mathcal{Z}(J) \cap \Omega_1(\mathbf{p}')$. Consequently, by (4.2.8), $\vartheta \in \Omega(\mathbf{p}')$. This completes the proof of (2.0.13).

Finally, we show (2.0.14). Let $\mathbf{p}, \mathbf{q} \in \mathfrak{P}$ with $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{q})$ and $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) \neq \emptyset$. If we have $s(\mathbf{p}) \geq s(\mathbf{q})$, then it follows from (2.0.8) that $I = J$, thus $\mathbf{p}, \mathbf{q} \in \mathfrak{P}(I)$. By (2.0.13) we have then either $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) = \emptyset$ or $\Omega(\mathbf{p}) = \Omega(\mathbf{q})$. By the assumption in (2.0.14) we have $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}) \neq \emptyset$, so we must have $\Omega(\mathbf{p}) = \Omega(\mathbf{q})$ and in particular $\Omega(\mathbf{q}) \subset \Omega(\mathbf{p})$.

So it remains to show (2.0.14) under the additional assumption that $s(\mathbf{q}) > s(\mathbf{p})$. In this case, we argue by induction on $s(\mathbf{q}) - s(\mathbf{p})$. By (2.0.7), there exists a cube $J \in \mathcal{D}$ with $s(J) = s(\mathbf{q}) - 1$ and $J \cap \mathcal{J}(\mathbf{p}) \neq \emptyset$. We pick one such J . By (2.0.8), we have $\mathcal{J}(\mathbf{p}) \subset J \subset \mathcal{J}(\mathbf{q})$.

Thus, by (4.2.3), there exists $z' \in \mathcal{Z}(J)$ with $\mathcal{Q}(\mathbf{q}) \in B_{J'}(z', 0.7)$. Then by Lemma 4.2.3 there exists $\mathbf{q}' \in \mathfrak{P}(J)$ with $\mathcal{Q}(\mathbf{q}) \in \Omega_1(\mathbf{q}')$. By (4.2.8), it follows that $\Omega(\mathbf{q}) \subset \Omega(\mathbf{q}')$. Note that then $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{q}')$ and $\Omega(\mathbf{p}) \cap \Omega(\mathbf{q}') \neq \emptyset$ and $s(\mathbf{q}') - s(\mathbf{p}) = s(\mathbf{q}) - s(\mathbf{p}) - 1$. Thus, we have by the induction hypothesis that $\Omega(\mathbf{q}') \subset \Omega(\mathbf{p})$. This completes the proof. \square

5. PROOF OF DISCRETE CARLESON

Let a grid structure (\mathcal{D}, c, s) and a tile structure $(\mathfrak{P}, \mathcal{J}, \Omega, \mathcal{Q})$ for this grid structure be given. In Section 5.1, we decompose the set \mathfrak{P} of tiles into subsets. Each subset

will be controlled by one of three methods. The guiding principle of the decomposition is to be able to apply the forest estimate of Proposition 2.0.4 to the final subsets defined in (5.1.23). This application is done in Section 5.4. The miscellaneous subsets along the construction of the forests will either be thrown into exceptional sets, which are defined and controlled in Section 5.2, or will be controlled by the antichain estimate of Proposition 2.0.3, which is done in Section 5.5. Section 5.3 contains some auxiliary lemmas needed for the proofs in Subsections 5.4-5.5.

5.1. Organisation of the tiles. In the following definitions, k, n , and j will be nonnegative integers. Define $\mathcal{C}(G, k)$ to be the set of $I \in \mathcal{D}$ such that there exists a $J \in \mathcal{D}$ with $I \subset J$ and

$$\mu(G \cap J) > 2^{-k-1} \mu(J), \quad (5.1.1)$$

but there does not exist a $J \in \mathcal{D}$ with $I \subset J$ and

$$\mu(G \cap J) > 2^{-k} \mu(J). \quad (5.1.2)$$

Let

$$\mathfrak{P}(k) = \{\mathfrak{p} \in \mathfrak{P} : \mathcal{J}(\mathfrak{p}) \in \mathcal{C}(G, k)\} \quad (5.1.3)$$

Define $\mathfrak{M}(k, n)$ to be the set of $\mathfrak{p} \in \mathfrak{P}(k)$ such that

$$\mu(E_1(\mathfrak{p})) > 2^{-n} \mu(\mathcal{J}(\mathfrak{p})) \quad (5.1.4)$$

and there does not exist $\mathfrak{p}' \in \mathfrak{P}(k)$ with $\mathfrak{p}' \neq \mathfrak{p}$ and $\mathfrak{p} \leq \mathfrak{p}'$ such that

$$\mu(E_1(\mathfrak{p}')) > 2^{-n} \mu(\mathcal{J}(\mathfrak{p}')). \quad (5.1.5)$$

Define for a collection $\mathfrak{P}' \subset \mathfrak{P}(k)$

$$\text{dens}'_k(\mathfrak{P}') := \sup_{\mathfrak{p}' \in \mathfrak{P}'} \sup_{\lambda \geq 2} \lambda^{-a} \sup_{\mathfrak{p} \in \mathfrak{P}(k) : \lambda \mathfrak{p}' \leq \lambda \mathfrak{p}} \frac{\mu(E_2(\lambda, \mathfrak{p}))}{\mu(\mathcal{J}(\mathfrak{p}))}. \quad (5.1.6)$$

Sorting by density, we define

$$\mathfrak{C}(k, n) := \{\mathfrak{p} \in \mathfrak{P}(k) : 2^{4a} 2^{-n} < \text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{4a} 2^{-n+1}\}. \quad (5.1.7)$$

Following Fefferman [Fef73], we define for $\mathfrak{p} \in \mathfrak{C}(k, n)$

$$\mathfrak{B}(\mathfrak{p}) := \{\mathfrak{m} \in \mathfrak{M}(k, n) : 100\mathfrak{p} \lesssim \mathfrak{m}\} \quad (5.1.8)$$

and

$$\mathfrak{C}_1(k, n, j) := \{\mathfrak{p} \in \mathfrak{C}(k, n) : 2^j \leq |\mathfrak{B}(\mathfrak{p})| < 2^{j+1}\}. \quad (5.1.9)$$

and

$$\mathfrak{L}_0(k, n) := \{\mathfrak{p} \in \mathfrak{C}(k, n) : |\mathfrak{B}(\mathfrak{p})| < 1\}. \quad (5.1.10)$$

Together with the following removal of minimal layers, the splitting into $\mathfrak{C}_1(k, n, j)$ will lead to a separation of trees. Define recursively for $0 \leq l \leq Z(n+1)$

$$\mathfrak{L}_1(k, n, j, l) \quad (5.1.11)$$

to be the set of minimal elements with respect to \leq in

$$\mathfrak{C}_1(k, n, j) \setminus \bigcup_{0 \leq l' < l} \mathfrak{L}_1(k, n, j, l'). \quad (5.1.12)$$

Define

$$\mathfrak{C}_2(k, n, j) := \mathfrak{C}_1(k, n, j) \setminus \bigcup_{0 \leq l' \leq Z(n+1)} \mathfrak{L}_1(k, n, j, l'). \quad (5.1.13)$$

The remaining tile organization will be relative to prospective tree tops, which we define now. Define

$$\mathfrak{U}_1(k, n, j) \quad (5.1.14)$$

to be the set of all $\mathbf{u} \in \mathfrak{C}_1(k, n, j)$ such that for all $\mathbf{p} \in \mathfrak{C}_1(k, n, j)$ with $\mathcal{J}(\mathbf{u})$ strictly contained in $\mathcal{J}(\mathbf{p})$ we have $B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100) \cap B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 100) = \emptyset$.

We first remove the pairs that are outside the immediate reach of any of the prospective tree tops. Define

$$\mathfrak{L}_2(k, n, j) \quad (5.1.15)$$

to be the set of all $\mathbf{p} \in \mathfrak{C}_2(k, n, j)$ such that there does not exist $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $\mathcal{J}(\mathbf{p}) \neq \mathcal{J}(\mathbf{u})$ and $2\mathbf{p} \lesssim \mathbf{u}$. Define

$$\mathfrak{C}_3(k, n, j) := \mathfrak{C}_2(k, n, j) \setminus \mathfrak{L}_2(k, n, j). \quad (5.1.16)$$

We next remove the maximal layers. Define recursively for $0 \leq l \leq Z(n+1)$

$$\mathfrak{L}_3(k, n, j, l) \quad (5.1.17)$$

to be the set of all maximal elements with respect to \leq in

$$\mathfrak{C}_3(k, n, j) \setminus \bigcup_{0 \leq l' < l} \mathfrak{L}_3(k, n, j, l'). \quad (5.1.18)$$

Define

$$\mathfrak{C}_4(k, n, j) := \mathfrak{C}_3(k, n, j) \setminus \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_3(k, n, j, l). \quad (5.1.19)$$

Finally, we remove the boundary pairs relative to the prospective tree tops. Define

$$\mathcal{L}(\mathbf{u}) \quad (5.1.20)$$

to be the set of all $I \in \mathcal{D}$ with $I \subset \mathcal{J}(\mathbf{u})$ and $s(I) = s(\mathbf{u}) - Z(n+1) - 1$ and

$$B(c(I), 8D^{s(I)}) \not\subset \mathcal{J}(\mathbf{u}). \quad (5.1.21)$$

Define

$$\mathfrak{L}_4(k, n, j) \quad (5.1.22)$$

to be the set of all $\mathbf{p} \in \mathfrak{C}_4(k, n, j)$ such that there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $\mathcal{J}(\mathbf{p}) \subset \bigcup \mathcal{L}(\mathbf{u})$, and define

$$\mathfrak{C}_5(k, n, j) := \mathfrak{C}_4(k, n, j) \setminus \mathfrak{L}_4(k, n, j). \quad (5.1.23)$$

We define three exceptional sets. The first exceptional set G_1 takes into account the ratio of the measures of F and G . Define $\mathfrak{P}_{F,G}$ to be the set of all $\mathbf{p} \in \mathfrak{P}$ with

$$\text{dens}_2(\{\mathbf{p}\}) > 2^{2a+5} \frac{\mu(F)}{\mu(G)}. \quad (5.1.24)$$

Define

$$G_1 := \bigcup_{\mathbf{p} \in \mathfrak{P}_{F,G}} \mathcal{J}(\mathbf{p}). \quad (5.1.25)$$

For an integer $\lambda \geq 0$, define $A(\lambda, k, n)$ to be the set of all $x \in X$ such that

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{J}(\mathbf{p})}(x) > \lambda 2^{n+1} \quad (5.1.26)$$

and define

$$G_2 := \bigcup_{k \geq 0} \bigcup_{k \leq n} A(2n + 6, k, n). \quad (5.1.27)$$

Define

$$G_3 := \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{\mathfrak{p} \in \mathfrak{L}_4(k, n, j)} \mathcal{J}(\mathfrak{p}). \quad (5.1.28)$$

Define $G' = G_1 \cup G_2 \cup G_3$. The following bound of the measure of G' will be proven in Section 5.2.

Lemma 5.1.1 (exceptional set). *We have*

$$\mu(G') \leq 2^{-1} \mu(G). \quad (5.1.29)$$

In Section 5.4, we identify each set $\mathfrak{C}_5(k, n, j)$ outside G' as forest and use Proposition 2.0.4 to prove the following lemma.

Lemma 5.1.2 (forest union). *Let*

$$\mathfrak{F}_1 = \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \mathfrak{C}_5(k, n, j) \quad (5.1.30)$$

For all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{F}_1} T_{\mathfrak{p}} f \right| d\mu \leq \frac{2^{441a^3}}{(q-1)^4} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (5.1.31)$$

In Section 5.5, we decompose the complement of the set of tiles in Lemma 5.1.2 and apply the antichain estimate of Proposition 2.0.3 to prove the following lemma.

Lemma 5.1.3 (forest complement). *Let*

$$\mathfrak{F}_2 = \mathfrak{F} \setminus \mathfrak{F}_1. \quad (5.1.32)$$

For all $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$ we have

$$\int_{G \setminus G'} \left| \sum_{\mathfrak{p} \in \mathfrak{F}_2} T_{\mathfrak{p}} f \right| d\mu \leq \frac{2^{120a^3}}{(q-1)^5} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (5.1.33)$$

Proof of Proposition 2.0.2. Proposition 2.0.2 follows by applying the triangle inequality to (2.0.22) according to the splitting in Lemma 5.1.2 and Lemma 5.1.3 and using both Lemmas as well as the bound on the set G' given by Lemma 5.1.1. \square

5.2. Proof of the Exceptional Sets Lemma. We prove separate bounds for G_1 , G_2 , and G_3 in Lemmas 5.2.1, 5.2.6, and 5.2.10. Adding up these bounds proves Lemma 5.1.1.

The bound for G_1 follows from the Vitali covering lemma, Proposition 2.0.6.

Lemma 5.2.1 (first exception). *We have*

$$\mu(G_1) \leq 2^{-5} \mu(G). \quad (5.2.1)$$

Proof. Let

$$K = 2^{2a+5} \frac{\mu(F)}{\mu(G)}.$$

For each $\mathbf{p} \in \mathfrak{P}_{F,G}$ pick a $r(\mathbf{p}) > 4D^{s(\mathbf{p})}$ with

$$\mu(F \cap B(c(\mathbf{p}), r(\mathbf{p}))) \geq K\mu(B(c(\mathbf{p}), r(\mathbf{p}))).$$

This ball exists by definition of $\mathfrak{P}_{F,G}$ and dens_2 . By applying Proposition 2.0.6 to the collection of balls

$$\mathcal{B} = \{B(c(\mathbf{p}), r(\mathbf{p})) : \mathbf{p} \in \mathfrak{P}_{F,G}\}$$

and the function $u = \mathbf{1}_F$, we obtain

$$\mu\left(\bigcup \mathcal{B}\right) \leq 2^{2a} K^{-1} \mu(F).$$

We conclude with (2.0.10) and $r(\mathbf{p}) > 4D^{s(\mathbf{p})}$

$$\mu(G_1) = \mu\left(\bigcup_{\mathbf{p} \in \mathfrak{P}_{F,G}} \mathcal{J}(\mathbf{p})\right) \leq \mu\left(\bigcup \mathcal{B}\right) \leq 2^{2a} K^{-1} \mu(F) = 2^{-5} \mu(G).$$

□

We turn to the bound of G_2 , which relies on the Dyadic Covering Lemma 5.2.2 and the John-Nirenberg Lemma 5.2.5 below.

Lemma 5.2.2 (dense cover). *For each $k \geq 0$, the union of all dyadic cubes in $\mathcal{C}(G, k)$ has measure at most $2^{k+1} \mu(G)$.*

Proof. The union of dyadic cubes in $\mathcal{C}(G, k)$ is contained the union of elements of the set $\mathcal{M}(k)$ of all dyadic cubes J with $\mu(G \cap J) > 2^{-k-1} \mu(J)$. The union of elements in the set $\mathcal{M}(k)$ is contained in the union of elements in the set $\mathcal{M}^*(k)$ of maximal elements in $\mathcal{M}(k)$ with respect to set inclusion. Hence

$$\mu\left(\bigcup \mathcal{C}(G, k)\right) \leq \mu\left(\bigcup \mathcal{M}^*(k)\right) \leq \sum_{J \in \mathcal{M}^*(k)} \mu(J) \quad (5.2.2)$$

Using the definition of $\mathcal{M}(k)$ and then the pairwise disjointedness of elements in $\mathcal{M}^*(k)$, we estimate (5.2.2) by

$$\leq 2^{k+1} \sum_{J \in \mathcal{M}^*(k)} \mu(J \cap G) \leq 2^{k+1} \mu(G). \quad (5.2.3)$$

This proves the lemma. □

Lemma 5.2.3 (pairwise disjoint). *If $\mathbf{p}, \mathbf{p}' \in \mathfrak{M}(k, n)$ and*

$$E_1(\mathbf{p}) \cap E_1(\mathbf{p}') \neq \emptyset, \quad (5.2.4)$$

then $\mathbf{p} = \mathbf{p}'$.

Proof. Let \mathbf{p}, \mathbf{p}' be as in the lemma. By definition of E_1 , we have $E_1(\mathbf{p}) \subset \mathcal{J}(\mathbf{p})$ and analogously for \mathbf{p}' , we conclude from (5.2.4) that $\mathcal{J}(\mathbf{p}) \cap \mathcal{J}(\mathbf{p}') \neq \emptyset$. Let without loss of generality $\mathcal{J}(\mathbf{p})$ be maximal in $\{\mathcal{J}(\mathbf{p}), \mathcal{J}(\mathbf{p}')\}$, then $\mathcal{J}(\mathbf{p}') \subset \mathcal{J}(\mathbf{p})$. By (5.2.4), we conclude by definition of E_1 that $\Omega(\mathbf{p}) \cap \Omega(\mathbf{p}') \neq \emptyset$. By (2.0.14) we conclude $\Omega(\mathbf{p}) \subset \Omega(\mathbf{p}')$. It follows that $\mathbf{p}' \leq \mathbf{p}$. By maximality (5.1.5) of \mathbf{p}' , we have $\mathbf{p}' = \mathbf{p}$. This proves the lemma. □

Lemma 5.2.4 (dyadic union). *For each $x \in A(\lambda, k, n)$, there is a dyadic cube I that contains x and is a subset of $A(\lambda, k, n)$.*

Proof. Fix k, n, λ, x as in the lemma such that $x \in A(\lambda, k, n)$. Let \mathcal{M} be the set of dyadic cubes $\mathcal{J}(\mathbf{p})$ with \mathbf{p} in $\mathfrak{M}(k, n)$ and $x \in \mathcal{J}(\mathbf{p})$. By definition of $A(\lambda, k, n)$, the cardinality of \mathcal{M} is at least $1 + \lambda 2^{n+1}$. Let I be a cube of smallest scale in \mathcal{M} . Then I is contained in all cubes of \mathcal{M} . It follows that $I \subset A(\lambda, k, n)$. \square

Lemma 5.2.5 (John Nirenberg). *For all integers $k, n, \lambda \geq 0$, we have*

$$\mu(A(\lambda, k, n)) \leq 2^{k+1-\lambda} \mu(G). \quad (5.2.5)$$

Proof. Fix k, n as in the lemma and suppress notation to write $A(\lambda)$ for $A(\lambda, k, n)$. We prove the lemma by induction on λ . For $\lambda = 0$, we use that $A(\lambda)$ by definition of $\mathfrak{M}(k, n)$ is contained in the union of elements in $\mathcal{C}(G, k)$. Lemma 5.2.2 then completes the base of the induction.

Now assume that the statement of Lemma 5.2.5 is proven for some integer $\lambda \geq 0$. The set $A(\lambda + 1)$ is contained in the set $A(\lambda)$. Let \mathcal{M} be the set of dyadic cubes which are a subset of $A(\lambda)$. By Lemma 5.2.4, the union of \mathcal{M} is $A(\lambda)$. Let \mathcal{M}^* be the set of maximal dyadic cubes in \mathcal{M} .

Let $x \in A(\lambda + 1)$ and $L \in \mathcal{M}^*$ such that $x \in L$. Then by the dyadic property (2.0.8)

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{J}(\mathbf{p})}(x) = \sum_{\mathbf{p} \in \mathfrak{M}(k, n): \mathcal{J}(\mathbf{p}) \subset L} \mathbf{1}_{\mathcal{J}(\mathbf{p})}(x) + \sum_{\mathbf{p} \in \mathfrak{M}(k, n): L \subsetneq \mathcal{J}(\mathbf{p})} \mathbf{1}_{\mathcal{J}(\mathbf{p})}(x). \quad (5.2.6)$$

We now show

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n): \mathcal{J}(\mathbf{p}) \subset L} \mathbf{1}_{\mathcal{J}(\mathbf{p})}(x) \geq 2^{n+1}. \quad (5.2.7)$$

The left-hand side of (5.2.6) is strictly greater than $(\lambda + 1)2^{n+1}$. If L is the top cube the second sum Q_2 on the right-hand side of (5.2.6) is zero and (5.2.7) follows immediately. Otherwise consider the inclusion-minimal cube \hat{L} with $L \subsetneq \hat{L}$; all tiles \mathbf{p} over which Q_2 is summed over satisfy $\hat{L} \subset \mathbf{p}$, so Q_2 is constant for all $x \in \hat{L}$. By maximality of L , Q_2 is at most $\lambda 2^{n+1}$ somewhere on \hat{L} , thus on all of \hat{L} and consequently also at x . Rearranging the inequality yields (5.2.7).

By Lemma 5.2.3, we have

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n): \mathcal{J}(\mathbf{p}) \subset L} \mu(E_1(\mathbf{p})) \leq \mu(L). \quad (5.2.8)$$

Multiplying by 2^n and applying (5.1.4), we obtain

$$\sum_{\mathbf{p} \in \mathfrak{M}(k, n): \mathcal{J}(\mathbf{p}) \subset L} \mu(\mathcal{J}(\mathbf{p})) \leq 2^n \mu(L). \quad (5.2.9)$$

We then have with (5.2.7) and (5.2.9)

$$2^{n+1} \mu(A(\lambda + 1) \cap L) = \int_{A(\lambda+1) \cap L} 2^{n+1} d\mu \quad (5.2.10)$$

$$\leq \int \sum_{\mathbf{p} \in \mathfrak{M}(k, n): \mathcal{J}(\mathbf{p}) \subset L} \hat{\mathbf{1}}_{\mathcal{J}(\mathbf{p})} d\mu \leq 2^n \mu(L). \quad (5.2.11)$$

Hence

$$2\mu(A(\lambda + 1)) = 2 \sum_{L \in \mathcal{M}^*} \mu(A(\lambda + 1) \cap L) \leq \sum_{L \in \mathcal{M}^*} \mu(L) = \mu(A(\lambda)). \quad (5.2.12)$$

Using the induction hypothesis, this proves (5.2.5) for $\lambda + 1$ and completes the proof of the lemma. \square

Lemma 5.2.6 (second exception). *We have*

$$\mu(G_2) \leq 2^{-2}\mu(G). \quad (5.2.13)$$

Proof. We use Lemma 5.2.5 and sum twice a geometric series to obtain

$$\sum_{0 \leq k \leq n} \sum_{k \leq n} \mu(A(2n + 6, k, n)) \leq \sum_{0 \leq k \leq n} \sum_{k \leq n} 2^{k-5-2n} \mu(G) \quad (5.2.14)$$

$$\leq \sum_{0 \leq k} 2^{-k-3} \mu(G) \leq 2^{-2} \mu(G). \quad (5.2.15)$$

This proves the lemma. \square

We turn to the set G_3 .

Lemma 5.2.7 (top tiles). *We have*

$$\sum_{\mathfrak{m} \in \mathfrak{M}(k, n)} \mu(\mathcal{J}(\mathfrak{m})) \leq 2^{n+k+3} \mu(G). \quad (5.2.16)$$

Proof. We write the left-hand side of (5.2.16)

$$\int \sum_{\mathfrak{m} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{J}(\mathfrak{m})}(x) d\mu(x) \leq 2^{n+1} \sum_{\lambda=0}^{|\mathfrak{M}|} \mu(A(\lambda, k, n)). \quad (5.2.17)$$

Using Lemma 5.2.5 and then summing a geometric series, we estimate this by

$$\leq 2^{n+1} \sum_{\lambda=0}^{|\mathfrak{M}|} 2^{k+1-\lambda} \mu(G) \leq 2^{n+1} 2^{k+2} \mu(G). \quad (5.2.18)$$

This proves the lemma. \square

Lemma 5.2.8 (tree count). *Let $k, n, j \geq 0$. We have for every $x \in X$*

$$\sum_{\mathfrak{u} \in \mathfrak{U}_1(k, n, j)} \mathbf{1}_{\mathcal{J}(\mathfrak{u})}(x) \leq 2^{-j} 2^{9a} \sum_{\mathfrak{m} \in \mathfrak{M}(k, n)} \mathbf{1}_{\mathcal{J}(\mathfrak{m})}(x) \quad (5.2.19)$$

Proof. Let $x \in X$. For each $\mathfrak{u} \in \mathfrak{U}_1(k, n, j)$ with $x \in \mathcal{J}(\mathfrak{u})$, as $\mathfrak{u} \in \mathfrak{C}_1(k, n, j)$, there are at least 2^j elements $\mathfrak{m} \in \mathfrak{M}(k, n)$ with $100\mathfrak{u} \lesssim \mathfrak{m}$ and in particular $x \in \mathcal{J}(\mathfrak{m})$. Hence

$$\mathbf{1}_{\mathcal{J}(\mathfrak{u})}(x) \leq 2^{-j} \sum_{\mathfrak{m} \in \mathfrak{M}(k, n): 100\mathfrak{u} \lesssim \mathfrak{m}} \mathbf{1}_{\mathcal{J}(\mathfrak{m})}(x). \quad (5.2.20)$$

Conversely, for each $\mathfrak{m} \in \mathfrak{M}(k, n)$ with $x \in \mathcal{J}(\mathfrak{m})$, let $\mathfrak{U}(\mathfrak{m})$ be the set of $\mathfrak{u} \in \mathfrak{U}_1(k, n, j)$ with $x \in \mathcal{J}(\mathfrak{u})$ and $100\mathfrak{u} \lesssim \mathfrak{m}$. Summing (5.2.20) over \mathfrak{u} and counting the pairs $(\mathfrak{u}, \mathfrak{m})$ with $100\mathfrak{u} \lesssim \mathfrak{m}$ differently gives

$$\sum_{\mathfrak{u} \in \mathfrak{U}_1(k, n, j)} \mathbf{1}_{\mathcal{J}(\mathfrak{u})}(x) \leq 2^{-j} \sum_{\mathfrak{m} \in \mathfrak{M}(k, n)} \sum_{\mathfrak{u} \in \mathfrak{U}(\mathfrak{m})} \mathbf{1}_{\mathcal{J}(\mathfrak{m})}(x). \quad (5.2.21)$$

We estimate the number of elements in $\mathfrak{U}(\mathfrak{m})$. Let $\mathfrak{u} \in \mathfrak{U}(\mathfrak{m})$. Then by definition of $\mathfrak{U}(\mathfrak{m})$

$$d_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), \mathcal{Q}(\mathfrak{m})) \leq 100. \quad (5.2.22)$$

If \mathfrak{u}' is a further element in $\mathfrak{U}(\mathfrak{m})$ with $\mathfrak{u} \neq \mathfrak{u}'$, then

$$\mathcal{Q}(\mathfrak{m}) \in B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100). \quad (5.2.23)$$

By the last display and definition of $\mathfrak{U}_1(k, n, j)$, none of $\mathcal{J}(\mathfrak{u})$, $\mathcal{J}(\mathfrak{u}')$ is strictly contained in the other. As both contain x , we have $\mathcal{J}(\mathfrak{u}) = \mathcal{J}(\mathfrak{u}')$. We then have $d_{\mathfrak{u}} = d_{\mathfrak{u}'}$.

By (2.0.15), the balls $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 0.2)$ and $B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 0.2)$ are contained respectively in $\Omega(\mathfrak{u})$ and $\Omega(\mathfrak{u}')$ and thus are disjoint by (2.0.13). By (5.2.22) and the triangle inequality, both balls are contained in $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{m}), 100.2)$.

By (1.1.8) applied nine times, there is a collection of at most 2^{9a} balls of radius 0.2 with respect to the metric $d_{\mathfrak{u}}$ which cover the ball $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{m}), 100.2)$. Let B' be a ball in this cover. As the center of B' can be in at most one of the disjoint balls $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 0.2)$ and $B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 0.2)$, the ball B' can contain at most one of the points $\mathcal{Q}(\mathfrak{u})$, $\mathcal{Q}(\mathfrak{u}')$.

Hence the image of $\mathfrak{U}(\mathfrak{m})$ under \mathcal{Q} has at most 2^{9a} elements; since \mathcal{Q} is injective on $\mathfrak{U}(\mathfrak{m})$, the same is true of $\mathfrak{U}(\mathfrak{m})$. Inserting this into (5.2.21) proves the lemma. \square

Lemma 5.2.9 (boundary exception). *Let $\mathcal{L}(\mathfrak{u})$ be as defined in (5.1.20). We have for each $\mathfrak{u} \in \mathfrak{U}_1(k, n, l)$,*

$$\mu\left(\bigcup_{I \in \mathcal{L}(\mathfrak{u})} I\right) \leq D^{1-\kappa Z(n+1)} \mu(\mathcal{J}(\mathfrak{u})). \quad (5.2.24)$$

Proof. Let $\mathfrak{u} \in \mathfrak{U}_1(k, n, l)$. Let $I \in \mathcal{L}(\mathfrak{u})$. Then we have $s(I) = s(\mathfrak{u}) - Z(n+1) - 1$ and $I \subset \mathcal{J}(\mathfrak{u})$ and $B(c(I), 8D^{s(I)}) \not\subset \mathcal{J}(\mathfrak{u})$. By (2.0.10), the set I is contained in $B(c(I), 4D^{s(I)})$. By the triangle inequality, the set I is contained in

$$X(\mathfrak{u}) := \{x \in \mathcal{J}(\mathfrak{u}) : \rho(x, X \setminus \mathcal{J}(\mathfrak{u})) \leq 12D^{s(\mathfrak{u})-Z(n+1)-1}\}. \quad (5.2.25)$$

By the small boundary property (2.0.11), noting that

$$12D^{s(\mathfrak{u})-Z(n+1)-1} = 12D^{s(I)} > D^{-S},$$

we have

$$\mu(X(\mathfrak{u})) \leq 2 \cdot (12D^{-Z(n+1)-1})^{\kappa} \mu(\mathcal{J}(\mathfrak{u})).$$

Using $\kappa < 1$ and $D \geq 12$, this proves the lemma. \square

Lemma 5.2.10 (third exception). *We have*

$$\mu(G_3) \leq 2^{-4} \mu(G). \quad (5.2.26)$$

Proof. As each $\mathfrak{p} \in \mathfrak{L}_4(k, n, j)$ is contained in $\cup \mathcal{L}(\mathfrak{u})$ for some $\mathfrak{u} \in \mathfrak{U}_1(k, n, l)$, we have

$$\mu\left(\bigcup_{\mathfrak{p} \in \mathfrak{L}_4(k, n, j)} \mathcal{J}(\mathfrak{p})\right) \leq \sum_{\mathfrak{u} \in \mathfrak{U}_1(k, n, j)} \mu\left(\bigcup_{I \in \mathcal{L}(\mathfrak{u})} I\right). \quad (5.2.27)$$

Using Lemma 5.2.9 and then Lemma 5.2.8, we estimate this further by

$$\leq \sum_{\mathfrak{u} \in \mathfrak{U}_1(k, n, j)} D^{1-\kappa Z(n+1)} \mu(\mathcal{J}(\mathfrak{u})) \quad (5.2.28)$$

$$\leq 2^{100a^2+9a+1-j} \sum_{\mathbf{m} \in \mathfrak{M}(k,n)} D^{-\kappa Z(n+1)} \mu(\mathcal{J}(\mathbf{m})). \quad (5.2.29)$$

Using Lemma 5.2.7, we estimate this by

$$\leq 2^{100a^2+9a+1-j} D^{-\kappa Z(n+1)} 2^{n+k+3} \mu(G). \quad (5.2.30)$$

Now we estimate G_3 defined in (5.1.28) by

$$\mu(G_3) \leq \sum_{k \geq 0} \sum_{n \geq k} \sum_{0 \leq j \leq 2n+3} \mu\left(\bigcup_{\mathbf{p} \in \mathfrak{D}_4(k,n,j)} \mathcal{J}(\mathbf{p})\right) \quad (5.2.31)$$

$$\leq \sum_{k \geq 0} \sum_{n \geq k} \sum_{0 \leq j \leq 2n+3} 2^{100a^2+9a+3+n+k-j} D^{-\kappa Z(n+1)} \mu(G) \quad (5.2.32)$$

Summing geometric series, using that $D^{\kappa Z} \geq 8$ by (2.0.1), (2.0.2) and (2.0.3), we estimate this by

$$\leq \sum_{k \geq 0} \sum_{n \geq k} 2^{100a^2+9a+4+n+k} D^{-\kappa Z(n+1)} \mu(G) \quad (5.2.33)$$

$$= \sum_{k \geq 0} 2^{100a^2+9a+4+2k} D^{-\kappa Z(k+1)} \sum_{n \geq k} 2^{n-k} D^{-\kappa Z(n-k)} \mu(G) \quad (5.2.34)$$

$$\leq \sum_{k \geq 0} 2^{100a^2+9a+5+2k} D^{-\kappa Z(k+1)} \mu(G) \quad (5.2.35)$$

$$\leq 2^{100a^2+9a+6} D^{-\kappa Z} \mu(G) \quad (5.2.36)$$

Using $D = 2^{100a^2}$ and $a \geq 4$ and $\kappa Z \geq 2$ by (2.0.1) and (2.0.2) proves the lemma. \square

Proof of Lemma 5.1.1. Adding up the bounds in Lemmas 5.2.1, 5.2.6, and 5.2.10 proves Lemma 5.1.1. \square

5.3. Auxiliary lemmas. Before proving Lemma 5.1.2 and Lemma 5.1.3, we collect some useful properties of \lesssim .

Lemma 5.3.1 (wiggle order 1). *If $n\mathbf{p} \lesssim m\mathbf{p}'$ and $n' \geq n$ and $m \geq m'$ then $n'\mathbf{p} \lesssim m'\mathbf{p}'$.*

Proof. This follows immediately from the definition (2.0.24) of \lesssim and the two inclusions $B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), n) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), n')$ and $B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), m') \subset B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), m)$. \square

Lemma 5.3.2 (wiggle order 2).

Let $n, m \geq 1$ and $k > 0$. If $\mathbf{p}, \mathbf{p}' \in \mathfrak{P}$ with $\mathcal{J}(\mathbf{p}) \neq \mathcal{J}(\mathbf{p}')$ and

$$n\mathbf{p} \lesssim k\mathbf{p}' \quad (5.3.1)$$

then

$$(n + 2^{-95a}m)\mathbf{p} \lesssim m\mathbf{p}'. \quad (5.3.2)$$

Proof. The assumption (5.3.1) together with the definition (2.0.24) of \lesssim implies that $\mathcal{J}(\mathbf{p}) \subsetneq \mathcal{J}(\mathbf{p}')$. Let $\vartheta \in B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), m)$. Then we have by the triangle inequality

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}'), \vartheta)$$

The first summand is bounded by n since

$$\mathcal{Q}(\mathbf{p}') \in B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), k) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), n),$$

using (2.0.24). For the second summand we use Lemma 2.1.2 to show that the sum is estimated by

$$n + 2^{-95a}d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) < n + 2^{-95a}m.$$

Thus $B_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), m) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), n + 2^{-95a}m)$. Combined with $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{p}')$, this yields (5.3.2). \square

Lemma 5.3.3 (wiggle order 3). *The following implications hold for all $\mathbf{q}, \mathbf{q}' \in \mathfrak{P}$:*

$$\mathbf{q} \leq \mathbf{q}' \text{ and } \lambda \geq 1.1 \implies \lambda \mathbf{q} \lesssim \lambda \mathbf{q}', \quad (5.3.3)$$

$$10\mathbf{q} \lesssim \mathbf{q}' \text{ and } \mathcal{J}(\mathbf{q}) \neq \mathcal{J}(\mathbf{q}') \implies 100\mathbf{q} \lesssim 100\mathbf{q}', \quad (5.3.4)$$

$$2\mathbf{q} \lesssim \mathbf{q}' \text{ and } \mathcal{J}(\mathbf{q}) \neq \mathcal{J}(\mathbf{q}') \implies 4\mathbf{q} \lesssim 500\mathbf{q}'. \quad (5.3.5)$$

Proof. (5.3.4) and (5.3.5) are easy consequences of Lemma 5.3.1, Lemma 5.3.2 and the fact that $a \geq 4$. For (5.3.3), if $\mathcal{J}(\mathbf{q}) = \mathcal{J}(\mathbf{q}')$ then we get $\mathbf{q} = \mathbf{q}'$ by (2.0.13) and (2.0.23). If $\mathcal{J}(\mathbf{q}) \neq \mathcal{J}(\mathbf{q}')$, then from (2.0.23), (2.0.24) and (2.0.15) it follows that $\mathbf{q} \lesssim 0.2\mathbf{q}'$, and (5.3.3) follows from an easy calculation using Lemma 5.3.2. \square

We call a collection \mathfrak{A} of tiles convex if

$$\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}'' \text{ and } \mathbf{p}, \mathbf{p}'' \in \mathfrak{A} \implies \mathbf{p}' \in \mathfrak{A}. \quad (5.3.6)$$

Lemma 5.3.4 (P convex). *For each k , the collection $\mathfrak{P}(k)$ is convex.*

Proof. Suppose that $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ and $\mathbf{p}, \mathbf{p}'' \in \mathfrak{P}(k)$. By (5.1.3) we have $\mathcal{J}(\mathbf{p}), \mathcal{J}(\mathbf{p}'') \in \mathcal{C}(G, k)$, so there exists by (5.1.1) some $J \in \mathcal{D}$ with

$$\mathcal{J}(\mathbf{p}') \subset \mathcal{J}(\mathbf{p}'') \subset J$$

and $\mu(G \cap J) > 2^{-k-1}\mu(J)$. Thus (5.1.1) holds for $\mathcal{J}(\mathbf{p}')$. On the other hand, by (5.1.2), there exists no $J \in \mathcal{D}$ with $\mathcal{J}(\mathbf{p}) \subset J$ and $\mu(G \cap J) > 2^{-k}\mu(J)$. Since $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{p}')$, this implies that (5.1.2) holds for $\mathcal{J}(\mathbf{p}')$. Hence $\mathcal{J}(\mathbf{p}') \in \mathcal{C}(G, k)$, and therefore by (5.1.3) $\mathbf{p}' \in \mathfrak{P}(k)$. \square

Lemma 5.3.5 (C convex). *For each k, n , the collection $\mathfrak{C}(k, n)$ is convex.*

Proof. Let $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$ with $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}(k, n)$. Then, in particular, $\mathbf{p}, \mathbf{p}'' \in \mathfrak{P}(k)$, so, by Lemma 5.3.4, $\mathbf{p}' \in \mathfrak{P}(k)$. Next, we show that if $\mathbf{q} \leq \mathbf{q}' \in \mathfrak{P}(k)$ then $\text{dens}'_k(\{\mathbf{q}\}) \geq \text{dens}'_k(\{\mathbf{q}'\})$. If $\mathbf{p} \in \mathfrak{P}(k)$ and $\lambda \geq 2$ with $\lambda \mathbf{q}' \lesssim \lambda \mathbf{p}$, then it follows from $\mathbf{q} \leq \mathbf{q}'$, (5.3.3) of Lemma 5.3.3 and transitivity of \lesssim that $\lambda \mathbf{q} \lesssim \lambda \mathbf{p}$. Thus the supremum in the definition (5.1.6) of $\text{dens}'_k(\{\mathbf{q}\})$ is over a superset of the set the supremum in the definition of $\text{dens}'_k(\{\mathbf{q}'\})$ is taken over, which shows $\text{dens}'_k(\{\mathbf{q}\}) \geq \text{dens}'_k(\{\mathbf{q}'\})$. From $\mathbf{p}' \leq \mathbf{p}''$, $\mathbf{p}'' \in \mathfrak{C}(k, n)$ and (5.1.7) it then follows that

$$2^{4a}2^{-n} < \text{dens}'_k(\{\mathbf{p}''\}) \leq \text{dens}'_k(\{\mathbf{p}'\}).$$

Similarly, it follows from $\mathbf{p} \leq \mathbf{p}'$, $\mathbf{p} \in \mathfrak{C}(k, n)$ and (5.1.7) that

$$\text{dens}'_k(\{\mathbf{p}'\}) \leq \text{dens}'_k(\{\mathbf{p}\}) \leq 2^{4a}2^{-n+1}.$$

Thus $\mathbf{p}' \in \mathfrak{C}(k, n)$. \square

Lemma 5.3.6 (C1 convex). *For each k, n, j , the collection $\mathfrak{C}_1(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_1(k, n, j)$. By Lemma 5.3.5 and the inclusion $\mathfrak{C}_1(k, n, j) \subset \mathfrak{C}(k, n)$, which holds by definition (5.1.9), we have $\mathfrak{p}' \in \mathfrak{C}(k, n)$. By (5.3.3) and transitivity of \lesssim we have that $\mathfrak{q} \leq \mathfrak{q}'$ and $100\mathfrak{q}' \lesssim \mathfrak{m}$ imply $100\mathfrak{q} \lesssim \mathfrak{m}$. So, by (5.1.8), $\mathfrak{B}(\mathfrak{p}'') \subset \mathfrak{B}(\mathfrak{p}') \subset \mathfrak{B}(\mathfrak{p})$. Consequently, by (5.1.9)

$$2^j \leq |\mathfrak{B}(\mathfrak{p}'')| \leq |\mathfrak{B}(\mathfrak{p}')| \leq |\mathfrak{B}(\mathfrak{p})| < 2^{j+1},$$

thus $\mathfrak{p}' \in \mathfrak{C}_1(k, n, j)$. \square

Lemma 5.3.7 (C2 convex). *For each k, n, j , the collection $\mathfrak{C}_2(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_2(k, n, j)$. By (5.1.13), we have

$$\mathfrak{C}_2(k, n, j) \subset \mathfrak{C}_1(k, n, j).$$

Combined with Lemma 5.3.6, it follows that $\mathfrak{p}' \in \mathfrak{C}_1(k, n, j)$. If $\mathfrak{p} = \mathfrak{p}'$ the statement is trivially true, otherwise suppose that $\mathfrak{p}' \notin \mathfrak{C}_2(k, n, j)$. By (5.1.13), this implies that there exists $0 \leq l' \leq Z(n+1)$ with $\mathfrak{p}' \in \mathfrak{L}_1(k, n, j, l')$. By the definition (5.1.11) of $\mathfrak{L}_1(k, n, j, l')$, this implies that \mathfrak{p}' is minimal with respect to \leq in $\mathfrak{C}_1(k, n, j) \setminus \bigcup_{l < l'} \mathfrak{L}_1(k, n, j, l)$. Since $\mathfrak{p} \leq \mathfrak{p}'$ and $\mathfrak{p} \in \mathfrak{C}_2(k, n, j)$, $\mathfrak{p} = \mathfrak{p}'$, a contradiction. \square

Lemma 5.3.8 (C3 convex). *For each k, n, j , the collection $\mathfrak{C}_3(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_3(k, n, j)$. By (5.1.16) and Lemma 5.3.7 it follows that $\mathfrak{p}' \in \mathfrak{C}_2(k, n, j)$. By (5.1.16) and (5.1.15), there exists $\mathfrak{u} \in \mathfrak{U}_1(k, n, j)$ with $2\mathfrak{p}'' \lesssim \mathfrak{u}$ and $\mathcal{J}(\mathfrak{p}'') \subsetneq \mathcal{J}(\mathfrak{u})$. From $\mathfrak{p}' \leq \mathfrak{p}''$, (5.3.3) and transitivity of \lesssim we then have $2\mathfrak{p}' \lesssim \mathfrak{u}$ and $\mathcal{J}(\mathfrak{p}') \subsetneq \mathcal{J}(\mathfrak{u})$, so $\mathfrak{p}' \in \mathfrak{C}_3(k, n, j)$. \square

Lemma 5.3.9 (C4 convex). *For each k, n, j , the collection $\mathfrak{C}_4(k, n, j)$ is convex.*

Proof. The proof is entirely analogous to Lemma 5.3.7, substituting \mathfrak{C}_4 for \mathfrak{C}_2 , \mathfrak{C}_3 for \mathfrak{C}_1 and $\mathfrak{p}' \leq \mathfrak{p}''$ for $\mathfrak{p} \leq \mathfrak{p}'$. \square

Lemma 5.3.10 (C5 convex). *For each k, n, j , the collection $\mathfrak{C}_5(k, n, j)$ is convex.*

Proof. Let $\mathfrak{p} \leq \mathfrak{p}' \leq \mathfrak{p}''$ with $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_5(k, n, j)$. Then $\mathfrak{p}, \mathfrak{p}'' \in \mathfrak{C}_4(k, n, j)$ by (5.1.23), and thus by Lemma 5.3.9 $\mathfrak{p}' \in \mathfrak{C}_4(k, n, j)$. It suffices to show that if $\mathfrak{p}' \in \mathfrak{L}_4(k, n, j)$ then $\mathfrak{p} \in \mathfrak{L}_4(k, n, j)$ by contraposition; this is true by (5.1.22) and $\mathfrak{p} \leq \mathfrak{p}'$. \square

Lemma 5.3.11 (dens compare). *We have for every $k \geq 0$ and $\mathfrak{P}' \subset \mathfrak{P}(k)$*

$$\text{dens}_1(\mathfrak{P}') \leq \text{dens}'_k(\mathfrak{P}'). \quad (5.3.7)$$

Proof. It suffices to show that for all $\mathfrak{p}' \in \mathfrak{P}'$ and $\lambda \geq 2$ and $\mathfrak{p} \in \mathfrak{P}(\mathfrak{P}')$ with $\lambda\mathfrak{p}' \lesssim \lambda\mathfrak{p}$ we have

$$\frac{\mu(E_2(\lambda, \mathfrak{p}))}{\mu(\mathcal{J}(\mathfrak{p}))} \leq \sup_{\mathfrak{p}'' \in \mathfrak{P}(k): \lambda\mathfrak{p}' \lesssim \lambda\mathfrak{p}''} \frac{\mu(E_2(\lambda, \mathfrak{p}''))}{\mu(\mathcal{J}(\mathfrak{p}''))}. \quad (5.3.8)$$

Let such \mathfrak{p}' , λ , \mathfrak{p} be given. It suffices to show that $\mathfrak{p} \in \mathfrak{P}(k)$, that is, it satisfies (5.1.1) and (5.1.2).

We show (5.1.1). As $\mathfrak{p} \in \mathfrak{P}(\mathfrak{P}')$, there exists $\mathfrak{p}'' \in \mathfrak{P}'$ with $\mathcal{J}(\mathfrak{p}') \subset \mathcal{J}(\mathfrak{p}'')$. By assumption on \mathfrak{P}' , we have $\mathfrak{p}'' \in \mathfrak{P}(k)$ and there exists $J \in \mathcal{D}$ with $\mathcal{J}(\mathfrak{p}'') \subset J$ and

$$\mu(G \cap J) > 2^{-k-1}\mu(J). \quad (5.3.9)$$

Then also $\mathcal{J}(\mathfrak{p}') \subset J$, which proves (5.1.1) for \mathfrak{p} .

We show (5.1.2). Assume to get a contradiction that there exists $J \in \mathcal{D}$ with $\mathcal{J}(\mathfrak{p}) \subset J$ and

$$\mu(G \cap J) > 2^{-k} \mu(J). \quad (5.3.10)$$

As $\lambda \mathfrak{p}' \lesssim \lambda \mathfrak{p}$, we have $\mathcal{J}(\mathfrak{p}') \subset \mathcal{J}(\mathfrak{p})$, and therefore $\mathcal{J}(\mathfrak{p}') \subset J$. This contradicts $\mathfrak{p}' \in \mathfrak{P}' \subset \mathfrak{P}(k)$. This proves (5.1.2) for \mathfrak{p} . \square

Lemma 5.3.12 (C dens1). *For each set $\mathfrak{A} \subset \mathfrak{C}(k, n)$, we have*

$$\text{dens}_1(\mathfrak{A}) \leq 2^{4a} 2^{-n+1}.$$

Proof. We have by Lemma 5.3.11 that $\text{dens}_1(\mathfrak{A}) \leq \text{dens}'_k(\mathfrak{A})$. Since $\mathfrak{A} \subset \mathfrak{C}(k, n)$, it follows from monotonicity of suprema and the definition (5.1.6) that $\text{dens}'_k(\mathfrak{A}) \leq \text{dens}'_k(\mathfrak{C}(k, n))$. By (5.1.6) and (5.1.7), we have

$$\text{dens}'_k(\mathfrak{C}(k, n)) = \sup_{\mathfrak{p} \in \mathfrak{C}(k, n)} \text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{4a} 2^{-n+1}.$$

\square

5.4. Proof of the Forest Union Lemma. Fix $k, n, j \geq 0$. Define

$$\mathfrak{C}_6(k, n, j)$$

to be the set of all tiles $\mathfrak{p} \in \mathfrak{C}_5(k, n, j)$ such that $\mathcal{J}(\mathfrak{p}) \not\subset G'$. The following chain of lemmas establishes that the set $\mathfrak{C}_6(k, n, j)$ can be written as a union of a small number of n -forests.

For $\mathfrak{u} \in \mathfrak{U}_1(k, n, j)$, define

$$\mathfrak{T}_1(\mathfrak{u}) := \{\mathfrak{p} \in \mathfrak{C}_1(k, n, j) : \mathcal{J}(\mathfrak{p}) \neq \mathcal{J}(\mathfrak{u}), 2\mathfrak{p} \lesssim \mathfrak{u}\}. \quad (5.4.1)$$

Define

$$\mathfrak{U}_2(k, n, j) := \{\mathfrak{u} \in \mathfrak{U}_1(k, n, j) : \mathfrak{T}_1(\mathfrak{u}) \cap \mathfrak{C}_6(k, n, j) \neq \emptyset\}. \quad (5.4.2)$$

Define a relation \sim on $\mathfrak{U}_2(k, n, j)$ by setting $\mathfrak{u} \sim \mathfrak{u}'$ for $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}_2(k, n, j)$ if $\mathfrak{u} = \mathfrak{u}'$ or there exists \mathfrak{p} in $\mathfrak{T}_1(\mathfrak{u})$ with $10\mathfrak{p} \lesssim \mathfrak{u}'$.

Lemma 5.4.1 (relation geometry). *If $\mathfrak{u} \sim \mathfrak{u}'$, then $\mathcal{J}(\mathfrak{u}) = \mathcal{J}(\mathfrak{u}')$ and*

$$B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) \neq \emptyset.$$

Proof. Let $\mathfrak{u}, \mathfrak{u}' \in \mathfrak{U}_2(k, n, j)$ with $\mathfrak{u} \sim \mathfrak{u}'$. If $\mathfrak{u} = \mathfrak{u}'$ then the conclusion of the Lemma clearly holds. Else, there exists $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$ such that $\mathcal{J}(\mathfrak{p}) \neq \mathcal{J}(\mathfrak{u})$ and $2\mathfrak{p} \lesssim \mathfrak{u}$ and $10\mathfrak{p} \lesssim \mathfrak{u}'$. Using Lemma 5.3.1 and (5.3.4) of Lemma 5.3.3, we deduce that

$$100\mathfrak{p} \lesssim 100\mathfrak{u}, \quad 100\mathfrak{p} \lesssim 100\mathfrak{u}'. \quad (5.4.3)$$

Now suppose that $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) = \emptyset$. Then we have $\mathfrak{B}(\mathfrak{u}) \cap \mathfrak{B}(\mathfrak{u}') = \emptyset$, by the definition (5.1.8) of \mathfrak{B} and the definition (2.0.24) of \lesssim , but also $\mathfrak{B}(\mathfrak{u}) \subset \mathfrak{B}(\mathfrak{p})$ and $\mathfrak{B}(\mathfrak{u}') \subset \mathfrak{B}(\mathfrak{p})$, by (5.1.8), (2.0.24) and (5.4.3). Hence,

$$|\mathfrak{B}(\mathfrak{p})| \geq |\mathfrak{B}(\mathfrak{u})| + |\mathfrak{B}(\mathfrak{u}')| \geq 2^j + 2^j = 2^{j+1},$$

which contradicts $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$. Therefore we must have

$$B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) \neq \emptyset.$$

It follows from $2\mathbf{p} \lesssim \mathbf{u}$ and $10\mathbf{p} \lesssim \mathbf{u}'$ that $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u})$ and $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}')$. By (2.0.8), it follows that $\mathcal{J}(\mathbf{u})$ and $\mathcal{J}(\mathbf{u}')$ are nested. Combining this with the conclusion of the last paragraph and definition (5.1.14) of $\mathfrak{A}_1(k, n, j)$, we obtain that $\mathcal{J}(\mathbf{u}) = \mathcal{J}(\mathbf{u}')$. \square

Lemma 5.4.2 (equivalence relation). *For each k, n, j , the relation \sim on $\mathfrak{A}_2(k, n, j)$ is an equivalence relation.*

Proof. Reflexivity holds by definition. For transitivity, suppose that

$$\mathbf{u}, \mathbf{u}', \mathbf{u}'' \in \mathfrak{A}_1(k, n, j)$$

and $\mathbf{u} \sim \mathbf{u}'$, $\mathbf{u}' \sim \mathbf{u}''$. By Lemma 5.4.1, it follows that $\mathcal{J}(\mathbf{u}) = \mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u}'')$, that there exists

$$\vartheta \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100) \cap B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100)$$

and that there exists

$$\theta \in B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100) \cap B_{\mathbf{u}''}(\mathcal{Q}(\mathbf{u}''), 100).$$

If $\mathbf{u} = \mathbf{u}'$, then $\mathbf{u} \sim \mathbf{u}''$ holds by assumption. Else, there exists by the definition of \sim some $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u})$ with $10\mathbf{p} \lesssim \mathbf{u}'$. Then we have $2\mathbf{p} \lesssim \mathbf{u}$ and $\mathbf{p} \neq \mathbf{u}$ by definition of $\mathfrak{T}_1(\mathbf{u})$, so $4\mathbf{p} \lesssim 500\mathbf{u}$ by (5.3.5). For $q \in B_{\mathbf{u}''}(\mathcal{Q}(\mathbf{u}''), 1)$ it follows by the triangle inequality that

$$\begin{aligned} d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), q) &\leq d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}}(\vartheta, \mathcal{Q}(\mathbf{u}')) \\ &\quad + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \theta) + d_{\mathbf{u}}(\theta, \mathcal{Q}(\mathbf{u}'')) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}''), q). \end{aligned}$$

Using (2.0.17) and the fact that $\mathcal{J}(\mathbf{u}) = \mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u}'')$ this equals

$$\begin{aligned} &d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}'}(\vartheta, \mathcal{Q}(\mathbf{u}')) \\ &\quad + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \theta) + d_{\mathbf{u}''}(\theta, \mathcal{Q}(\mathbf{u}'')) + d_{\mathbf{u}''}(\mathcal{Q}(\mathbf{u}''), q) \\ &< 100 + 100 + 100 + 100 + 1 < 500. \end{aligned}$$

Since $4\mathbf{p} \lesssim 500\mathbf{u}$, it follows that $d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), q) < 4 < 10$. We have shown that $B_{\mathbf{u}''}(\mathcal{Q}(\mathbf{u}''), 1) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 10)$, combining this with $\mathcal{J}(\mathbf{u}'') = \mathcal{J}(\mathbf{u})$ gives $\mathbf{u} \sim \mathbf{u}''$.

For symmetry suppose that $\mathbf{u} \sim \mathbf{u}'$. By Lemma (5.4.1), it follows that $\mathcal{J}(\mathbf{u}) = \mathcal{J}(\mathbf{u}')$ and that there exists $\vartheta \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100) \cap B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100)$. Again, for $\mathbf{u} = \mathbf{u}'$ symmetry is obvious, so suppose that $\mathbf{u} \neq \mathbf{u}'$. There exists $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u}')$, which then satisfies $2\mathbf{p} \lesssim \mathbf{u}'$ and $\mathcal{J}(\mathbf{p}) \neq \mathcal{J}(\mathbf{u}')$. By Lemma 5.3.1 and (5.3.5), it follows that

$$10\mathbf{p} \lesssim 4\mathbf{p} \lesssim 500\mathbf{u}'. \quad (5.4.4)$$

If $q \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1)$ then we have from the triangle inequality and the fact that $\mathcal{J}(\mathbf{u}) = \mathcal{J}(\mathbf{u}')$:

$$\begin{aligned} d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), q) &\leq d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}'}(\vartheta, \mathcal{Q}(\mathbf{u})) + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}), q) \\ &= d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}}(\vartheta, \mathcal{Q}(\mathbf{u})) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), q) \\ &< 100 + 100 + 1 < 500. \end{aligned}$$

Combining this with (5.4.4) and (2.0.24), we get

$$B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 10).$$

Since $2\mathbf{p} \lesssim \mathbf{u}'$, we have $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u})$. Thus, $10\mathbf{p} \lesssim \mathbf{u}$ which completes the proof of $\mathbf{u}' \sim \mathbf{u}$. \square

Choose a set $\mathfrak{U}_3(k, n, j)$ of representatives for the equivalence classes of \sim in $\mathfrak{U}_2(k, n, j)$. Define for each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$

$$\mathfrak{T}_2(\mathbf{u}) := \bigcup_{\mathbf{u} \sim \mathbf{u}'} \mathfrak{T}_1(\mathbf{u}') \cap \mathfrak{C}_6(k, n, j). \quad (5.4.5)$$

Lemma 5.4.3 (C6 forest). *We have*

$$\mathfrak{C}_6(k, n, j) = \bigcup_{\mathbf{u} \in \mathfrak{U}_3(k, n, j)} \mathfrak{T}_2(\mathbf{u}). \quad (5.4.6)$$

Proof. Let $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$. By (5.1.19) and (5.1.23), we have $\mathbf{p} \in \mathfrak{C}_3(k, n, j)$. By (5.1.15) and (5.1.16), there exists $\mathbf{u} \in \mathfrak{U}_1(k, n, j)$ with $2\mathbf{p} \lesssim \mathbf{u}$ and $\mathcal{J}(\mathbf{p}) \neq \mathcal{J}(\mathbf{u})$, that is, with $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u})$. Then $\mathfrak{T}_1(\mathbf{u})$ is clearly nonempty, so $\mathbf{u} \in \mathfrak{U}_2(k, n, j)$. By the definition of $\mathfrak{U}_3(k, n, j)$, there exists $\mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $\mathbf{u} \sim \mathbf{u}'$. By (5.4.5), we have $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u}')$. \square

Lemma 5.4.4 (forest geometry). *For each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$, the set $\mathfrak{T}_2(\mathbf{u})$ satisfies (2.0.32).*

Proof. Let $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$. By (5.4.5), there exists $\mathbf{u}' \sim \mathbf{u}$ with $\mathbf{p} \in \mathfrak{T}_1(\mathbf{u}')$. Then we have $2\mathbf{p} \lesssim \mathbf{u}'$ and $\mathcal{J}(\mathbf{p}) \neq \mathcal{J}(\mathbf{u}')$, so by (5.3.5) $4\mathbf{p} \lesssim 500\mathbf{u}'$. Further, by Lemma 5.4.1, we have that $\mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u})$ and there exists $\vartheta \in B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 100) \cap B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 100)$. Let $\theta \in B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1)$. Using the triangle inequality and the fact that $\mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u})$, we obtain

$$\begin{aligned} d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \theta) &\leq d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}), \theta) \\ &= d_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), \vartheta) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \vartheta) + d_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), \theta) \\ &< 100 + 100 + 1 < 500. \end{aligned}$$

Combining this with $4\mathbf{p} \lesssim 500\mathbf{u}'$, we obtain

$$B_{\mathbf{u}}(\mathcal{Q}(\mathbf{u}), 1) \subset B_{\mathbf{u}'}(\mathcal{Q}(\mathbf{u}'), 500) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 4).$$

Together with $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}') = \mathcal{J}(\mathbf{u})$, this gives $4\mathbf{p} \lesssim \mathbf{u}$, which is (2.0.32). \square

Lemma 5.4.5 (forest convex). *For each $\mathbf{u} \in \mathfrak{U}_3(k, n, j)$, the set $\mathfrak{T}_2(\mathbf{u})$ satisfies the convexity condition (2.0.33).*

Proof. Let $\mathbf{p}, \mathbf{p}'' \in \mathfrak{T}_2(\mathbf{u})$ and $\mathbf{p}' \in \mathfrak{P}$ with $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$. By (5.4.5) we have $\mathbf{p}, \mathbf{p}'' \in \mathfrak{C}_6(k, n, j) \subset \mathfrak{C}_5(k, n, j)$. By Lemma 5.3.10, we have $\mathbf{p}' \in \mathfrak{C}_5(k, n, j)$. Since $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$ we have $\mathcal{J}(\mathbf{p}) \not\subset G'$, so $\mathcal{J}(\mathbf{p}') \not\subset G'$ and therefore also $\mathbf{p}' \in \mathfrak{C}_6(k, n, j)$.

By (5.4.5) there exists $\mathbf{u}' \in \mathfrak{U}_2(k, n, j)$ with $\mathbf{p}'' \in \mathfrak{T}_1(\mathbf{u}')$ and hence $2\mathbf{p}'' \lesssim \mathbf{u}'$ and $\mathcal{J}(\mathbf{p}'') \neq \mathcal{J}(\mathbf{u}')$. Together this implies $\mathcal{J}(\mathbf{p}'') \subsetneq \mathcal{J}(\mathbf{u}')$. With the inclusion $\mathcal{J}(\mathbf{p}') \subset \mathcal{J}(\mathbf{p}'')$ from $\mathbf{p}' \leq \mathbf{p}''$, it follows that $\mathcal{J}(\mathbf{p}') \subsetneq \mathcal{J}(\mathbf{u}')$ and hence $\mathcal{J}(\mathbf{p}') \neq \mathcal{J}(\mathbf{u}')$. By (5.3.3) and transitivity of \lesssim we further have $2\mathbf{p}' \lesssim \mathbf{u}'$, so $\mathbf{p}' \in \mathfrak{T}_1(\mathbf{u}')$. It follows that $\mathbf{p}' \in \mathfrak{T}_2(\mathbf{u})$, which shows (2.0.33). \square

Lemma 5.4.6 (forest separation). *For each $\mathbf{u}, \mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $\mathbf{u} \neq \mathbf{u}'$ and each $\mathbf{p} \in \mathfrak{T}_2(\mathbf{u})$ with $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}')$ we have*

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) > 2^{Z(n+1)}. \quad (5.4.7)$$

Proof. By the definition (5.1.13) of $\mathfrak{C}_2(k, n, j)$, there exists a tile $\mathfrak{p}' \in \mathfrak{C}_1(k, n, j)$ with $\mathfrak{p}' \leq \mathfrak{p}$ and $s(\mathfrak{p}') \leq s(\mathfrak{p}) - Z(n+1)$. By Lemma 2.1.2 we have

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{u}')) \geq 2^{95aZ(n+1)} d_{\mathfrak{p}'}.(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{u}')).$$

By (5.3.3) we have $2\mathfrak{p}' \lesssim 2\mathfrak{p}$, so by transitivity of \lesssim there exists $\mathfrak{v} \sim \mathfrak{u}$ with $2\mathfrak{p}' \lesssim \mathfrak{v}$ and $\mathcal{J}(\mathfrak{p}') \neq \mathcal{J}(\mathfrak{v})$. Since $\mathfrak{u}, \mathfrak{u}'$ are not equivalent under \sim , we have $\mathfrak{v} \approx \mathfrak{u}'$, thus $10\mathfrak{p}' \not\lesssim \mathfrak{u}'$. This implies that there exists $q \in B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 1) \setminus B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), 10)$.

From $\mathfrak{p}' \leq \mathfrak{p}$, $\mathcal{J}(\mathfrak{p}') \subset \mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{u}')$ and Lemma 2.1.2 it then follows that

$$\begin{aligned} & d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{u}')) \\ & \geq -d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{p}')) + d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), q) - d_{\mathfrak{p}'}(q, \mathcal{Q}(\mathfrak{u}')) \\ & \geq -d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}), \mathcal{Q}(\mathfrak{p}')) + d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), q) - d_{\mathfrak{u}'}(q, \mathcal{Q}(\mathfrak{u}')) \\ & > -1 + 10 - 1 = 8. \end{aligned}$$

The lemma follows by combining the two displays with the fact that $95a \geq 1$. \square

Lemma 5.4.7 (forest inner). *For each $\mathfrak{u} \in \mathfrak{U}_3(k, n, j)$ and each $\mathfrak{p} \in \mathfrak{T}_2(\mathfrak{u})$ we have*

$$B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})}) \subset \mathcal{J}(\mathfrak{u}). \quad (5.4.8)$$

Proof. Let $\mathfrak{p} \in \mathfrak{T}_2(\mathfrak{u})$. Then $\mathfrak{p} \in \mathfrak{C}_4(k, n, j)$, hence there exists a chain

$$\mathfrak{p} \leq \mathfrak{p}_{Z(n+1)} \leq \cdots \leq \mathfrak{p}_0$$

of distinct tiles in $\mathfrak{C}_3(n, k, j)$. We pick such a chain and set $\mathfrak{q} = \mathfrak{p}_0$. Then we have from distinctness of the tiles in the chain that $s(\mathfrak{p}) \leq s(\mathfrak{q}) - Z(n+1)$. By (5.1.16) there exists $\mathfrak{u}'' \in \mathfrak{U}_1(k, n, j)$ with $2\mathfrak{q} \lesssim \mathfrak{u}''$ and $s(\mathfrak{q}) < s(\mathfrak{u}'')$. Then we have in particular by Lemma 5.3.1 that $10\mathfrak{p} \lesssim \mathfrak{u}''$. Let $\mathfrak{u}' \sim \mathfrak{u}$ be such that $\mathfrak{p} \in \mathfrak{T}_1(\mathfrak{u}')$. By the definition of \sim , it follows that $\mathfrak{u}' \sim \mathfrak{u}''$. By transitivity of \sim , we have $\mathfrak{u} \sim \mathfrak{u}''$. By Lemma 5.4.1, we have $s(\mathfrak{u}'') = s(\mathfrak{u})$, hence $s(\mathfrak{q}) < s(\mathfrak{u})$ and $s(\mathfrak{p}) \leq s(\mathfrak{q}) - Z(n+1) \leq s(\mathfrak{u}) - Z(n+1) - 1$.

Thus, there exists some cube $I \in \mathcal{D}$ with $s(I) = s(\mathfrak{u}) - Z(n+1) - 1$ and $I \subset \mathcal{J}(\mathfrak{u})$ and $\mathcal{J}(\mathfrak{p}) \subset I$. Since $\mathfrak{p} \in \mathfrak{C}_5(k, n, j)$, we have that $I \notin \mathcal{L}(\mathfrak{u})$, so $B(c(I), 8D^{s(I)}) \subset \mathcal{J}(\mathfrak{u})$. By the triangle inequality, (2.0.1) and $a \geq 4$, the same then holds for the subcube $\mathcal{J}(\mathfrak{p}) \subset I$. \square

Lemma 5.4.8 (forest stacking). *It holds for $k \leq n$ that*

$$\sum_{\mathfrak{u} \in \mathfrak{U}_3(k, n, j)} \mathbf{1}_{\mathcal{J}(\mathfrak{u})} \leq (4n+12)2^n. \quad (5.4.9)$$

Proof. Suppose that a point x is contained in more than $(4n+12)2^n$ cubes $\mathcal{J}(\mathfrak{u})$ with $\mathfrak{u} \in \mathfrak{U}_3(k, n, j)$. Since $\mathfrak{U}_3(k, n, j) \subset \mathfrak{C}_1(k, n, j)$ for each such \mathfrak{u} , there exists $\mathfrak{m} \in \mathfrak{M}(k, n)$ such that $100\mathfrak{u} \lesssim \mathfrak{m}$. We fix such an $\mathfrak{m}(\mathfrak{u}) := \mathfrak{m}$ for each \mathfrak{u} , and claim that the map $\mathfrak{u} \mapsto \mathfrak{m}(\mathfrak{u})$ is injective. Indeed, assume for $\mathfrak{u} \neq \mathfrak{u}'$ there is $\mathfrak{m} \in \mathfrak{M}(k, n)$ such that $100\mathfrak{u} \lesssim \mathfrak{m}$ and $100\mathfrak{u}' \lesssim \mathfrak{m}$. By (2.0.8), either $\mathcal{J}(\mathfrak{u}) \subset \mathcal{J}(\mathfrak{u}')$ or $\mathcal{J}(\mathfrak{u}') \subset \mathcal{J}(\mathfrak{u})$. By (5.1.14), $B_{\mathfrak{u}}(\mathcal{Q}(\mathfrak{u}), 100) \cap B_{\mathfrak{u}'}(\mathcal{Q}(\mathfrak{u}'), 100) = \emptyset$. This contradicts $\Omega(\mathfrak{m})$ being contained in both sets by (2.0.15). Thus x is contained in more than $(4n+12)2^n$ cubes $\mathcal{J}(\mathfrak{m})$, $\mathfrak{m} \in \mathfrak{M}(k, n)$. Consequently, we have by (5.1.26) that $x \in A(2n+6, k, n) \subset G_2$. Let $\mathcal{J}(\mathfrak{u})$ be an inclusion minimal cube among the

$\mathcal{J}(\mathbf{u}'), \mathbf{u}' \in \mathfrak{U}_3(k, n, j)$ with $x \in \mathcal{J}(\mathbf{u})$. By the dyadic property (2.0.8), we have $\mathcal{J}(\mathbf{u}) \subset \mathcal{J}(\mathbf{u}')$ for all cubes $\mathcal{J}(\mathbf{u}')$ containing x . Thus

$$\mathcal{J}(\mathbf{u}) \subset \{y : \sum_{\mathbf{u} \in \mathfrak{U}_3(k, n, j)} \mathbf{1}_{\mathcal{J}(\mathbf{u})}(y) > 1 + (4n + 12)2^n\} \subset G_2.$$

Thus $\mathfrak{T}_1(\mathbf{u}) \cap \mathfrak{C}_6(k, n, j) = \emptyset$. This contradicts $\mathbf{u} \in \mathfrak{U}_2(k, n, j)$. \square

We now turn to the proof of Lemma 5.1.2.

Proof of Lemma 5.1.2. We first fix k, n, j . By (2.0.21) and (2.0.20), we have that $\mathbf{1}_{\mathcal{J}(\mathbf{p})} T_{\mathbf{p}} f(x) = T_{\mathbf{p}} f(x)$ and hence $\mathbf{1}_{G \setminus G'} T_{\mathbf{p}} f(x) = 0$ for all $\mathbf{p} \in \mathfrak{C}_5(k, n, j) \setminus \mathfrak{C}_6(k, n, j)$. Thus it suffices to estimate the contribution of the sets $\mathfrak{C}_6(k, n, j)$. By Lemma 5.4.8, we can decompose $\mathfrak{U}_3(k, n, j)$ as a disjoint union of at most $4n + 12$ collections $\mathfrak{U}_4(k, n, j, l)$, $1 \leq l \leq 4n + 12$, each satisfying

$$\sum_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathbf{1}_{\mathcal{J}(\mathbf{u})} \leq 2^n.$$

By Lemmas 5.4.4, 5.4.5, 5.4.6, 5.4.7 and 5.3.12, the pairs

$$(\mathfrak{U}_4(k, n, j, l), \mathfrak{T}_2|_{\mathfrak{U}_4(k, n, j, l)})$$

are n -forests for each k, n, j, l , and by Lemma 5.4.3, we have

$$\mathfrak{C}_6(k, n, j) = \bigcup_{l=1}^{4n+12} \bigcup_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathfrak{T}_2(\mathbf{u}).$$

Since $\mathcal{J}(\mathbf{p}) \not\subset G_1$ for all $\mathbf{p} \in \mathfrak{C}_6(k, n, j)$, we have $\mathfrak{C}_6(k, n, j) \cap \mathfrak{P}_{F, G} = \emptyset$ and hence

$$\text{dens}_2\left(\bigcup_{\mathbf{u} \in \mathfrak{U}_4(k, n, j, l)} \mathfrak{T}_2(\mathbf{u})\right) \leq 2^{2a+5} \frac{\mu(F)}{\mu(G)}.$$

Using the triangle inequality according to the splitting by k, n, j and l in (5.1.31) and applying Proposition 2.0.4 to each term, we obtain the estimate

$$\sum_{k \geq 0} \sum_{n \geq k} (2n + 3)(4n + 12) 2^{440a^3} 2^{-(1-\frac{1}{q})n} (2^{2a+5} \frac{\mu(F)}{\mu(G)})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|\mathbf{1}_{G \setminus G'}\|_2$$

for the left hand side of (5.1.31). Since $|f| \leq \mathbf{1}_F$, we have $\|f\|_2 \leq \mu(F)^{1/2}$, and we have $\|\mathbf{1}_{G \setminus G'}\|_2 \leq \mu(G)^{1/2}$. We get a bound

$$2^{440a^3} \mu(F)^{\frac{1}{q}} \mu(G)^{1-\frac{1}{q}} 2^{2a+5} \sum_{k \geq 0} \sum_{n \geq k} (2n + 3)(4n + 12) 2^{-(1-\frac{1}{q})n}.$$

Interchanging the order of summation, the last factor equals

$$2^{2a+5} \sum_{n \geq 0} (2n + 3)(4n + 12)(n + 1) 2^{-\frac{q-1}{q}n}.$$

Up to an explicit constant, the sum is bounded by $\sum n^3 2^{-\frac{q-1}{q}n}$, which is at most some constant times $1/(q-1)^4$ by comparing to an integral. Since $a \geq 4$, this is overall bounded by $2^{a^3}/(q-1)^4$, which completes the proof of the lemma. \square

5.5. Proof of the Forest Complement Lemma. Define $\mathfrak{P}_{G \setminus G'}$ to be the set of all $\mathfrak{p} \in \mathfrak{P}$ such that $\mu(\mathcal{J}(\mathfrak{p}) \cap (G \setminus G')) > 0$.

Lemma 5.5.1 (antichain decomposition). *We have that*

$$\mathfrak{P}_2 \cap \mathfrak{P}_{G \setminus G'} \quad (5.5.1)$$

$$= \bigcup_{k \geq 0} \bigcup_{n \geq k} \mathfrak{L}_0(k, n) \cap \mathfrak{P}_{G \setminus G'} \quad (5.5.2)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \mathfrak{L}_2(k, n, j) \cap \mathfrak{P}_{G \setminus G'} \quad (5.5.3)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_1(k, n, j, l) \cap \mathfrak{P}_{G \setminus G'} \quad (5.5.4)$$

$$\cup \bigcup_{k \geq 0} \bigcup_{n \geq k} \bigcup_{0 \leq j \leq 2n+3} \bigcup_{0 \leq l \leq Z(n+1)} \mathfrak{L}_3(k, n, j, l) \cap \mathfrak{P}_{G \setminus G'}. \quad (5.5.5)$$

Proof. Let $\mathfrak{p} \in \mathfrak{P}_2 \cap \mathfrak{P}_{G \setminus G'}$. Clearly, for every cube $J = \mathcal{J}(\mathfrak{p})$ with $\mathfrak{p} \in \mathfrak{P}_{G \setminus G'}$ there exists some $k \geq 0$ such that (5.1.1) holds, and for no cube $J \in \mathcal{D}$ and no $k < 0$ does (5.1.2) hold. Thus $\mathfrak{p} \in \mathfrak{P}(k)$ for some $k \geq 0$.

Next, since $E_2(\lambda, \mathfrak{p}') \subset \mathcal{J}(\mathfrak{p}') \cap G$ for every $\lambda \geq 2$ and every tile $\mathfrak{p}' \in \mathfrak{P}(k)$ with $\lambda \mathfrak{p} \lesssim \lambda \mathfrak{p}'$, it follows from (5.1.2) that $\mu(E_2(\lambda, \mathfrak{p}')) \leq 2^{-k} \mu(\mathcal{J}(\mathfrak{p}'))$ for every such \mathfrak{p}' , so $\text{dens}'_k(\{\mathfrak{p}\}) \leq 2^{-k}$. Combining this with $a \geq 0$, it follows from (5.1.7) that there exists $n \geq k$ with $\mathfrak{p} \in \mathfrak{C}(k, n)$.

Since $\mathfrak{p} \in \mathfrak{P}_{G \setminus G'}$, we have in particular $\mathcal{J}(\mathfrak{p}) \not\subset A(2n+6, k, n)$, so there exist at most $1 + (4n+12)2^n < 2^{2n+4}$ tiles $\mathfrak{m} \in \mathfrak{M}(k, n)$ with $\mathfrak{p} \leq \mathfrak{m}$. It follows that $\mathfrak{p} \in \mathfrak{L}_0(k, n)$ or $\mathfrak{p} \in \mathfrak{C}_1(k, n, j)$ for some $1 \leq j \leq 2n+3$. In the former case we are done, in the latter case the equality to be shown follows from the definitions of the collections \mathfrak{C}_i and \mathfrak{L}_i . \square

Lemma 5.5.2 (L0 antichain). *We have that*

$$\mathfrak{L}_0(k, n) = \bigcup_{0 \leq l < n} \mathfrak{L}_0(k, n, l),$$

where each $\mathfrak{L}_0(k, n, l)$ is an antichain.

Proof. It suffices to show that $\mathfrak{L}_0(k, n)$ contains no chain of length $n+1$. Suppose that we had such a chain $\mathfrak{p}_0 \leq \mathfrak{p}_1 \leq \dots \leq \mathfrak{p}_n$ with $\mathfrak{p}_i \neq \mathfrak{p}_{i+1}$ for $i = 0, \dots, n-1$. By (5.1.7), we have that $\text{dens}'_k(\{\mathfrak{p}_n\}) > 2^{-n}$. Thus, by (5.1.6), there exists $\mathfrak{p}' \in \mathfrak{P}(k)$ and $\lambda \geq 2$ with $\lambda \mathfrak{p}_n \leq \lambda \mathfrak{p}'$ and

$$\frac{\mu(E_2(\lambda, \mathfrak{p}'))}{\mu(\mathcal{J}(\mathfrak{p}'))} > \lambda^a 2^{4a} 2^{-n}. \quad (5.5.6)$$

Let \mathfrak{D} be the set of all $\mathfrak{p}'' \in \mathfrak{P}(k)$ such that we have $\mathcal{J}(\mathfrak{p}'') = \mathcal{J}(\mathfrak{p}')$ and $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \lambda) \cap \Omega(\mathfrak{p}'') \neq \emptyset$. We now show that

$$|\mathfrak{D}| \leq 2^{4a} \lambda^a. \quad (5.5.7)$$

The balls $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}''), 0.2)$, $\mathfrak{p}'' \in \mathfrak{D}$ are disjoint by (2.0.15), and by the triangle inequality contained in $B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \lambda + 1.2)$. By assumption (1.1.8) on Θ , this ball can

be covered with

$$2^{a\lceil \log_2(\lambda+1.2)+\log_2(5) \rceil} \leq 2^{a(\log_2(\lambda)+4)} = 2^{4a}\lambda^a$$

many $d_{\mathbf{p}'}$ -balls of radius 0.2. Here we have used that for $\lambda \geq 2$

$$\lceil \log_2(\lambda + 1.2) + \log_2(5) \rceil \leq 1 + \log_2(1.6\lambda) + \log_2(5) = 4 + \log_2(\lambda).$$

By the triangle inequality, each such ball contains at most one $\mathcal{Q}(\mathbf{p}'')$, and each $\mathcal{Q}(\mathbf{p}'')$ is contained in one of the balls. Thus we get (5.5.7).

By (2.0.26) and (2.0.27) we have $E_2(\lambda, \mathbf{p}') \subset \bigcup_{\mathbf{p}'' \in \mathfrak{D}} E_1(\mathbf{p}'')$, thus

$$2^{4a}\lambda^a 2^{-n} < \sum_{\mathbf{p}'' \in \mathfrak{D}} \frac{\mu(E_1(\mathbf{p}''))}{\mu(\mathcal{J}(\mathbf{p}''))}.$$

Hence there exists a tile $\mathbf{p}'' \in \mathfrak{D}$ with

$$\mu(E_1(\mathbf{p}'')) \geq 2^{-n}\mu(\mathcal{J}(\mathbf{p}')).$$

By the definition (5.1.5) of $\mathfrak{M}(k, n)$, there exists a tile $\mathbf{m} \in \mathfrak{M}(k, n)$ with $\mathbf{p}' \leq \mathbf{m}$. From (5.5.6), the inclusion $E_2(\lambda, \mathbf{p}') \subset \mathcal{J}(\mathbf{p}')$ and $a \geq 1$ we obtain

$$2^n \geq 2^{4a}\lambda^a \geq \lambda.$$

From the triangle inequality, Lemma 2.1.2 and $a \geq 1$, we now obtain for all $\vartheta \in B_{\mathbf{m}}(\mathcal{Q}(\mathbf{m}), 1)$ that

$$\begin{aligned} & d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_0), \vartheta) \\ & \leq d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_0), \mathcal{Q}(\mathbf{p}_n)) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}_n), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p}'')) \\ & \quad + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{p}''), \mathcal{Q}(\mathbf{m})) + d_{\mathbf{p}_0}(\mathcal{Q}(\mathbf{m}), \vartheta) \\ & \leq 1 + 2^{-95an}(d_{\mathbf{p}_n}(\mathcal{Q}(\mathbf{p}_n), \mathcal{Q}(\mathbf{p}')) + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p}''))) \\ & \quad + d_{\mathbf{p}''}(\mathcal{Q}(\mathbf{p}''), \mathcal{Q}(\mathbf{m})) + d_{\mathbf{m}}(\mathcal{Q}(\mathbf{m}), \vartheta) \\ & \leq 1 + 2^{-95an}(\lambda + (\lambda + 1) + 1 + 1) \leq 100. \end{aligned}$$

Thus, by (2.0.23), $100\mathbf{p}_0 \lesssim \mathbf{m}$, a contradiction to $\mathbf{p}_0 \notin \mathfrak{C}(k, n)$. \square

Lemma 5.5.3 (L2 antichain). *Each of the sets $\mathfrak{L}_2(k, n, j)$ is an antichain.*

Proof. Suppose that there are $\mathbf{p}_0, \mathbf{p}_1 \in \mathfrak{L}_2(k, n, j)$ with $\mathbf{p}_0 \neq \mathbf{p}_1$ and $\mathbf{p}_0 \leq \mathbf{p}_1$. By Lemma 5.3.1 and Lemma 5.3.2, it follows that $2\mathbf{p}_0 \lesssim 200\mathbf{p}_1$. Since $\mathfrak{L}_2(k, n, j)$ is finite, there exists a maximal $l \geq 1$ such that there exists a chain $2\mathbf{p}_0 \lesssim 200\mathbf{p}_1 \lesssim \dots \lesssim 200\mathbf{p}_l$ with all \mathbf{p}_i in $\mathfrak{C}_1(k, n, j)$ and $\mathbf{p}_i \neq \mathbf{p}_{i+1}$ for $i = 0, \dots, l-1$. If we have $\mathbf{p}_l \in \mathfrak{U}_1(k, n, j)$, then it follows from $2\mathbf{p}_0 \lesssim 200\mathbf{p}_l \lesssim \mathbf{p}_l$ and (5.1.15) that $\mathbf{p}_0 \notin \mathfrak{L}_2(k, n, j)$, a contradiction. Thus, by the definition (5.1.14) of $\mathfrak{U}_1(k, n, j)$, there exists $\mathbf{p}_{l+1} \in \mathfrak{C}_1(k, n, j)$ with $\mathcal{J}(\mathbf{p}_l) \subsetneq \mathcal{J}(\mathbf{p}_{l+1})$ and $\vartheta \in B_{\mathbf{p}_l}(\mathcal{Q}(\mathbf{p}_l), 100) \cap B_{\mathbf{p}_{l+1}}(\mathcal{Q}(\mathbf{p}_{l+1}), 100)$. Using the triangle inequality and Lemma 2.1.2, one deduces that $200\mathbf{p}_l \lesssim 200\mathbf{p}_{l+1}$. This contradicts maximality of l . \square

Lemma 5.5.4 (L1 L3 antichain). *Each of the sets $\mathfrak{L}_1(k, n, j, l)$ and $\mathfrak{L}_3(k, n, j, l)$ is an antichain.*

Proof. By its definition (5.1.11), each set $\mathfrak{L}_1(k, n, j, l)$ is a set of minimal elements in some set of tiles with respect to \leq . If there were distinct $\mathfrak{p}, \mathfrak{q} \in \mathfrak{L}_1(k, n, j, l)$ with $\mathfrak{p} \leq \mathfrak{q}$, then \mathfrak{q} would not be minimal. Hence such $\mathfrak{p}, \mathfrak{q}$ do not exist. Similarly, by (5.1.17), each set $\mathfrak{L}_3(k, n, j, l)$ is a set of maximal elements in some set of tiles with respect to \leq . If there were distinct $\mathfrak{p}, \mathfrak{q} \in \mathfrak{L}_3(k, n, j, l)$ with $\mathfrak{p} \leq \mathfrak{q}$, then \mathfrak{p} would not be maximal. \square

We now turn to the proof of Lemma 5.1.3.

Proof of Lemma 5.1.3. If $\mathfrak{p} \notin \mathfrak{P}_{G \setminus G'}$, then $\mu(\mathcal{J}(\mathfrak{p}) \cap (G \setminus G')) = 0$. By (2.0.21) and (2.0.26), it follows that $\mathbf{1}_{G \setminus G'} T_{\mathfrak{p}} f(x) = 0$ for almost every x . We thus have, almost everywhere,

$$\mathbf{1}_{G \setminus G'} \sum_{\mathfrak{p} \in \mathfrak{P}_2} T_{\mathfrak{p}} f(x) = \mathbf{1}_{G \setminus G'} \sum_{\mathfrak{p} \in \mathfrak{P}_2 \cap \mathfrak{P}_{G \setminus G'}} T_{\mathfrak{p}} f(x).$$

Let $\mathfrak{L}(k, n)$ denote any of the terms $\mathfrak{L}_i(k, n, j, l) \cap \mathfrak{P}_{G \setminus G'}$ on the right hand side of (5.5.1), where the indices j, l may be void. Then $\mathfrak{L}(k, n)$ is an antichain, by Lemmas 5.5.2, 5.5.3, 5.5.4. Further, we have

$$\text{dens}_1(\mathfrak{L}(k, n)) \leq 2^{4a+1-n}$$

by Lemma 5.3.12, and we have

$$\text{dens}_2(\mathfrak{L}(k, n)) \leq 2^{2a+5} \frac{\mu(F)}{\mu(G)},$$

since

$$\mathfrak{L}(k, n) \cap \mathfrak{P}_{F, G} \subset \mathfrak{P}_{G \setminus G'} \cap \mathfrak{P}_{F, G} = \emptyset.$$

Applying now the triangle inequality according to the decomposition coming from Lemma 5.5.1, and then applying Proposition 2.0.3 to each term, we obtain the estimate

$$\begin{aligned} &\leq \sum_{k \geq 0} \sum_{n \geq k} (n + (2n + 4) + 2(2n + 4)(1 + Z(n + 1))) \\ &\quad \times 2^{117a^3} (q - 1)^{-1} (2^{4a+1-n})^{\frac{q-1}{8a^4}} (2^{2a+5} \frac{\mu(F)}{\mu(G)})^{\frac{1}{q} - \frac{1}{2}} \|f\|_2 \|\mathbf{1}_{G \setminus G'}\|_2. \end{aligned}$$

Because $|f| \leq \mathbf{1}_F$, we have $\|f\|_2 \leq \mu(F)^{1/2}$, and we have $\|\mathbf{1}_{G \setminus G'}\|_2 \leq \mu(G)^{1/2}$. Using this and (2.0.3), we bound

$$\leq 2^{118a^3} (q - 1)^{-1} \mu(F)^{\frac{1}{q}} \mu(G)^{\frac{1}{q}} \sum_{k \geq 0} \sum_{n \geq k} n^2 2^{-n \frac{q-1}{8a^4}}.$$

The last sum equals, by changing the order of summation,

$$\sum_{n \geq 0} n^2 (n + 1) 2^{-n \frac{q-1}{8a^4}} \leq \frac{2^{2a^3}}{(q - 1)^4}.$$

This completes the proof. \square

6. PROOF OF THE ANTICHAIN OPERATOR PROPOSITION

Let an antichain \mathfrak{A} and functions f, g as in Proposition 2.0.3 be given. We prove (2.0.30) in Section 6.1 as the geometric mean of two inequalities, each involving one of the two densities. One of these two inequalities will need a careful estimate formulated in Lemma 6.1.5 of the TT^* correlation between two tile operators. Lemma 6.1.5 will be proven in Section 6.2.

The summation of the contributions of these individual correlations will require a geometric Lemma 6.1.6 counting the relevant tile pairs. Lemma 6.1.6 will be proven in Subsection 6.3.

6.1. The density arguments. We begin with the following crucial disjointness property of the sets $E(\mathfrak{p})$ with $\mathfrak{p} \in \mathfrak{A}$.

Lemma 6.1.1 (tile disjointness). *Let $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{A}$. If there exists an $x \in X$ with $x \in E(\mathfrak{p}) \cap E(\mathfrak{p}')$, then $\mathfrak{p} = \mathfrak{p}'$.*

Proof. Let $\mathfrak{p}, \mathfrak{p}'$ and x be given. Assume without loss of generality that $s(\mathfrak{p}) \leq s(\mathfrak{p}')$. As we have $x \in E(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p})$ and $x \in E(\mathfrak{p}') \subset \mathcal{J}(\mathfrak{p}')$ by Definition (2.0.20), we conclude for $i = 1, 2$ that $Q(x) \in \Omega(\mathfrak{p})$ and $Q(x) \in \Omega(\mathfrak{p}')$. By (2.0.14) we have $\Omega(\mathfrak{p}') \subset \Omega(\mathfrak{p})$. By Definition (2.0.23), we conclude $\mathfrak{p} \leq \mathfrak{p}'$. As \mathfrak{A} is an antichain, we conclude $\mathfrak{p} = \mathfrak{p}'$. This proves the lemma. \square

Let \mathcal{B} be the collection of balls

$$B(c(\mathfrak{p}), 8D^{s(\mathfrak{p})}) \quad (6.1.1)$$

with $\mathfrak{p} \in \mathfrak{A}$ and recall the definition of $M_{\mathcal{B}}$ from Definition 2.0.41.

Lemma 6.1.2 (maximal bound antichain). *Let $x \in X$. Then*

$$\left| \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) \right| \leq 2^{102a^3} M_{\mathcal{B}} f(x). \quad (6.1.2)$$

Proof. Fix $x \in X$. By Lemma 6.1.1, there is at most one $\mathfrak{p} \in \mathfrak{A}$ such that $T_{\mathfrak{p}} f(x)$ is not zero. If there is no such \mathfrak{p} , the estimate (6.1.2) follows.

Assume there is such a \mathfrak{p} . By definition of $T_{\mathfrak{p}}$ we have $x \in E(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p})$ and by the squeezing property (2.0.10)

$$\rho(x, c(\mathfrak{p})) \leq 4D^{s(\mathfrak{p})}. \quad (6.1.3)$$

Let $y \in X$ with $K_{s(\mathfrak{p})}(x, y) \neq 0$. By Definition (2.0.5) of $K_{s(\mathfrak{p})}$ we have

$$\frac{1}{4}D^{s(\mathfrak{p})-1} \leq \rho(x, y) \leq \frac{1}{2}D^{s(\mathfrak{p})}. \quad (6.1.4)$$

The triangle inequality with (6.1.3) and (6.1.4) implies

$$\rho(c(\mathfrak{p}), y) < 8D^{s(\mathfrak{p})}. \quad (6.1.5)$$

Using the kernel bound (1.1.11) and the lower bound in (2.1.2) we obtain

$$|K_{s(\mathfrak{p})}(x, y)| \leq \frac{2^{a^3}}{\mu(B(x, \frac{1}{4}D^{s(\mathfrak{p})-1}))}. \quad (6.1.6)$$

Using $D = 2^{100a^2}$ and the doubling property (1.1.2) $5 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{5a+101a^3}}{\mu(B(x, 8D^s(\mathfrak{p})))} \quad (6.1.7)$$

which, thanks to the closeness of the points x and $c(\mathfrak{p})$ shown in (6.1.3), is in turn bounded by

$$\leq \frac{2^{6a+101a^3}}{\mu(B(c(\mathfrak{p}), 8D^s(\mathfrak{p})))}. \quad (6.1.8)$$

Using that $|e(\vartheta)|$ is bounded by 1 for every $\vartheta \in \Theta$, we estimate with the triangle inequality and the above information

$$|T_{\mathfrak{p}}f(x)| \leq \frac{2^{6a+101a^3}}{\mu(B(c(\mathfrak{p}), 8D^s(\mathfrak{p})))} \int_{\mu(B(c(\mathfrak{p}), 8D^s(\mathfrak{p})))} |f(y)| dy \quad (6.1.9)$$

This together with $a \geq 4$ proves the Lemma. \square

Set

$$\tilde{q} = \frac{2q}{1+q}. \quad (6.1.10)$$

Since $1 < q \leq 2$, we have $1 < \tilde{q} < q \leq 2$.

Lemma 6.1.3 (dens2 antichain). *We have that*

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}}f(x) d\mu(x) \right| \leq 2^{103a^3} (q-1)^{-1} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}} \|f\|_2 \|g\|_2. \quad (6.1.11)$$

Proof. We have $f = \mathbf{1}_F f$. Using Hölder's inequality, we obtain for each $x \in B'$ and each $B' \in \mathcal{B}$ using $1 < \tilde{q} \leq 2$

$$\frac{1}{\mu(B')} \int_{B'} |f(y)| d\mu(y) \quad (6.1.12)$$

$$\leq \left(\frac{1}{\mu(B')} \int_{B'} |f(y)|^{\frac{2\tilde{q}}{3\tilde{q}-2}} d\mu(y) \right)^{\frac{3}{2}-\frac{1}{\tilde{q}}} \left(\frac{1}{\mu(B')} \int_{B'} \mathbf{1}_F(y) d\mu(y) \right)^{\frac{1}{\tilde{q}}-\frac{1}{2}} \quad (6.1.13)$$

$$\leq \left(M_{\mathcal{B}}(|f|^{\frac{2\tilde{q}}{3\tilde{q}-2}})(x) \right)^{\frac{3}{2}-\frac{1}{\tilde{q}}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}}. \quad (6.1.14)$$

Taking the maximum over all B' containing x , we obtain

$$M_{\mathcal{B}}|f| \leq M_{\mathcal{B}, \frac{2\tilde{q}}{3\tilde{q}-2}} |f| \text{dens}_2(\mathfrak{A})^{\frac{1}{q}-\frac{1}{2}}. \quad (6.1.15)$$

We have with Proposition 2.0.6

$$\left\| M_{\mathcal{B}, \frac{2\tilde{q}}{3\tilde{q}-2}} f \right\|_2 \leq 2^{2a} (3\tilde{q}-2)(2\tilde{q}-2)^{-1} \|f\|_2. \quad (6.1.16)$$

Using $1 < \tilde{q} \leq 2$ estimates the last display by

$$2^{2a+2} (\tilde{q}-1)^{-1} \|f\|_2. \quad (6.1.17)$$

We obtain with Cauchy-Schwarz and then Lemma 6.1.2

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x) \right| \quad (6.1.18)$$

$$\leq \|g\|_2 \left\| \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f \right\|_2 \quad (6.1.19)$$

$$\leq 2^{102a^3} \|g\|_2 \|M_{\mathcal{B}} f\|_2 \quad (6.1.20)$$

With (6.1.15) and (6.1.17) we can estimate the last display by

$$\leq 2^{102a^3+2a+2} (\tilde{q}-1)^{-1} \|g\|_2 \|f\|_2 \text{dens}_2(\mathfrak{A})^{\frac{1}{\tilde{q}}-\frac{1}{2}} \quad (6.1.21)$$

Using $a \geq 4$ and $(\tilde{q}-1)^{-1} = (q+1)/(q-1) \leq 3(q-1)^{-1}$ proves the lemma. \square

Lemma 6.1.4 (dens1 antichain). *Set $p := 4a^4$. We have*

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x) \right| \leq 2^{117a^3} \text{dens}_1(\mathfrak{A})^{\frac{1}{2p}} \|f\|_2 \|g\|_2. \quad (6.1.22)$$

Proof. We write for the expression inside the absolute values on the left-hand side of (6.1.22)

$$\sum_{\mathfrak{p} \in \mathfrak{A}} \iint \overline{g(x)} \mathbf{1}_{E(\mathfrak{p})}(x) K_{s(\mathfrak{p})}(x, y) e(Q(x)(y) - Q(x)(x)) f(y) d\mu(y) d\mu(x) \quad (6.1.23)$$

$$= \int \sum_{\mathfrak{p} \in \mathfrak{A}} \overline{T_{\mathfrak{p}}^* g(y)} f(y) d\mu(y) \quad (6.1.24)$$

with the adjoint operator

$$T_{\mathfrak{p}}^* g(y) = \int_{E(\mathfrak{p})} \overline{K_{s(\mathfrak{p})}(x, y)} e(-Q(x)(y) + Q(x)(x)) g(x) d\mu(x). \quad (6.1.25)$$

We have by expanding the square

$$\int \left| \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}}^* g(y) \right|^2 d\mu(y) = \int \left(\sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}}^* g(y) \right) \left(\sum_{\mathfrak{p}' \in \mathfrak{A}} \overline{T_{\mathfrak{p}'}^* g(y)} \right) d\mu(y) \quad (6.1.26)$$

$$\leq \sum_{\mathfrak{p} \in \mathfrak{A}} \sum_{\mathfrak{p}' \in \mathfrak{A}} \left| \int T_{\mathfrak{p}}^* g(y) \overline{T_{\mathfrak{p}'}^* g(y)} d\mu(y) \right|. \quad (6.1.27)$$

We split the sum into the terms with $s(\mathfrak{p}') \leq s(\mathfrak{p})$ and $s(\mathfrak{p}) < s(\mathfrak{p}')$. Using the symmetry of each summand, we may switch \mathfrak{p} and \mathfrak{p}' in the second sum. Using further positivity of each summand to replace the condition $s(\mathfrak{p}') < s(\mathfrak{p})$ by $s(\mathfrak{p}') \leq s(\mathfrak{p})$ in the second sum, we estimate (6.1.27) by

$$\leq 2 \sum_{\mathfrak{p} \in \mathfrak{A}} \sum_{\mathfrak{p}' \in \mathfrak{A} : s(\mathfrak{p}') \leq s(\mathfrak{p})} \left| \int T_{\mathfrak{p}}^* g(y) \overline{T_{\mathfrak{p}'}^* g(y)} d\mu(y) \right|. \quad (6.1.28)$$

Define for $\mathfrak{p} \in \mathfrak{P}$

$$B(\mathfrak{p}) := B(c(\mathfrak{p}), 14D^{s(\mathfrak{p})}) \quad (6.1.29)$$

and define

$$\mathfrak{A}(\mathfrak{p}) := \{\mathfrak{p}' \in \mathfrak{A} : s(\mathfrak{p}') \leq s(\mathfrak{p}) \wedge \mathcal{J}(\mathfrak{p}') \subset B(\mathfrak{p})\}. \quad (6.1.30)$$

Note that by the squeezing property (2.0.10) and the doubling property (1.1.2) applied 6 times we have

$$\mu(B(\mathbf{p})) \leq 2^{6a} \mu(\mathcal{J}(\mathbf{p})). \quad (6.1.31)$$

Using Lemma 6.1.5 and (6.1.31), we estimate (6.1.28) by

$$\leq 2^{232a^3+6a+1} \sum_{\mathbf{p} \in \mathfrak{A}} \int_{E(\mathbf{p})} |g|(y) h(\mathbf{p}) d\mu(y) \quad (6.1.32)$$

with $h(\mathbf{p})$ defined as

$$\frac{1}{\mu(B(\mathbf{p}))} \int \sum_{\mathbf{p}' \in \mathfrak{A}(\mathbf{p})} (1 + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p})))^{-1/(2a^2+a^3)} (\mathbf{1}_{E(\mathbf{p}')} |g|)(y') d\mu(y'). \quad (6.1.33)$$

Note that $p > 4$ since $a \geq 4$. Let p' be the dual exponent of p , satisfying $1/p + 1/p' = 1$. We estimate $h(\mathbf{p})$ as defined in (6.1.33) with Hölder using $|g| \leq \mathbf{1}_G$ and $E(\mathbf{p}') \subset B(\mathbf{p})$ by

$$\frac{\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'}}{\mu(B(\mathbf{p}))} \left\| \sum_{\mathbf{p} \in \mathfrak{A}(\mathbf{p})} (1 + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{p}')))^{-1/(2a^2+a^3)} \mathbf{1}_{E(\mathbf{p})} \mathbf{1}_G \right\|_p. \quad (6.1.34)$$

Then we apply Lemma 6.1.6 to estimate this by

$$\leq 2^{5a} \frac{\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'}}{\mu(B(\mathbf{p}))} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(B(\mathbf{p}))^{\frac{1}{p}}. \quad (6.1.35)$$

Let \mathcal{B}' be the collection of all balls $B(\mathbf{p})$ with $\mathbf{p} \in \mathfrak{A}$. Then for each $\mathbf{p} \in \mathfrak{A}$ and $x \in B(\mathbf{p})$ we have by definition (2.0.41) of $M_{\mathcal{B}', p'}$

$$\|g \mathbf{1}_{B(\mathbf{p})}\|_{p'} \leq \mu(B(\mathbf{p}))^{\frac{1}{p'}} M_{\mathcal{B}', p'} g(x). \quad (6.1.36)$$

Hence we can estimate (6.1.35) by

$$\leq 2^{5a} (M_{\mathcal{B}', p'} g(x)) \text{dens}_1(\mathfrak{A})^{\frac{1}{p}}. \quad (6.1.37)$$

With this estimate of $h(\mathbf{p})$, using $E(\mathbf{p}) \subset B(\mathbf{p})$ by construction of $B(\mathbf{p})$, we estimate (6.1.32) by

$$\leq 2^{232a^3+11a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \sum_{\mathbf{p} \in \mathfrak{A}} \int_{E(\mathbf{p})} |g|(y) M_{\mathcal{B}', p'} g(y) dy. \quad (6.1.38)$$

Using Lemma 6.1.1, the last display is observed to be

$$\leq 2^{232a^3+11a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \int |g|(y) (M_{\mathcal{B}', p'} g)(y) dy. \quad (6.1.39)$$

Applying Cauchy-Schwarz and using Proposition 2.0.6 and $1 < p' < \frac{3}{2}$ estimates the last display by

$$\leq 2^{232a^3+11a+1} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \|g\|_2 \|M_{\mathcal{B}', p'} g\|_2 \quad (6.1.40)$$

$$\leq 2^{232a^3+12a+3} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \|g\|_2^2. \quad (6.1.41)$$

Now Lemma 6.1.4 follows by applying Cauchy-Schwarz on the left-hand side and using $a \geq 4$. \square

The following basic TT^* estimate will be proved in Section 6.2.

Lemma 6.1.5 (tile correlation). *Let $\mathfrak{p}, \mathfrak{p}' \in \mathfrak{P}$ with $s(\mathfrak{p}') \leq s(\mathfrak{p})$. Then*

$$\left| \int T_{\mathfrak{p}'}^* g \overline{T_{\mathfrak{p}}^* g} \right| \quad (6.1.42)$$

$$\leq 2^{232a^3} \frac{(1 + d_{\mathfrak{p}'}(Q(\mathfrak{p}'), Q(\mathfrak{p})))^{-1/(2a^2+a^3)}}{\mu(\mathcal{J}(\mathfrak{p}))} \int_{E(\mathfrak{p}')} |g| \int_{E(\mathfrak{p})} |g|. \quad (6.1.43)$$

Moreover, the term (6.1.42) vanishes unless

$$\mathcal{J}(\mathfrak{p}') \subset B(c(\mathfrak{p}), 14D^{s(\mathfrak{p})}). \quad (6.1.44)$$

The following lemma will be proved in Section 6.3.

Lemma 6.1.6 (antichain tile count). *Set $p := 4a^4$. For every $\vartheta \in \Theta$ and every antichain \mathfrak{A} we have*

$$\left\| \sum_{\mathfrak{p} \in \mathfrak{A}} (1 + d_{\mathfrak{p}}(Q(\mathfrak{p}), \vartheta))^{-1/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})} \mathbf{1}_G \right\|_p \quad (6.1.45)$$

$$\leq 2^{5a} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu(\cup_{\mathfrak{p} \in \mathfrak{A}} I_{\mathfrak{p}})^{\frac{1}{p}}. \quad (6.1.46)$$

From these lemmas it is easy to prove Proposition 2.0.3.

Proof of Proposition 2.0.3. We have

$$\left(\frac{1}{\tilde{q}} - \frac{1}{2} \right) (2 - q) = \frac{1}{q} - \frac{1}{2}. \quad (6.1.47)$$

Multiplying the $(2 - q)$ -th power of (6.1.11) and the $(q - 1)$ -th power of (6.1.22) and estimating gives after simplification of some factors

$$\left| \int \overline{g(x)} \sum_{\mathfrak{p} \in \mathfrak{A}} T_{\mathfrak{p}} f(x) d\mu(x) \right| \quad (6.1.48)$$

$$\leq 2^{117a^3} (q - 1)^{-1} \text{dens}_1(\mathfrak{A})^{\frac{q-1}{2p}} \text{dens}_2(\mathfrak{A})^{\frac{1}{q} - \frac{1}{2}} \|f\|_2 \|g\|_2. \quad (6.1.49)$$

With the definition of p , this implies Proposition 2.0.3. \square

6.2. Proof of the Tile Correlation Lemma. The next lemma prepares an application of Proposition 2.0.5.

Lemma 6.2.1 (correlation kernel bound). *Let $-S \leq s_1 \leq s_2 \leq S$ and let $x_1, x_2 \in X$. Define*

$$\varphi(y) := \overline{K_{s_1}(x_1, y)} K_{s_2}(x_2, y). \quad (6.2.1)$$

If $\varphi(y) \neq 0$, then

$$y \in B(x_1, D^{s_1}). \quad (6.2.2)$$

Moreover, we have with $\tau = 1/a$

$$\|\varphi\|_{C^\tau(B(x_1, 2D^{s_1}))} \leq \frac{2^{231a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))}. \quad (6.2.3)$$

Proof. If $\varphi(y)$ is not zero, then $K_{s_1}(x_1, y)$ is not zero and thus (2.1.2) gives (6.2.2).

We next have for y with (2.1.3)

$$|\varphi(y)| \leq \frac{2^{204a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \quad (6.2.4)$$

and for $y' \neq y$ additionally with (2.1.4)

$$|\varphi(y) - \varphi(y')| \quad (6.2.5)$$

$$\leq |K_{s_1}(x_1, y) - K_{s_1}(x_1, y')| |K_{s_2}(x_2, y)| \quad (6.2.6)$$

$$+ |K_{s_1}(x_1, y')| |K_{s_2}(x_2, y) - K_{s_2}(x_2, y')| \quad (6.2.7)$$

$$\leq \frac{2^{229a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \left(\left(\frac{\rho(y, y')}{D^{s_1}} \right)^{1/a} + \left(\frac{\rho(y, y')}{D^{s_2}} \right)^{1/a} \right) \quad (6.2.8)$$

$$\leq \frac{2^{230a^3}}{\mu(B(x_1, D^{s_1}))\mu(B(x_2, D^{s_2}))} \left(\frac{\rho(y, y')}{D^{s_1}} \right)^{1/a}. \quad (6.2.9)$$

Adding the estimates (6.2.4) and (6.2.9) gives (6.2.3). This proves the lemma. \square

The following auxiliary statement about the support of $T_{\mathbf{p}}^*g$ will be used repeatedly.

Lemma 6.2.2 (tile range support). *For each $\mathbf{p} \in \mathfrak{P}$, and each $y \in X$, we have that*

$$T_{\mathbf{p}}^*g(y) \neq 0 \quad (6.2.10)$$

implies

$$y \in B(c(\mathbf{p}), 5D^{s(\mathbf{p})}). \quad (6.2.11)$$

Proof. Fix \mathbf{p} and y with (6.2.10). Then there exists $x \in E(\mathbf{p})$ with

$$\overline{K_{s(\mathbf{p})}(x, y)} e(-Q(x)(y) + Q(x)(x))g(x) \neq 0. \quad (6.2.12)$$

As $E(\mathbf{p}) \subset \mathcal{J}(\mathbf{p})$ and by the squeezing property (2.0.10), we have

$$\rho(x, c(\mathbf{p})) < 4D^{s(\mathbf{p})}. \quad (6.2.13)$$

As $K_{s(\mathbf{p})}(x, y) \neq 0$, we have by (2.1.2) that

$$\rho(x, y) \leq \frac{1}{2}D^{s(\mathbf{p})}. \quad (6.2.14)$$

Now (6.2.11) follows by the triangle inequality. \square

The next lemma is a geometric estimate for two tiles.

Lemma 6.2.3. *Let $\mathbf{p}_1, \mathbf{p}_2 \in \mathfrak{P}$ with $B(c(\mathbf{p}_1), 5D^{s(\mathbf{p}_1)}) \cap B(c(\mathbf{p}_2), 5D^{s(\mathbf{p}_2)}) \neq \emptyset$ and $s(\mathbf{p}_1) \leq s(\mathbf{p}_2)$. For each $x_1 \in E(\mathbf{p}_1)$ and $x_2 \in E(\mathbf{p}_2)$ we have*

$$1 + d_{\mathbf{p}_1}(Q(\mathbf{p}_1), Q(\mathbf{p}_2)) \leq 2^{8a}(1 + d_{B(x_1, D^{s(\mathbf{p}_1)})}(Q(x_1), Q(x_2))). \quad (6.2.15)$$

Proof. Let $i \in \{1, 2\}$. By Definition (2.0.20) of E , we have $Q(x_i) \in \Omega(\mathbf{p}_i)$. With (2.0.15) we then conclude

$$d_{\mathbf{p}_i}(Q(x_i), \mathcal{Q}(\mathbf{p}_i)) < 1. \quad (6.2.16)$$

We have by the triangle inequality and (2.0.10) that $\mathcal{J}(\mathbf{p}_1) \subset B(c(\mathbf{p}_2), 14D^{s(\mathbf{p}_2)})$. Thus, using again (2.0.10) and the doubling property (1.1.5)

$$d_{\mathbf{p}_1}(Q(x_2), \mathcal{Q}(\mathbf{p}_2)) \leq 2^{6a} d_{\mathbf{p}_2}(Q(x_2), \mathcal{Q}(\mathbf{p}_2)) \leq 2^{6a}. \quad (6.2.17)$$

By the triangle inequality, we obtain from (6.2.16) and (6.2.17)

$$1 + d_{\mathbf{p}_1}(Q(\mathbf{p}_1), \mathcal{Q}(\mathbf{p}_2)) \leq 2 + 2^{6a} + d_{\mathbf{p}_1}(Q(x_1), Q(x_2)). \quad (6.2.18)$$

As $x_1 \in \mathcal{J}(\mathbf{p}_1)$ by Definition (2.0.20) of E , we have by the squeezing property (2.0.10)

$$d(x_1, c(\mathbf{p}_1)) \leq 4D^{s(\mathbf{p}_1)} \quad (6.2.19)$$

and thus by (2.0.10) again and the triangle inequality

$$\mathcal{J}(\mathbf{p}_1) \subset B(x_1, 8D^{s(\mathbf{p}_1)}). \quad (6.2.20)$$

We thus estimate the right-hand side of (6.2.18) with monotonicity (1.1.6) of the metrics d_B by

$$\leq 2 + 2^{6a} + d_{B(x_1, 8D^{s(\mathbf{p}_1)})}(Q(x_1), Q(x_2)). \quad (6.2.21)$$

This is further estimated by applying the doubling property (1.1.5) three times by

$$\leq 2 + 2^{6a} + 2^{3a} d_{B_1(x_1, D^{s(\mathbf{p}_1)})}(Q(x_1), Q(x_2)). \quad (6.2.22)$$

Now (6.2.15) follows with $a \geq 4$. \square

We now prove Lemma 6.1.5.

Proof of Lemma 6.1.5. We begin with (6.1.42). By Lemma 6.2.2, the left-hand side of (6.1.42) vanishes if $B(c(\mathbf{p}'), 5D^{s(\mathbf{p}')} \cap B(c(\mathbf{p}), 5D^{s(\mathbf{p})}) = \emptyset$. Thus we can assume for the remainder of the proof that

$$B(c(\mathbf{p}'), 5D^{s(\mathbf{p}')} \cap B(c(\mathbf{p}), 5D^{s(\mathbf{p})}) \neq \emptyset. \quad (6.2.23)$$

We expand the left-hand side of (6.1.42) as

$$\left| \int \int_{E(\mathbf{p}')} \overline{K_{s(\mathbf{p}')} (x_1, y)} e(-Q(x_1)(y) + Q(x_1)(x_1)) g(x_1) d\mu(x_1) \right. \quad (6.2.24)$$

$$\left. \times \int_{E(\mathbf{p})} K_{s(\mathbf{p})} (x_2, y) e(Q(x_2)(y) - Q(x_2)(x_2)) \overline{g(x_2)} d\mu(x_2) d\mu(y) \right|. \quad (6.2.25)$$

By Fubini and the triangle inequality and the fact $|e(Q(x_i)(x_i))| = 1$ for $i = 1, 2$, we can estimate (6.2.24) and (6.2.25) from above by

$$\int_{E(\mathbf{p}')} \int_{E(\mathbf{p})} \mathbf{I}(x_1, x_2) d\mu(x_1) d\mu(x_2). \quad (6.2.26)$$

with

$$\mathbf{I}(x_1, x_2) := \left| \int e(-Q(x_1)(y) + Q(x_2)(y)) \varphi_{x_1, x_2}(y) d\mu(y) g(x_1) g(x_2) \right| \quad (6.2.27)$$

We estimate for fixed $x_1 \in E(\mathbf{p}')$ and $x_2 \in E(\mathbf{p})$ the inner integral of (6.2.26) with Proposition 2.0.5. The function $\varphi := \varphi_{x_1, x_2}$ satisfies the assumptions of Proposition 2.0.5 with $z = x_1$ and $R = D^{s_1}$ by Lemma 6.2.1. We obtain with $B' := B(x_1, D^{s(\mathbf{p}')})$,

$$\begin{aligned} \mathbf{I}(x_1, x_2) &\leq 2^{8a} \mu(B') \|\varphi\|_{C^\tau(B')} (1 + d_{B'}(Q(x_1), Q(x_2)))^{-1/(2a^2+a^3)} |g(x_1)g(x_2)| \\ &\leq \frac{2^{231a^3+8a}}{\mu(B(x_2, D^{s(\mathbf{p})}))} (1 + d_{B'}(Q(x_1), Q(x_2)))^{-1/(2a^2+a^3)}. \end{aligned} \quad (6.2.28)$$

Using (6.2.23), Lemma 6.2.3 and $a \geq 1$ estimates (6.2.28) by

$$\leq \frac{2^{231a^3+8a+1}}{\mu(B(x_2, D^{s(\mathbf{p})}))} (1 + d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{p})))^{-1/(2a^2+a^3)} |g(x_1)g(x_2)|. \quad (6.2.29)$$

As $x_2 \in \mathcal{J}(\mathbf{p})$ by Definition (2.0.20) of E , we have by (2.0.10)

$$\rho(x_2, c(\mathbf{p})) < 4D^{s(\mathbf{p})} \quad (6.2.30)$$

and thus by (2.0.10) again and the triangle inequality

$$\mathcal{J}(\mathbf{p}) \subset B(x_2, 8D^{s(\mathbf{p})}). \quad (6.2.31)$$

Using three iterations of the doubling property (1.1.2) give

$$\mu(\mathcal{J}(\mathbf{p})) \leq 2^{3a} \mu(B(x_2, D^{s(\mathbf{p})})). \quad (6.2.32)$$

With $a \geq 4$ and (6.2.29) we conclude (6.1.42).

Now assume the left-hand side of (6.1.42) is not zero. There is a $y \in X$ with

$$T_{\mathbf{p}'}^* g(y) \overline{T_{\mathbf{p}}^* g(y)} \neq 0 \quad (6.2.33)$$

By the triangle inequality and Lemma 6.2.2, we conclude

$$\rho(c(\mathbf{p}), c(\mathbf{p}')) \leq \rho(c(\mathbf{p}), y) + \rho(c(\mathbf{p}'), y) \leq 5D^{s(\mathbf{p})} + 5D^{s(\mathbf{p}')} \leq 10D^{s(\mathbf{p})}. \quad (6.2.34)$$

By the squeezing property (2.0.10) and the triangle inequality, we conclude

$$\mathcal{J}(\mathbf{p}') \subset B(c(\mathbf{p}), 14D^{s(\mathbf{p})}). \quad (6.2.35)$$

This completes the proof of Lemma 6.1.5. \square

6.3. Proof of the Antichain Tile Count Lemma.

Lemma 6.3.1 (tile reach). *Let $\vartheta \in \Theta$ and $N \geq 0$ be an integer. Let $\mathbf{p}, \mathbf{p}' \in \mathfrak{P}$ with*

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \vartheta) \leq 2^N \quad (6.3.1)$$

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \vartheta) \leq 2^N. \quad (6.3.2)$$

Assume $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{p}')$ and $s(\mathbf{p}) < s(\mathbf{p}')$. Then

$$2^{N+2}\mathbf{p} \lesssim 2^{N+2}\mathbf{p}'. \quad (6.3.3)$$

Proof. By Lemma 2.1.2, we have

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}'), \vartheta) \leq d_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), \vartheta) \leq 2^N. \quad (6.3.4)$$

Together with (6.3.1) and the triangle inequality, we obtain

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}'), \mathcal{Q}(\mathfrak{p})) \leq 2^{N+1}. \quad (6.3.5)$$

Now assume

$$\vartheta' \in B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), 2^{N+2}). \quad (6.3.6)$$

By the doubling property (1.1.5), applied five times, we have

$$d_{B(c(\mathfrak{p}'), 8D^{s(\mathfrak{p}')})}(\mathcal{Q}(\mathfrak{p}'), \vartheta') < 2^{5a+N+2}. \quad (6.3.7)$$

We have by the squeezing property (2.0.10)

$$c(\mathfrak{p}) \in B(c(\mathfrak{p}'), 4D^{s(\mathfrak{p}')}). \quad (6.3.8)$$

Hence by the triangle inequality

$$B(c(\mathfrak{p}), 4D^{s(\mathfrak{p}')}) \subseteq B(c(\mathfrak{p}'), 8D^{s(\mathfrak{p}')}). \quad (6.3.9)$$

Together with (6.3.7) and monotonicity (1.1.6) of d

$$d_{B(c(\mathfrak{p}), 4D^{s(\mathfrak{p}')})}(\mathcal{Q}(\mathfrak{p}'), \vartheta') < 2^{5a+N+2}. \quad (6.3.10)$$

Using the doubling property (1.1.7) $5a + 2$ times gives

$$d_{B(c(\mathfrak{p}), 2^{2-5a-2}D^{s(\mathfrak{p}')})}(\mathcal{Q}(\mathfrak{p}'), \vartheta') < 2^N. \quad (6.3.11)$$

Using $s(\mathfrak{p}) < s(\mathfrak{p}')$ and $D = 2^{100a^2}$ and $a \geq 4$ gives

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}'), \vartheta') < 2^N. \quad (6.3.12)$$

With the triangle inequality and (6.3.5),

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \vartheta') < 2^{N+2}. \quad (6.3.13)$$

This shows

$$B_{\mathfrak{p}'}(\mathcal{Q}(\mathfrak{p}'), 2^{N+2}) \subset B_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), 2^{N+2}). \quad (6.3.14)$$

This implies (6.3.3) and completes the proof of the lemma. \square

For $\vartheta \in \Theta$ and $N \geq 0$ define

$$\mathfrak{A}_{\vartheta, N} := \{\mathfrak{p} \in \mathfrak{A} : 2^N \leq 1 + d_{\mathfrak{p}}(\mathcal{Q}(\mathfrak{p}), \vartheta) < 2^{N+1}\}. \quad (6.3.15)$$

Lemma 6.3.2 (stack density). *Let $\vartheta \in \Theta$, $N \geq 0$ and $L \in \mathcal{D}$. Then*

$$\sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N} : \mathcal{J}(\mathfrak{p}) = L} \mu(E(\mathfrak{p}) \cap G) \leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \mu(L). \quad (6.3.16)$$

Proof. Let ϑ, N, L be given and set

$$\mathfrak{A}' := \{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N} : \mathcal{J}(\mathfrak{p}) = L\}. \quad (6.3.17)$$

Let $\mathfrak{p} \in \mathfrak{A}'$. We have by Definition (2.0.28) using $\lambda = 2$ and the squeezing property (2.0.15)

$$\mu(E(\mathfrak{p}) \cap G) \leq \mu(E_2(2, \mathfrak{p})) \leq 2^a \text{dens}_1(\mathfrak{A}') \mu(L). \quad (6.3.18)$$

By the covering property (1.1.8), applied $N + 4$ times, there is a collection Θ' of at most $2^{a(N+4)}$ elements such that

$$B_{\mathbf{p}}(\vartheta, 2^{N+1}) \subset \bigcup_{\vartheta' \in \Theta'} B_{\mathbf{p}}(\vartheta', 0.2). \quad (6.3.19)$$

As each $\mathcal{Q}(\mathbf{p}')$ with $\mathbf{p}' \in \mathfrak{A}'$ is contained in the left-hand-side of (6.3.19) by definition (because $\mathcal{J}(\mathbf{p}') = \mathcal{J}(\mathbf{p})$), it is in at least one $B_{\mathbf{p}}(\vartheta', 0.2)$ with $\vartheta' \in \Theta'$.

For two different $\mathbf{p}', \mathbf{p}'' \in \mathfrak{A}'$, we have by (2.0.13) that $\Omega(\mathbf{p}')$ and $\Omega(\mathbf{p}'')$ are disjoint and thus by the squeezing property (2.0.15) we have for every $\vartheta' \in \Theta'$

$$\vartheta' \notin B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}'), 0.2) \cap B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}''), 0.2). \quad (6.3.20)$$

Hence at most one of $\mathcal{Q}(\mathbf{p}')$ and $\mathcal{Q}(\mathbf{p}'')$ is in $B_{\mathbf{p}}(\vartheta', 0.2)$. It follows that there are at most $2^{a(N+4)}$ elements in \mathfrak{A}' . Adding (6.3.18) over \mathfrak{A}' proves (6.3.16). \square

Lemma 6.3.3 (local antichain density). *Let $\vartheta \in \Theta$ and N be an integer. Let \mathbf{p}_{ϑ} be a tile with $\vartheta \in B_{\mathbf{p}_{\vartheta}}(\mathcal{Q}(\mathbf{p}_{\vartheta}), 2^{N+1})$. Then we have*

$$\sum_{\mathbf{p} \in \mathfrak{A}_{\vartheta, N}: s(\mathbf{p}_{\vartheta}) < s(\mathbf{p})} \mu(E(\mathbf{p}) \cap G \cap \mathcal{J}(\mathbf{p}_{\vartheta})) \leq \mu(E_2(2^{N+3}, \mathbf{p}_{\vartheta})). \quad (6.3.21)$$

Proof. Let \mathbf{p} be any tile in $\mathfrak{A}_{\vartheta, N}$ with $s(\mathbf{p}_{\vartheta}) < s(\mathbf{p})$. By definition of E , the tile contributes zero to the sum on the left-hand side of (6.3.21) unless $\mathcal{J}(\mathbf{p}) \cap \mathcal{J}(\mathbf{p}_{\vartheta}) \neq \emptyset$, which we may assume. With $s(\mathbf{p}_{\vartheta}) < s(\mathbf{p})$ and the dyadic property (2.0.8) we conclude $\mathcal{J}(\mathbf{p}_{\vartheta}) \subset \mathcal{J}(\mathbf{p})$. We conclude from $\mathbf{p} \in \mathfrak{A}_{\vartheta, N}$ that

$$\vartheta \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 2^{N+1}). \quad (6.3.22)$$

With Lemma 6.3.1 and the assumption on \mathbf{p}_{ϑ} , we conclude

$$2^{N+3}\mathbf{p}_{\vartheta} \lesssim 2^{N+3}\mathbf{p}. \quad (6.3.23)$$

By Definition (2.0.27) of E_2 , we conclude

$$E(\mathbf{p}) \cap G \cap \mathcal{J}(\mathbf{p}_{\vartheta}) \subset E_2(2^{N+3}, \mathbf{p}_{\vartheta}). \quad (6.3.24)$$

Using disjointedness of the various $E(\mathbf{p})$ with $\mathbf{p} \in \mathfrak{A}$ by Lemma 6.1.1, we obtain (6.3.21). This proves the lemma. \square

Lemma 6.3.4 (global antichain density). *Let $\vartheta \in Q(X)$ and let $N \geq 0$ be an integer. Then we have*

$$\sum_{\mathbf{p} \in \mathfrak{A}_{\vartheta, N}} \mu(E(\mathbf{p}) \cap G) \leq 2^{101a^3 + Na} \text{dens}_1(\mathfrak{A}) \mu(\cup_{\mathbf{p} \in \mathfrak{A}} I_{\mathbf{p}}). \quad (6.3.25)$$

Proof. Fix ϑ and N . Let \mathfrak{A}' be the set of $\mathbf{p} \in \mathfrak{A}_{\vartheta, N}$ such that $\mathcal{J}(\mathbf{p}) \cap G$ is not empty and $s(\mathbf{p}) > -S$. Let \mathfrak{A}_{-S} be the set of $\mathbf{p} \in \mathfrak{A}_{\vartheta, N}$ such that $\mathcal{J}(\mathbf{p}) \cap G$ is not empty and $s(\mathbf{p}) = -S$. Then we have

$$\sum_{\mathbf{p} \in \mathfrak{A}_{\vartheta, N}} \mu(E(\mathbf{p}) \cap G) = \sum_{\mathbf{p} \in \mathfrak{A}'} \mu(E(\mathbf{p}) \cap G) + \sum_{\mathbf{p} \in \mathfrak{A}_{-S}} \mu(E(\mathbf{p}) \cap G)$$

We start by estimating the contribution of \mathfrak{A}_{-S} . Let \mathcal{L}_{-S} be the collection of dyadic cubes $\mathcal{J}(\mathbf{p})$ with $\mathbf{p} \in \mathfrak{A}_{-S}$. They all have scale $-S$, by definition of \mathcal{L}_{-S} , and hence they are pairwise disjoint by the dyadic property (2.0.8). We write

$$\sum_{\mathbf{p} \in \mathfrak{A}_{-S}} \mu(E(\mathbf{p}) \cap G) = \sum_{L \in \mathcal{L}_{-S}} \sum_{\mathbf{p} \in \mathfrak{A}_{-S}, \mathcal{J}(\mathbf{p})=L} \mu(E(\mathbf{p}) \cap G),$$

and using Lemma 6.3.2, we estimate

$$\leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \sum_{L \in \mathcal{L}_{-S}} \mu(L) \leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \mu(\cup_{\mathbf{p} \in \mathfrak{A}} I_{\mathbf{p}}). \quad (6.3.26)$$

We turn to \mathfrak{A}' . Let \mathcal{L} be the collection of dyadic cubes $I \in \mathcal{D}$ such that $I \leq \mathcal{J}(\mathbf{p})$ for some $\mathbf{p} \in \mathfrak{A}'$ and such that $\mathcal{J}(\mathbf{p}) \not\subset I$ for all $\mathbf{p} \in \mathfrak{A}'$. By (2.0.7), for each $\mathbf{p} \in \mathfrak{A}'$ and each $x \in \mathcal{J}(\mathbf{p}) \cap G$, there is $I \in \mathcal{D}$ with $s(I) = -S$ and $x \in I$. By (2.0.8), we have $I \subset \mathcal{J}(\mathbf{p})$. Since for all $\mathbf{p}' \in \mathfrak{A}'$ we have $s(\mathbf{p}') > -S$, we have $I \in \mathcal{L}$. Hence

$$\mathcal{J}(\mathbf{p}) \subset \bigcup \{I \in \mathcal{D} : s(I) = -S, I \subset \mathcal{J}(\mathbf{p})\} \subset \bigcup \mathcal{L}. \quad (6.3.27)$$

As each $I \in \mathcal{L}$ satisfies $I \subset \mathcal{J}(\mathbf{p})$ for some \mathbf{p} in \mathfrak{A}' , we conclude

$$\bigcup \mathcal{L} = \bigcup_{\mathbf{p} \in \mathfrak{A}'} \mathcal{J}(\mathbf{p}). \quad (6.3.28)$$

Let \mathcal{L}^* be the set of maximal elements in \mathcal{L} with respect to set inclusion. By (2.0.8), the elements in \mathcal{L}^* are pairwise disjoint and we have

$$\bigcup \mathcal{L}^* = \bigcup_{\mathbf{p} \in \mathfrak{A}'} \mathcal{J}(\mathbf{p}). \quad (6.3.29)$$

Using the partition (6.3.29) into elements of \mathcal{L} in (6.3.30), it suffices to show for each $L \in \mathcal{L}^*$

$$\sum_{\mathbf{p} \in \mathfrak{A}'} \mu(E(\mathbf{p}) \cap G \cap L) \leq 2^{101a^3+aN} \text{dens}_1(\mathfrak{A}) \mu(L). \quad (6.3.30)$$

Fix $L \in \mathcal{L}^*$. By definition of \mathcal{L}^* , there exists an element $\mathbf{p}' \in \mathfrak{A}'$ such that $L \subset \mathcal{J}(\mathbf{p}')$. Pick such an element \mathbf{p}' in \mathfrak{A} with minimal $s(\mathbf{p}')$. As $\mathcal{J}(\mathbf{p}') \not\subset L$ by definition of L , we have with (2.0.8) that $s(L) < s(\mathbf{p}')$. In particular $s(L) < S$, thus $L \neq I_0$ and hence by (2.0.9) there exists a cube $J \in \mathcal{D}$ with $L \subsetneq J$. By (2.0.7), there is an $L' \in \mathcal{D}$ with $s(L') = s(L) + 1$ and $L \leq L'$. By (2.0.8), we have $L \subset L'$.

We split the left-hand side of (6.3.30) as

$$\sum_{\mathbf{p} \in \mathfrak{A}': \mathcal{J}(\mathbf{p})=L'} \mu(E(\mathbf{p}) \cap G \cap L) \quad (6.3.31)$$

$$+ \sum_{\mathbf{p} \in \mathfrak{A}': \mathcal{J}(\mathbf{p}) \neq L'} \mu(E(\mathbf{p}) \cap G \cap L), \quad (6.3.32)$$

We first estimate (6.3.31) with Lemma 6.3.2 by

$$\leq \sum_{\mathbf{p} \in \mathfrak{A}': \mathcal{J}(\mathbf{p})=L'} \mu(E(\mathbf{p}) \cap G \cap L') \leq 2^{a(N+5)} \text{dens}_1(\mathfrak{A}) \mu(L'). \quad (6.3.33)$$

We turn to (6.3.32). Consider the element $\mathbf{p}' \in \mathfrak{A}'$ as above with $L \subset \mathcal{J}(\mathbf{p}')$ and $s(L) < s(\mathbf{p}')$. As $L \subset L'$ and $s(L') = s(L) + 1$, we conclude with the dyadic property that $L' \subset \mathcal{J}(\mathbf{p}')$. By maximality of L , we have $L' \notin \mathcal{L}$. This together with the

existence of the given $\mathbf{p}' \in \mathfrak{A}$ with $L' \subset \mathcal{J}(\mathbf{p}')$ shows by definition of \mathcal{L} that there exists $\mathbf{p}'' \in \mathfrak{A}'$ with $\mathcal{J}(\mathbf{p}'') \subset L'$.

If $\mathcal{J}(\mathbf{p}'') = L'$, then we set $\mathbf{p}_\vartheta = \mathbf{p}''$ and note that as $\mathbf{p}'' \in \mathfrak{A}_{\vartheta, N}$

$$\vartheta \in B(Q(\mathbf{p}''), 2^{N+1}). \quad (6.3.34)$$

If $\mathcal{J}(\mathbf{p}'') \neq L'$, then it follows that $s(\mathbf{p}'') < s(L')$. By the covering property (2.0.13), there exists a unique \mathbf{p}_ϑ with

$$\mathcal{J}(\mathbf{p}_\vartheta) = L'$$

such that $\vartheta \in \Omega(\mathbf{p}_\vartheta)$. We take this as the definition of \mathbf{p}_ϑ in this case. Note that

$$\vartheta \in B(Q(\mathbf{p}_\vartheta), 1) \quad (6.3.35)$$

so by Lemma 6.3.1, we conclude

$$2^{N+3}\mathbf{p}'' \lesssim 2^{N+3}\mathbf{p}_\vartheta. \quad (6.3.36)$$

This clearly also holds in the case $\mathcal{J}(\mathbf{p}'') = L'$, since then $\mathbf{p}'' = \mathbf{p}_\vartheta$. Furthermore, in both cases it also holds that

$$\vartheta \in B_{\mathbf{p}_\vartheta}(Q(\mathbf{p}_\vartheta), 2^{N+1}). \quad (6.3.37)$$

As $\mathbf{p}'' \in \mathfrak{A}'$, we have by Definition (2.0.28) of dens_1 that

$$\mu(E_2(2^{N+3}, \mathbf{p}_\vartheta)) \leq 2^{Na+3a} \text{dens}_1(\mathfrak{A})\mu(L'). \quad (6.3.38)$$

Now let \mathbf{p} be any tile in the summation set in (6.3.32), that is, $\mathbf{p} \in \mathfrak{A}'$ and $\mathcal{J}(\mathbf{p}) \neq L'$. Then $\mathcal{J}(\mathbf{p}) \cap L \neq \emptyset$. It follows by the dyadic property (2.0.8) and the definition of L that $L \subset \mathcal{J}(\mathbf{p})$ and $L \neq \mathcal{J}(\mathbf{p})$. By the dyadic property (2.0.8), we have $s(L) < s(\mathbf{p})$ and thus $s(L') \leq s(\mathbf{p})$. By the dyadic property (2.0.8) again, we have $L' \subset \mathcal{J}(\mathbf{p})$. As $L' \neq \mathcal{J}(\mathbf{p})$, we conclude $s(L) < s(\mathbf{p})$. By Lemma 6.3.3, we can thus estimate (6.3.32) by

$$\sum_{\mathbf{p} \in \mathfrak{A}': \mathcal{J}(\mathbf{p}) \neq L'} \mu(E(\mathbf{p}) \cap G \cap L') \leq \mu(E_2(2^{N+3}, \mathbf{p}_\vartheta)). \quad (6.3.39)$$

With the decomposition in (6.3.31) and (6.3.32) and the estimates (6.3.33), (6.3.21), (6.3.38) we obtain the estimate

$$\sum_{\mathbf{p} \in \mathfrak{A}'} \mu(E(\mathbf{p}) \cap G \cap L) \leq (2^{a(N+5)} + 2^{Na+3a}) \text{dens}_1(\mathfrak{A})\mu(L'). \quad (6.3.40)$$

Using $s(L') = s(L) + 1$ and $D = 2^{100a^2}$ and the squeezing property (2.0.10) and the doubling property (1.1.2) $100a^2 + 4$ times, we obtain

$$\mu(L') \leq 2^{100a^3+4a} \mu(L). \quad (6.3.41)$$

Inserting in (6.3.40), adding the estimate (6.3.26) and using $a \geq 4$ gives (6.3.30). This completes the proof of the lemma. \square

We turn to the proof of Lemma 6.1.6.

Proof of Lemma 6.1.6. Using that \mathfrak{A} is the disjoint union of the $\mathfrak{A}_{\vartheta, N}$ with $N \geq 0$, we estimate the left-hand side (6.1.45) with the triangle inequality by

$$\leq \sum_{N \geq 0} \left\| \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})} \mathbf{1}_G \right\|_p \quad (6.3.42)$$

We consider each individual term in this sum and estimate its p -th power. Using that for each $x \in X$ by Lemma 6.1.1 there is at most one $\mathfrak{p} \in \mathfrak{A}$ with $x \in E(\mathfrak{p})$, we have

$$\left\| \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p})} \mathbf{1}_G \right\|_p^p \quad (6.3.43)$$

$$= \int \left(\sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-N/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p}) \cap G}(x) \right)^p d\mu(x) \quad (6.3.44)$$

$$= \int \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} 2^{-pN/(2a^2+a^3)} \mathbf{1}_{E(\mathfrak{p}) \cap G}(x) d\mu(x) \quad (6.3.45)$$

$$= 2^{-pN/(2a^2+a^3)} \sum_{\mathfrak{p} \in \mathfrak{A}_{\vartheta, N}} \mu(E(\mathfrak{p}) \cap G) \quad (6.3.46)$$

Using Lemma 6.3.4, we estimate the last display by

$$\leq 2^{-pN/(2a^2+a^3)+101a^3+Na} \text{dens}_1(\mathfrak{A}) \mu \left(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{J}(\mathfrak{p}) \right) \quad (6.3.47)$$

Using that $a \geq 4$ and since $p = 4a^4$, we have

$$pN/(2a^2 + a^3) \geq 4a^4 N/(3a^3) \geq Na + N.$$

Hence we have for (6.3.47) the upper bound

$$\leq 2^{101a^3-N} \text{dens}_1(\mathfrak{A}) \mu \left(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{J}(\mathfrak{p}) \right).$$

Taking the p -th root and summing over $N \geq 0$ gives for (6.3.42) the upper bound

$$\begin{aligned} &\leq \left(\sum_{N \geq 0} 2^{-N/p} \right) 2^{101a^3/p} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu \left(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{J}(\mathfrak{p}) \right)^{\frac{1}{p}} \\ &= (1 - 2^{-1/p})^{-1} 2^{101a^3/p} \text{dens}_1(\mathfrak{A})^{\frac{1}{p}} \mu \left(\cup_{\mathfrak{p} \in \mathfrak{A}} \mathcal{J}(\mathfrak{p}) \right)^{\frac{1}{p}}. \end{aligned}$$

Using that $p = 4a^4$ and $a \geq 4$, this proves the lemma. \square

7. PROOF OF THE FOREST OPERATOR PROPOSITION

7.1. The pointwise tree estimate. Fix a forest $(\mathfrak{U}, \mathfrak{T})$. The main result of this subsection is Lemma 7.1.3, we begin this section with some definitions necessary to state the lemma.

For $\mathfrak{u} \in \mathfrak{U}$ and $x \in X$, we define

$$\sigma(\mathfrak{u}, x) := \{s(\mathfrak{p}) : \mathfrak{p} \in \mathfrak{T}(\mathfrak{u}), x \in E(\mathfrak{p})\}.$$

This is a subset of $\mathbb{Z} \cap [-S, S]$, so has a minimum and a maximum. We set

$$\bar{\sigma}(\mathfrak{u}, x) := \max \sigma(\mathfrak{T}(\mathfrak{u}), x)$$

$$\underline{\sigma}(\mathbf{u}, x) := \min \sigma(\mathfrak{T}(\mathbf{u}), x).$$

Lemma 7.1.1 (convex scales). *For each $\mathbf{u} \in \mathfrak{U}$, we have*

$$\sigma(\mathbf{u}, x) = \mathbb{Z} \cap [\underline{\sigma}(\mathbf{u}, x), \bar{\sigma}(\mathbf{u}, x)].$$

Proof. Let $s \in \mathbb{Z}$ with $\underline{\sigma}(\mathbf{u}, x) \leq s \leq \bar{\sigma}(\mathbf{u}, x)$. By definition of σ , there exists $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}) = \underline{\sigma}(\mathbf{u}, x)$ and $x \in E(\mathbf{p})$, and there exists $\mathbf{p}'' \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}'') = \bar{\sigma}(\mathbf{u}, x)$ and $x \in E(\mathbf{p}'') \subset J(\mathbf{p}'')$. By property (2.0.7) of the dyadic grid, there exists a cube $I \in \mathcal{D}$ of scale s with $x \in I$. By property (2.0.13), there exists a tile $\mathbf{p}' \in \mathfrak{P}(I)$ with $Q(x) \in \Omega(\mathbf{p}')$. By the dyadic property (2.0.8) we have $J(\mathbf{p}) \subset J(\mathbf{p}') \subset J(\mathbf{p}'')$, and by (2.0.14), we have $\Omega(\mathbf{p}'') \subset \Omega(\mathbf{p}') \subset \Omega(\mathbf{p})$. Thus $\mathbf{p} \leq \mathbf{p}' \leq \mathbf{p}''$, which gives with the convexity property (2.0.33) of $\mathfrak{T}(\mathbf{u})$ that $\mathbf{p}' \in \mathfrak{T}(\mathbf{u})$, so $s \in \sigma(\mathbf{u}, x)$. \square

For a nonempty collection of tiles $\mathfrak{S} \subset \mathfrak{P}$ we define

$$\mathcal{J}_0(\mathfrak{S})$$

to be the collection of all dyadic cubes $J \in \mathcal{D}$ such that $s(J) = -S$ or

$$J(\mathbf{p}) \not\subset B(c(J), 100D^{s(J)+1})$$

for all $\mathbf{p} \in \mathfrak{S}$. We define $\mathcal{J}(\mathfrak{S})$ to be the collection of inclusion maximal cubes in $\mathcal{J}_0(\mathfrak{S})$.

We further define

$$\mathcal{L}_0(\mathfrak{S})$$

to be the collection of dyadic cubes $L \in \mathcal{D}$ such that $s(L) = -S$, or there exists $\mathbf{p} \in \mathfrak{S}$ with $L \subset J(\mathbf{p})$ and there exists no $\mathbf{p} \in \mathfrak{S}$ with $J(\mathbf{p}) \subset L$. We define $\mathcal{L}(\mathfrak{S})$ to be the collection of inclusion maximal cubes in $\mathcal{L}_0(\mathfrak{S})$.

Lemma 7.1.2 (dyadic partitions). *For each $\mathfrak{S} \subset \mathfrak{P}$, we have*

$$\bigcup_{I \in \mathcal{D}} I = \bigcup_{J \in \mathcal{J}(\mathfrak{S})} J \tag{7.1.1}$$

and

$$\bigcup_{I \in \mathcal{D}} I = \bigcup_{L \in \mathcal{L}(\mathfrak{S})} L. \tag{7.1.2}$$

Proof. Since $\mathcal{J}(\mathfrak{S})$ is the set of inclusion maximal cubes in $\mathcal{J}_0(\mathfrak{S})$, cubes in $\mathcal{J}(\mathfrak{S})$ are pairwise disjoint by (2.0.8). The same applies to $\mathcal{L}(\mathfrak{S})$.

If $x \in \bigcup_{I \in \mathcal{D}} I$, then there exists by (2.0.7) a cube $I \in \mathcal{D}$ with $x \in I$ and $s(I) = -S$. Then $I \in \mathcal{J}_0(\mathfrak{S})$. There exists an inclusion maximal cube in $\mathcal{J}_0(\mathfrak{S})$ containing I . This cube contains x and is contained in $\mathcal{J}(\mathfrak{S})$. This shows one inclusion in (7.1.1), the other one follows from $\mathcal{J}(\mathfrak{S}) \subset \mathcal{D}$.

The proof of the two inclusions in (7.1.2) is similar. \square

For a finite collection of pairwise disjoint cubes \mathcal{C} , define the projection operator

$$P_{\mathcal{C}}f(x) := \sum_{J \in \mathcal{C}} \mathbf{1}_J(x) \frac{1}{\mu(J)} \int_J f(y) d\mu(y).$$

Given a scale $-S \leq s \leq S$ and a point $x \in \bigcup_{I \in \mathcal{D}, s(I)=s} I$, there exists a unique cube in \mathcal{D} of scale s containing x by (2.0.7). We denote it by $I_s(x)$. Define for $\vartheta \in \Theta$ the nontangential maximal operator

$$T_{\mathcal{N}}^{\vartheta} f(x) := \sup_{-S \leq s_1 < S} \sup_{x' \in I_{s_1}(x)} \sup_{\substack{s_1 \leq s_2 \leq S \\ D^{s_2-1} \leq R_Q(\vartheta, x')}} \left| \sum_{s=s_1}^{s_2} \int K_s(x', y) f(y) d\mu(y) \right|. \quad (7.1.3)$$

Define for each $\mathbf{u} \in \mathfrak{U}$ the auxiliary operator

$$S_{1, \mathbf{u}} f(x) := \sum_{I \in \mathcal{D}} \mathbf{1}_I(x) \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(c(I), 16D^{s(I)}) \\ s(J) \leq s(I)}} \frac{D^{(s(J)-s(I))/a}}{\mu(B(c(I), 16D^{s(I)}))} \int_J |f(y)| d\mu(y). \quad (7.1.4)$$

Define also the collection of balls

$$\mathcal{B} = \{B(c(I), 2^s D^{s(I)+t}) : I \in \mathcal{D}, 0 \leq s \leq S+5, 0 \leq t \leq 2S+3\}.$$

The following pointwise estimate for operators associated to sets $\mathfrak{T}(\mathbf{u})$ is the main result of this subsection.

Lemma 7.1.3 (pointwise tree estimate). *Let $\mathbf{u} \in \mathfrak{U}$ and $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$. Let $x, x' \in L$. Then for all bounded functions f with bounded support*

$$\left| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}[e(-\mathcal{Q}(\mathbf{u}))f](x) \right| \leq 2^{129a^3} (M_{\mathcal{B}, 1} + S_{1, \mathbf{u}}) P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x') + |T_{\mathcal{N}}^{\mathcal{Q}(\mathbf{u})} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(x')|, \quad (7.1.5)$$

Proof. By (2.0.21), if $T_{\mathbf{p}}[e(-\mathcal{Q}(\mathbf{u}))f](x) \neq 0$, then $x \in E(\mathbf{p})$. Combining this with $|e(\mathcal{Q}(\mathbf{u})(x))| = 1$, we obtain

$$\begin{aligned} & \left| \sum_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathbf{p}}[e(-\mathcal{Q}(\mathbf{u}))f](x) \right| \\ &= \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int e(-\mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(x)(y) + \mathcal{Q}(\mathbf{u})(x) - \mathcal{Q}(x)(x)) \times \right. \\ & \qquad \qquad \qquad \left. K_s(x, y) f(y) d\mu(y) \right|. \end{aligned}$$

Using the triangle inequality, we bound this by the sum of three terms:

$$\leq \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int (e(-\mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(x)(y) + \mathcal{Q}(\mathbf{u})(x) - \mathcal{Q}(x)(x)) - 1) \times \right. \\ \left. K_s(x, y) f(y) d\mu(y) \right| \quad (7.1.6)$$

$$+ \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int K_s(x, y) P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y) \, d\mu(y) \right| \quad (7.1.7)$$

$$+ \left| \sum_{s \in \sigma(\mathbf{u}, x)} \int K_s(x, y) (f(y) - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y)) \, d\mu(y) \right|. \quad (7.1.8)$$

The proof is completed using the bounds for these three terms proven respectively in Lemma 7.1.4, Lemma 7.1.5 and Lemma 7.1.6. \square

Lemma 7.1.4 (first tree pointwise). *For all $\mathbf{u} \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$(7.1.6) \leq 10 \cdot 2^{104a^3} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x').$$

Proof. Let $s \in \sigma(\mathbf{u}, x)$. If $x, y \in X$ are such that $K_s(x, y) \neq 0$, then, by (2.1.2), we have $\rho(x, y) \leq 1/2D^s$. By 1-Lipschitz continuity of the function $t \mapsto \exp(it) = e(t)$ and the property (1.1.4) of the metrics d_B , it follows that

$$\begin{aligned} & |e(-Q(\mathbf{u})(y) + Q(x)(y) + Q(\mathbf{u})(x) - Q(x)(x)) - 1| \\ & \leq d_{B(x, 1/2D^s)}(Q(\mathbf{u}), Q(x)). \end{aligned}$$

Let $\mathbf{p}_s \in \mathfrak{T}(\mathbf{u})$ be a tile with $s(\mathbf{p}_s) = s$ and $x \in E(\mathbf{p}_s)$, and let \mathbf{p}' be a tile with $s(\mathbf{p}') = \bar{\sigma}(\mathbf{u}, x)$ and $x \in E(\mathbf{p}')$. Using the monotonicity property (1.1.6), the doubling property (1.1.5) repeatedly, the definition of $d_{\mathbf{p}}$ and Lemma 2.1.2, we can bound the previous display by

$$d_{B(x, 4D^s)}(Q(\mathbf{u}), Q(x)) \leq 2^{4a} d_{\mathbf{p}_s}(Q(\mathbf{u}), Q(x)) \leq 2^{4a} 2^{s - \bar{\sigma}(\mathbf{u}, x)} d_{\mathbf{p}'}(Q(\mathbf{u}), Q(x)).$$

Since $Q(\mathbf{u}) \in B_{\mathbf{p}'}(Q(\mathbf{p}'), 4)$ by (2.0.32) and $Q(x) \in \Omega(\mathbf{p}') \subset B_{\mathbf{p}'}(Q(\mathbf{p}'), 1)$ by (2.0.15), this is estimated by

$$\leq 5 \cdot 2^{4a} 2^{s - \bar{\sigma}(\mathbf{u}, x)}.$$

Using (2.1.3), it follows that

$$(7.1.6) \leq 5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s - \bar{\sigma}(\mathbf{u}, x)} \frac{1}{\mu(B(x, D^s))} \int_{B(x, 0.5D^s)} |f(y)| \, d\mu(y).$$

By (7.1.1), the collection \mathcal{J} is a partition of $\bigcup_{I \in \mathcal{D}} I$, so this is estimated by

$$5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s - \bar{\sigma}(\mathbf{u}, x)} \frac{1}{\mu(B(x, D^s))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \cap B(x, 0.5D^s) \neq \emptyset}} \int_J |f(y)| \, d\mu(y).$$

This expression does not change if we replace $|f|$ by $P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|$.

Let $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$ with $B(x, 0.5D^s) \cap J \neq \emptyset$. By the triangle inequality and since $x \in E(\mathbf{p}_s) \subset B(c(\mathbf{p}_s), 4D^s)$, it follows that $B(c(\mathbf{p}_s), 4.5D^s) \cap J \neq \emptyset$. If $s(J) \geq s$ and $s(J) > -S$, then it follows from the triangle inequality, (2.0.10) and (2.0.1) that $J(\mathbf{p}_s) \subset B(c(J), 100D^{s(J)+1})$, contradicting $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$. Thus $s(J) \leq s - 1$ or $s(J) = -S$. If $s(J) = -S$ and $s(J) > s - 1$, then $s = -S$. Thus we always have $s(J) \leq s$. It then follows from the triangle inequality and (2.0.10) that $J \subset B(c(\mathbf{p}_s), 16D^s)$.

Thus we can continue our chain of estimates with

$$5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u,x)} \frac{1}{\mu(B(x, D^s))} \int_{B(c(\mathbf{p}_s), 16D^s)} P_{\mathcal{J}(\mathfrak{T}(u))} |f(y)| d\mu(y).$$

We have $B(c(\mathbf{p}_s), 16D^s) \subset B(x, 32D^s)$, by (2.0.10) and the triangle inequality, since $x \in \mathcal{J}(\mathbf{p}_s)$. Combining this with the doubling property (1.1.2), we obtain

$$\mu(B(c(\mathbf{p}_s), 16D^s)) \leq 2^{5a} \mu(B(x, D^s)).$$

Since $a \geq 4$, it follows that (7.1.6) is bounded by

$$5 \cdot 2^{103a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u,x)} \frac{1}{\mu(B(c(\mathbf{p}_s), 16D^s))} \int_{B(c(\mathbf{p}_s), 16D^s)} P_{\mathcal{J}(\mathfrak{T}(u))} |f(y)| d\mu(y).$$

Since $L \in \mathcal{L}(\mathfrak{T}(u))$ and $x \in L \cap \mathcal{J}(\mathbf{p}_s)$, we have $s(L) \leq s(\mathbf{p}_s)$. It follows by (2.0.8) that $L \subset \mathcal{J}(\mathbf{p}_s)$, in particular $x' \in \mathcal{J}(\mathbf{p}_s) \subset B(c(\mathbf{p}_s), 16D^s)$. Thus

$$\begin{aligned} &\leq 5 \cdot 2^{104a^3} \sum_{s \in \sigma(x)} 2^{s-\bar{\sigma}(u,x)} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(u))} |f|(x') \\ &\leq 10 \cdot 2^{104a^3} M_{\mathcal{B},1} P_{\mathcal{J}(\mathfrak{T}(u))} |f|(x'). \end{aligned}$$

This completes the estimate for term (7.1.6). \square

Lemma 7.1.5 (second tree pointwise). *For all $u \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(u))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$\left| \sum_{s \in \sigma(u,x)} \int K_s(x, y) P_{\mathcal{J}(\mathfrak{T}(u))} f(y) d\mu(y) \right| \leq T_{\mathcal{N}}^{Q(u)} P_{\mathcal{J}(\mathfrak{T}(u))} f(x').$$

Proof. Let $s_1 = \underline{\sigma}(u, x)$. By definition, there exists a tile $\mathbf{p} \in \mathfrak{T}(u)$ with $s(\mathbf{p}) = s_1$ and $x \in E(\mathbf{p})$. Then $x \in \mathcal{J}(\mathbf{p}) \cap L$. By (2.0.8) and the definition of $\mathcal{L}(\mathfrak{T}(u))$, it follows that $L \subset \mathcal{J}(\mathbf{p})$, in particular $x' \in \mathcal{J}(\mathbf{p})$, so $x \in I_{s_1}(x')$. Next, let $s_2 = \bar{\sigma}(u, x)$ and let $\mathbf{p}' \in \mathfrak{T}(u)$ with $s(\mathbf{p}') = s_2$ and $x \in E(\mathbf{p}')$. Since $\mathbf{p}' \in \mathfrak{T}(u)$, we have $4\mathbf{p}' \lesssim u$. Since $Q(x) \in \Omega(\mathbf{p}')$, it follows that

$$d_{\mathbf{p}}(Q(u), Q(x)) \leq 5.$$

Applying the doubling property (1.1.5) five times, we obtain

$$d_{B(c(\mathbf{p}), 8D^{s_2})}(Q(u), Q(x)) \leq 5 \cdot 2^{5a}.$$

By the triangle inequality, we have $B(x, D^{s_2}) \subset B(c(\mathbf{p}), 8D^{s_2})$, so by (1.1.6)

$$d_{B(x, D^{s_2})}(Q(u), Q(x)) \leq 5 \cdot 2^{5a}.$$

Finally, by applying (1.1.7) $100a$ times, we obtain

$$d_{B(x, D^{s_2-1})}(Q(u), Q(x)) \leq 5 \cdot 2^{-95a} < 1.$$

Consequently, $D^{s_2-1} < R_Q(Q(u), x)$. The lemma now follows from the definition of $T_{\mathcal{N}}$. \square

Lemma 7.1.6 (third tree pointwise). *For all $\mathbf{u} \in \mathfrak{U}$, all $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$, all $x, x' \in L$ and all bounded f with bounded support, we have*

$$\left| \sum_{s \in \sigma(\mathbf{u}, x)} \int K_s(x, y) (f(y) - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} f(y)) d\mu(y) \right| \leq 2^{128a^3} S_{1, \mathbf{u}} P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|(x').$$

Proof. We have for $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$:

$$\begin{aligned} & \int_J K_s(x, y) (1 - P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))}) f(y) d\mu(y) \\ &= \int_J \frac{1}{\mu(J)} \int_J K_s(x, y) - K_s(x, z) d\mu(z) f(y) d\mu(y). \end{aligned} \quad (7.1.9)$$

By (2.1.4) and (2.0.10), we have for $y, z \in J$

$$|K_s(x, y) - K_s(x, z)| \leq \frac{2^{127a^3}}{\mu(B(x, D^s))} \left(\frac{8D^{s(J)}}{D^s} \right)^{1/a}.$$

Suppose that $s \in \sigma(\mathbf{u}, x)$. If $K_s(x, y) \neq 0$ for some $y \in J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$ then, by (2.1.2), $y \in B(x, 0.5D^s) \cap J \neq \emptyset$. Let $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $s(\mathbf{p}) = s$ and $x \in E(\mathbf{p})$. Then $B(c(\mathbf{p}_s), 4.5D^s) \cap J \neq \emptyset$ by the triangle inequality. If $s(J) \geq s$ and $s(J) > -S$, then it follows from the triangle inequality, (2.0.10) and (2.0.1) that $\mathcal{J}(\mathbf{p}) \subset B(c(J), 100D^{s(J)+1})$, contradicting $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$. Thus $s(J) \leq s - 1$ or $s(J) = -S$. If $s(J) = -S$ and $s(J) > s - 1$, then $s = -S$. So in both cases, $s(J) \leq s$. It then follows from the triangle inequality and (2.0.10) that $J \subset B(x, 16D^s)$.

Thus, we can estimate (7.1.8) by

$$\begin{aligned} & 2^{127a^3+3/a} \sum_{\mathbf{p} \in \mathfrak{T}} \frac{\mathbf{1}_{E(\mathbf{p})}(x)}{\mu(B(x, D^{s(\mathbf{p})}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(x, 16D^{s(\mathbf{p})}) \\ s(J) \leq s(\mathbf{p})}} D^{(s(J)-s(\mathbf{p}))/a} \int_J |f|. \\ &= 2^{127a^3+3/a} \sum_{I \in \mathcal{D}} \sum_{\substack{\mathbf{p} \in \mathfrak{T} \\ \mathcal{J}(\mathbf{p})=I}} \frac{\mathbf{1}_{E(\mathbf{p})}(x)}{\mu(B(x, D^{s(I)}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(x, 16D^{s(I)}) \\ s(J) \leq s(I)}} D^{(s(J)-s(I))/a} \int_J |f|. \end{aligned}$$

By (2.0.13) and (2.0.20), the sets $E(\mathbf{p})$ for tiles \mathbf{p} with $\mathcal{J}(\mathbf{p}) = I$ are pairwise disjoint. It follows from the definition of $\mathcal{L}(\mathfrak{T}(\mathbf{u}))$ that $x \in \mathcal{J}(\mathbf{p})$ if and only if $x' \in \mathcal{J}(\mathbf{p})$, thus we can estimate the sum over $\mathbf{1}_{E(\mathbf{p})}(x)$ by $\mathbf{1}_I(x')$. If $x \in E(\mathbf{p})$ then in particular $x \in \mathcal{J}(\mathbf{p})$, so by (2.0.10) $B(c(I), 16D^{s(I)}) \subset B(x, 32D^{s(I)})$. By the doubling property (1.1.2)

$$\mu(B(c(I), 16D^{s(I)})) \leq 2^{5a} \mu(B(x, D^{s(I)})).$$

Since $a \geq 4$ we can continue our estimate with

$$\leq 2^{128a^3} \sum_{I \in \mathcal{D}} \frac{\mathbf{1}_I(x')}{\mu(B(c(I), 16D^{s(I)}))} \sum_{\substack{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \\ J \subset B(x, 16D^{s(I)}) \\ s(J) \leq s(I)}} D^{(s(J)-s(I))/a} \int_J |f|$$

$$= 2^{128a^3} S_{1,u} P_{\mathcal{J}(\mathfrak{T}(u))} |f|(x').$$

This completes the proof. \square

7.2. An auxiliary L^2 tree estimate. In this subsection we prove the following estimate on L^2 for operators associated to trees.

Lemma 7.2.1 (tree projection estimate). *Let $u \in \mathfrak{U}$. Then we have for all f, g bounded with bounded support*

$$\begin{aligned} & \left| \int_X \sum_{\mathfrak{p} \in \mathfrak{T}(u)} \bar{g}(y) T_{\mathfrak{p}} f(y) \, d\mu(y) \right| \\ & \leq 2^{130a^3} \|P_{\mathcal{J}(\mathfrak{T}(u))} |f|\|_2 \|P_{\mathcal{L}(\mathfrak{T}(u))} |g|\|_2. \end{aligned} \quad (7.2.1)$$

Below, we deduce Lemma 7.2.1 from Lemma 7.1.3 and the following estimates for the operators in Lemma 7.1.3.

Lemma 7.2.2 (nontangential operator bound). *For all bounded f with bounded support and all $\vartheta \in \Theta$*

$$\|T_{\mathcal{N}}^{\vartheta} f\|_2 \leq 2^{102a^3} \|f\|_2.$$

Lemma 7.2.3 (boundary operator bound). *For all $u \in \mathfrak{U}$ and all bounded functions f with bounded support*

$$\|S_{1,u} f\|_2 \leq 2^{12a} \|f\|_2. \quad (7.2.2)$$

Proof of Lemma 7.2.1. Let $L \in \mathcal{L}(\mathfrak{T}(u))$. Let $b(x')$ denote the right-hand side of Lemma 7.1.3. Apply this lemma to $e(\mathcal{Q}(u))f$, to obtain for all $y, x' \in L$

$$\left| \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}} f(y) \right| \leq b(x').$$

Hence, by taking an infimum, we have for $y \in L$

$$\left| \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}} f(y) \right| \leq \inf_{x' \in L} b(x').$$

Integrating this estimate yields

$$\begin{aligned} & \int_L |g(y)| \left| \sum_{\mathfrak{p} \in \mathfrak{T}(u)} T_{\mathfrak{p}} f(y) \right| \, d\mu(y) \\ & \leq \int_L |g(y)| \times \inf_{x' \in L} b(x') \, d\mu(y) \\ & = \int_L P_{\mathcal{L}(\mathfrak{T}(u))} |g|(y) \times \inf_{x' \in L} b(x') \, d\mu(y) \\ & \leq \int_L P_{\mathcal{L}(\mathfrak{T}(u))} |g|(y) \times b(y) \, d\mu(y) \end{aligned}$$

By (2.0.21), we have $T_{\mathfrak{p}}f = \mathbf{1}_{\mathcal{J}(\mathfrak{p})}T_{\mathfrak{p}}f$ for all $\mathfrak{p} \in \mathfrak{P}$, so

$$\left| \int \bar{g}(y) \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}f(y) d\mu(y) \right| = \left| \int_{\bigcup_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} \mathcal{J}(\mathfrak{p})} \bar{g}(y) \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}f(y) d\mu(y) \right|.$$

Since $\mathcal{L}(\mathfrak{T}(\mathfrak{u}))$ partitions $\bigcup_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} \mathcal{J}(\mathfrak{p})$ by Lemma 7.1.2, we get from the triangle inequality

$$\leq \sum_{L \in \mathcal{L}(\mathfrak{T}(\mathfrak{u}))} \int_L |g(y)| \left| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}f(y) \right| d\mu(y)$$

which by the above computation is bounded by

$$\begin{aligned} & \sum_{L \in \mathcal{L}(\mathfrak{T}(\mathfrak{u}))} \int_L P_{\mathcal{L}(\mathfrak{T}(\mathfrak{u}))} |g|(y) \times b(y) d\mu(y) \\ &= \int_X P_{\mathcal{L}(\mathfrak{T}(\mathfrak{u}))} |g|(y) \times b(y) d\mu(y) \end{aligned}$$

Applying Cauchy-Schwarz, this is bounded by $\|P_{\mathcal{L}(\mathfrak{T}(\mathfrak{u}))}|g|\|_2 \times \|b\|_2$. By Minkowski's inequality, Proposition 2.0.6, Lemma 7.2.2 and Lemma 7.2.3, $\|b\|_2$ is at most

$$2^{129a^3} (2^{2a+1} + 2^{12a}) \|P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))}|f|\|_2 + 2^{103a^3} \|P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))}[e(Q(\mathfrak{u}))f]\|_2.$$

By the triangle inequality we have for all $x \in X$ that $|P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))}[e(Q(\mathfrak{u}))f]|(x) \leq P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))}|f|(x)$, thus we can further estimate the above by

$$(2^{129a^3} (2^{2a+1} + 2^{12a}) + 2^{103a^3}) \|P_{\mathcal{J}(\mathfrak{T}(\mathfrak{u}))}|f|\|_2.$$

This completes the proof since $a \geq 4$. \square

Now we prove the two auxiliary lemmas. We begin with the nontangential maximal operator $T_{\mathcal{N}}$.

Proof of Lemma 7.2.2. Fix s_1, s_2 . By (2.0.4) we have for all $x \in (0, \infty)$

$$\sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 1 - \sum_{s < s_1} \psi(D^{-s}x) - \sum_{s > s_1} \psi(D^{-s}x).$$

Since ψ is supported in $[\frac{1}{4D}, \frac{1}{2}]$, the two sums on the right hand side are zero for all $x \in [\frac{1}{2}D^{s_1-1}, \frac{1}{4}D^{s_2-1}]$, hence

$$x \in [\frac{1}{2}D^{s_1-1}, \frac{1}{4}D^{s_2}] \implies \sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 1.$$

Since ψ is supported in $[\frac{1}{4D}, \frac{1}{2}]$, we further have

$$x \notin [\frac{1}{4}D^{s_1-1}, \frac{1}{2}D^{s_2}] \implies \sum_{s=s_1}^{s_2} \psi(D^{-s}x) = 0.$$

Finally, since $\psi \geq 0$ and $\sum_{s \in \mathbb{Z}} \psi(D^{-s}x) = 1$, we have for all x

$$0 \leq \sum_{s=s_1}^{s_2} \psi(D^{-s}x) \leq 1.$$

Let $x' \in I_{s_1}(x)$ and suppose that $D^{s_2-1} \leq R_Q(\vartheta, x')$. By the triangle inequality and (2.0.10), it holds that $\rho(x, x') \leq 8D^{s_1}$. We have

$$\begin{aligned} & \left| \sum_{s=s_1}^{s_2} \int K_s(x', y) f(y) \, d\mu(y) \right| \\ &= \left| \int \sum_{s=s_1}^{s_2} \psi(D^{-s}\rho(x', y)) K(x', y) f(y) \, d\mu(y) \right| \\ &\leq \left| \int_{8D^{s_1} < \rho(x', y) \leq \frac{1}{4}D^{s_2-1}} K(x', y) f(y) \, d\mu(y) \right| \end{aligned} \quad (7.2.3)$$

$$+ \int_{\frac{1}{4}D^{s_1-1} \leq \rho(x', y) \leq 8D^{s_1}} |K(x', y)| |f(y)| \, d\mu(y) \quad (7.2.4)$$

$$+ \int_{\frac{1}{4}D^{s_2-1} \leq \rho(x', y) \leq \frac{1}{2}D^{s_2}} |K(x', y)| |f(y)| \, d\mu(y). \quad (7.2.5)$$

The first term (7.2.3) is at most $2T_Q^\vartheta f(x)$, using with $R_1 := 8D^{s_1}$, $R_2 := \frac{1}{4}D^{s_2-1}$ and $R_1 < R_2 < R_Q(\vartheta, x')$ the triangle inequality in the form

$$\left| \int_{R_1 < \rho(x', y) \leq R_2} K(x', y) f(y) \, d\mu(y) \right| \quad (7.2.6)$$

$$\leq \left| \int_{R_1 < \rho(x', y) < R_Q(\vartheta, x')} K(x', y) f(y) \, d\mu(y) \right| \quad (7.2.7)$$

$$+ \left| \int_{R_2 < \rho(x', y) < R_Q(\vartheta, x')} K(x', y) f(y) \, d\mu(y) \right|. \quad (7.2.8)$$

The other two terms will be estimated by the finitary maximal function from Proposition 2.0.6. For the second term (7.2.4) we use (1.1.11) which implies that for all y with $\rho(x', y) \geq \frac{1}{4}D^{s_1-1}$, we have

$$|K(x', y)| \leq \frac{2^{a^3}}{\mu(B(x', \frac{1}{4}D^{s_1-1}))}.$$

Using $D = 2^{100a^2}$ and the doubling property (1.1.2) $7 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{7a+101a^3}}{\mu(B(x', 32D^{s_1}))}. \quad (7.2.9)$$

By the triangle inequality and (2.0.10), we have

$$B(x', 8D^{s_1}) \subset B(c(I_{s_1}(x)), 16D^{s(I_{s_1}(x))}) \subset B(x', 32D^{s_1}).$$

Combining this with (7.2.9), we conclude that (7.2.4) is at most

$$2^{7a+101a^3} M_{\mathcal{B},1} f(x).$$

For (7.2.5) we argue similarly. We have for all y with $\rho(x', y) \geq \frac{1}{4}D^{s_2-1}$

$$|K(x', y)| \leq \frac{2^{a^3}}{\mu(B(x', \frac{1}{4}D^{s_2-1}))}.$$

Using the doubling property (1.1.2) $7 + 100a^2$ times estimates the last display by

$$\leq \frac{2^{7a+101a^3}}{\mu(B(x', 32D^{s_2}))}. \quad (7.2.10)$$

Note that by (2.0.8) we have $I_{s_1}(x) \subset I_{s_2}(x)$, in particular $x' \in I_{s_2}(x)$. By the triangle inequality and (2.0.10), we have

$$B(x', 8D^{s_2}) \subset B(c(I_{s_2}(x)), 16D^{s(I_{s_2}(x))}) \subset B(x', 32D^{s_2}).$$

Combining this, (7.2.5) is at most

$$2^{7a+101a^3} M_{\mathcal{B},1} f(x).$$

Using $a \geq 4$, taking a supremum over all $x' \in I_{s_1}(x)$ and then a supremum over all $-S \leq s_1 < s_2 \leq S$, we obtain

$$T_{\mathcal{N}} f(x) \leq 2T_Q^\vartheta f(x) + 2^{102a^3} M_{\mathcal{B},1} f(x).$$

The lemma now follows from assumption (1.1.20), Proposition 2.0.6 and $a \geq 4$. \square

We need the following lemma to prepare the L^2 -estimate for the auxiliary operators $S_{1,u}$.

Lemma 7.2.4 (boundary overlap). *For every cube $I \in \mathcal{D}$, there exist at most 2^{9a} cubes $J \in \mathcal{D}$ with $s(J) = s(I)$ and $B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset$.*

Proof. Suppose that $B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset$ and $s(I) = s(J)$. Then $B(c(I), 33D^{s(I)}) \subset B(c(J), 128D^{s(J)})$. Hence by the doubling property (1.1.2)

$$2^{9a} \mu(B(c(J), \frac{1}{4}D^{s(J)})) \geq \mu(B(c(I), 33D^{s(I)})),$$

and by the triangle inequality, $B(c(J), \frac{1}{4}D^{s(J)})$ is contained in $B(c(I), 33D^{s(I)})$.

If \mathcal{C} is any finite collection of cubes $J \in \mathcal{D}$ satisfying $s(J) = s(I)$ and

$$B(c(I), 16D^{s(I)}) \cap B(c(J), 16D^{s(J)}) \neq \emptyset,$$

then it follows from (2.0.10) and pairwise disjointness of cubes of the same scale (2.0.8) that the balls $B(c(J), \frac{1}{4}D^{s(J)})$ are pairwise disjoint. Hence

$$\begin{aligned} \mu(B(c(I), 33D^{s(I)})) &\geq \sum_{J \in \mathcal{C}} \mu(B(c(J), \frac{1}{4}D^{s(J)})) \\ &\geq |\mathcal{C}| 2^{-9a} \mu(B(c(I), 33D^{s(I)})). \end{aligned}$$

Since μ is doubling and $\mu \neq 0$, we have $\mu(B(c(I), 33D^{s(I)})) > 0$. The lemma follows after dividing by $2^{-9a} \mu(B(c(I), 33D^{s(I)}))$. \square

Now we can bound the operators $S_{1,u}$.

Proof of Lemma 7.2.3. Note that by definition, $S_{1,u}f$ is a finite sum of indicator functions of cubes $I \in \mathcal{D}$ for each locally integrable f , and hence is bounded, has bounded support and is integrable. Let g be another function with the same three properties. Then $\bar{g}S_{1,u}f$ is integrable, and we have

$$\begin{aligned} & \left| \int \bar{g}(y) S_{1,u}f(y) \, d\mu(y) \right| \\ &= \left| \sum_{I \in \mathcal{D}} \frac{1}{\mu(B(c(I), 16D^s(I)))} \int_I \bar{g}(y) \, d\mu(y) \right. \\ & \quad \times \left. \sum_{\substack{J \in \mathcal{J}(\mathfrak{X}(u)) : J \subseteq B(c(I), 16D^s(I)) \\ s(J) \leq s(I)}} D^{(s(J)-s(I))/a} \int_J |f(y)| \, d\mu(y) \right| \\ &\leq \sum_{I \in \mathcal{D}} \frac{1}{\mu(B(c(I), 16D^s(I)))} \int_{B(c(I), 16D^s(I))} |g(y)| \, d\mu(y) \\ & \quad \times \sum_{\substack{J \in \mathcal{J}(\mathfrak{X}(u)) : J \subseteq B(c(I), 16D^s(I)) \\ s(J) \leq s(I)}} D^{(s(J)-s(I))/a} \int_J |f(y)| \, d\mu(y). \end{aligned}$$

Changing the order of summation and using $J \subset B(c(I), 16D^s(I))$ to bound the first average integral by $M_{\mathcal{B},1}|g|(y)$ for any $y \in J$, we obtain

$$\leq \sum_{J \in \mathcal{J}(\mathfrak{X}(u))} \int_J |f(y)| M_{\mathcal{B},1} |g|(y) \, d\mu(y) \sum_{\substack{I \in \mathcal{D} : J \subseteq B(c(I), 16D^s(I)) \\ s(I) \geq s(J)}} D^{(s(J)-s(I))/a}. \quad (7.2.11)$$

By Lemma 7.2.4, there are at most 2^{9a} cubes I at each scale satisfying the inclusion $J \subset B(c(I), 16D^s(I))$. Since $D^{-1/a} \leq \frac{1}{2}$, $(1 - D^{-1/a})^{-1} \leq 2$. Using this estimate for the sum of the geometric series, we conclude that (7.2.11) is at most

$$2^{9a+1} \sum_{J \in \mathcal{J}(\mathfrak{X}(u))} \int_J |f(y)| M_{\mathcal{B},1} |g|(y) \, d\mu(y).$$

The collection \mathcal{J} is a partition of $\bigcup_{I \in \mathcal{D}} I$, so this equals

$$2^{9a+1} \int_X |f(y)| M_{\mathcal{B},1} |g|(y) \, d\mu(y).$$

Using Cauchy-Schwarz and Proposition 2.0.6, we conclude

$$\left| \int \bar{g} S_{1,u}f \, d\mu \right| \leq 2^{12a} \|g\|_2 \|f\|_2.$$

The lemma now follows by choosing $g = S_{1,u}f$ and dividing on both sides by the finite $\|S_{1,u}f\|_2$. \square

7.3. The quantitative L^2 tree estimate. The main result of this subsection is the following quantitative bound for operators associated to trees, with decay in the densities dens_1 and dens_2 .

Lemma 7.3.1 (densities tree bound). *Let $\mathbf{u} \in \mathfrak{U}$. Then for all bounded f with bounded support and bounded g supported on G we have*

$$\left| \int_X \bar{g} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}} f \, d\mu \right| \leq 2^{181a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2 \|g\|_2. \quad (7.3.1)$$

If additionally $\text{support}(f) \subseteq F$, then we have

$$\left| \int_X \bar{g} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}} f \, d\mu \right| \leq 2^{282a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \text{dens}_2(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2 \|g\|_2. \quad (7.3.2)$$

Below, we deduce this lemma from Lemma 7.2.1 and the following two estimates controlling the size of support of the operator and its adjoint.

Lemma 7.3.2 (local dens1 tree bound). *Let $\mathbf{u} \in \mathfrak{U}$ and $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$. Then*

$$\mu(L \cap G \cap \bigcup_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} E(\mathfrak{p})) \leq 2^{101a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u})) \mu(L). \quad (7.3.3)$$

Lemma 7.3.3 (local dens2 tree bound). *Let $\mathbf{u} \in \mathfrak{U}$ and $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))$. Then*

$$\mu(F \cap J) \leq 2^{201a^3} \text{dens}_2(\mathfrak{T}(\mathbf{u})) \mu(J).$$

Proof of Lemma 7.3.1. Denote

$$\mathcal{E}(\mathbf{u}) = \bigcup_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} E(\mathfrak{p}).$$

Then we have

$$\left| \int_X \bar{g} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}} f \, d\mu \right| = \left| \int_X \overline{g \mathbf{1}_{\mathcal{E}(\mathbf{u})}} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}} (\mathbf{1}_{\mathcal{J}(\mathbf{u})} f) \, d\mu \right|.$$

By Lemma 7.2.1, this is bounded by

$$\leq 2^{130a^3} \|P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))} |f|\|_2 \|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}))} |\mathbf{1}_{\mathcal{E}(\mathbf{u})} g|\|_2. \quad (7.3.4)$$

We bound the two factors separately. We have

$$\|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}))} |\mathbf{1}_{\mathcal{E}(\mathbf{u})} g|\|_2 = \left(\sum_{L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(L)} \left(\int_{L \cap \mathcal{E}(\mathbf{u})} |g(y)| \, d\mu(y) \right)^2 \right)^{1/2}.$$

By Cauchy-Schwarz and Lemma 7.3.2 this is at most

$$\leq \left(\sum_{L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))} 2^{101a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u})) \int_{L \cap \mathcal{E}(\mathbf{u})} |g(y)|^2 \, d\mu(y) \right)^{1/2}.$$

Since cubes $L \in \mathcal{L}(\mathfrak{T}(\mathbf{u}))$ are pairwise disjoint by Lemma 7.1.2, this is

$$\leq 2^{51a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))^{1/2} \|g\|_2. \quad (7.3.5)$$

Similarly, we have

$$\|P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}))}|f\|_2 = \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(J)} \left(\int_J |f(y)| \, d\mu(y) \right)^2 \right)^{1/2}. \quad (7.3.6)$$

By Cauchy-Schwarz, this is

$$\leq \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \int_J |f(y)|^2 \, d\mu(y) \right)^{1/2}.$$

Since cubes in $\mathcal{J}(\mathfrak{T}(\mathbf{u}))$ are pairwise disjoint by Lemma 7.1.2, this is at most

$$\|f\|_2. \quad (7.3.7)$$

Combining (7.3.4), (7.3.5) and (7.3.7) gives (7.3.1).

If $f \leq \mathbf{1}_F$ then $f = f\mathbf{1}_F$, so

$$\begin{aligned} & \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(J)} \left(\int_J |f(y)| \, d\mu(y) \right)^2 \right)^{1/2} \\ &= \left(\sum_{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}))} \frac{1}{\mu(J)} \left(\int_{J \cap F} |f(y)| \, d\mu(y) \right)^2 \right)^{1/2}. \end{aligned}$$

We estimate as before, using now Lemma 7.3.3 and Cauchy-Schwarz, and obtain that this is

$$\leq 2^{101a^3} \text{dens}_2(\mathfrak{T}(\mathbf{u}))^{1/2} \|f\|_2.$$

Combining this with (7.3.4) and (7.3.5) gives (7.3.2). \square

Now we prove the two auxiliary estimates.

Proof of Lemma 7.3.2. If the set on the right hand side is empty, then (7.3.3) holds. If not, then there exists $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $L \cap \mathcal{J}(\mathbf{p}) \neq \emptyset$.

Suppose first that there exists such \mathbf{p} with $s(\mathbf{p}) \leq s(L)$. Then by (2.0.8) $\mathcal{J}(\mathbf{p}) \subset L$, which gives by the definition of $\mathcal{L}(\mathfrak{T}(\mathbf{u}))$ that $s(L) = -S$ and hence $L = \mathcal{J}(\mathbf{p})$. Let $\mathbf{q} \in \mathfrak{T}(\mathbf{u})$ with $E(\mathbf{q}) \cap L \neq \emptyset$. Since $s(L) = -S \leq s(\mathbf{q})$ it follows from (2.0.8) that $\mathcal{J}(\mathbf{p}) = L \subset \mathcal{J}(\mathbf{q})$. We have then by Lemma 2.1.2

$$\begin{aligned} d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{q})) &\leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u})) + d_{\mathbf{p}}(\mathcal{Q}(\mathbf{q}), \mathcal{Q}(\mathbf{u})) \\ &\leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u})) + d_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), \mathcal{Q}(\mathbf{u})). \end{aligned}$$

Using that $\mathbf{p}, \mathbf{q} \in \mathfrak{T}(\mathbf{u})$ and (2.0.32), this is at most 8. Using again the triangle inequality and Lemma 2.1.2, we obtain that for each $q \in B_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), 1)$

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), q) \leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{q})) + d_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), q) \leq 9.$$

Thus $L \cap G \cap E(\mathbf{q}) \subset E_2(9, \mathbf{p})$. We obtain

$$\mu(L \cap G \cap \bigcup_{\mathbf{q} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{q})) \leq \mu(E_2(9, \mathbf{p})).$$

By the definition of dens_1 , this is bounded by

$$9^a \text{dens}_1(\mathfrak{T}(\mathbf{u}))\mu(\mathcal{J}(\mathbf{p})) = 9^a \text{dens}_1(\mathfrak{T}(\mathbf{u}))\mu(L).$$

Since $a \geq 4$, (7.3.3) follows in this case.

Now suppose that for each $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ with $L \cap E(\mathbf{p}) \neq \emptyset$, we have $s(\mathbf{p}) > s(L)$. Since there exists at least one such \mathbf{p} , there exists in particular at least one cube $L'' \in \mathcal{D}$ with $L \subset L''$ and $s(L'') > s(L)$. By (2.0.7), there exists $L' \in \mathcal{D}$ with $L \subset L'$ and $s(L') = s(L) + 1$. By the definition of $\mathcal{L}(\mathfrak{T}(\mathbf{u}))$ there exists a tile $\mathbf{p}'' \in \mathfrak{T}(\mathbf{u})$ with $\mathcal{J}(\mathbf{p}'') \subset L'$.

It suffices to show that there exists a tile $\mathbf{p}' \in \mathfrak{P}(\mathfrak{T}(\mathbf{u}))$ with $\mathcal{J}(\mathbf{p}') = L'$, $d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{u})) < 4$ and $9\mathbf{p}'' \lesssim 9\mathbf{p}'$. For then, let $\mathbf{q} \in \mathfrak{T}(\mathbf{u})$ with $L \cap E(\mathbf{q}) \neq \emptyset$. As shown above, this implies $s(\mathbf{q}) \geq s(L')$, so by (2.0.8) $L' \subset \mathcal{J}(\mathbf{q})$. If $q \in B_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), 1)$, then by a similar calculation as above, using the triangle inequality, Lemma 2.1.2 and (2.0.32), we obtain

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), q) \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{q})) + d_{\mathbf{q}}(\mathcal{Q}(\mathbf{q}), q) \leq 9.$$

Thus $L \cap G \cap E(\mathbf{q}) \subset E_2(9, \mathbf{p}')$. We deduce using the definition (2.0.28) of dens_1

$$\mu(L \cap G \cap \bigcup_{\mathbf{q} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{q})) \leq \mu(E_2(9, \mathbf{p}')) \leq 9^a \text{dens}_1(\mathfrak{T}(\mathbf{u}))\mu(L').$$

Using the doubling property (1.1.2), (2.0.10), and $a \geq 4$ this is estimated by

$$9^a 2^{100a^3+5a} \text{dens}_1(\mathfrak{T}(\mathbf{u}))\mu(L) \leq 2^{101a^3} \text{dens}_1(\mathfrak{T}(\mathbf{u}))\mu(L).$$

To show existence of \mathbf{p}' with the given properties, if $\mathcal{J}(\mathbf{p}'') = L'$ we can take $\mathbf{p}' = \mathbf{p}''$, which satisfies the distance property by (2.0.32) and the other properties trivially. Otherwise, let \mathbf{p}' be the unique tile such that $\mathcal{J}(\mathbf{p}') = L'$ and such that $\Omega(\mathbf{u}) \cap \Omega(\mathbf{p}') \neq \emptyset$. Since $\mathcal{J}(\mathbf{p}') \subset \mathcal{J}(\mathbf{p})$ and $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$, we have $\mathbf{p}' \in \mathfrak{P}(\mathfrak{T}(\mathbf{u}))$. Since by (2.0.32) $s(\mathbf{p}') = s(L') \leq s(\mathbf{p}) < s(\mathbf{u})$, we have by (2.0.8) and (2.0.14) that $\Omega(\mathbf{u}) \subset \Omega(\mathbf{p}')$, and hence the distance property. $9\mathbf{p}'' \lesssim 9\mathbf{p}'$ follows by the triangle inequality, (2.0.32), Lemma 2.1.2 and (2.0.15). This completes the proof. \square

Proof of Lemma 7.3.3. We prove the inequality with the constant 2^{201a^3} replaced by 2^{200a^3+14a} ; this is stronger because $a \geq 4$. It suffices to show the existence of a tile $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ and an $r \geq 4D^{s(\mathbf{p})}$ such that $J \subset B(c(\mathbf{p}), r)$ and $\mu(B(c(\mathbf{p}), r)) \leq 2^{200a^3+14a}\mu(J)$, because then it follows from the definition (2.0.29) of dens_2 that

$$\begin{aligned} \mu(F \cap J) &\leq \mu(F \cap B(c(\mathbf{p}), r)) \\ &\leq \text{dens}_2(\mathfrak{T}(\mathbf{u}))\mu(B(c(\mathbf{p}), r)) \leq 2^{200a^3+14a} \text{dens}_2(\mathfrak{T}(\mathbf{u}))\mu(J). \end{aligned}$$

In particular, these criteria are satisfied, with $r = 4D^{s(\mathbf{p})}$, by any $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ such that $J \subseteq B(c(\mathbf{p}), 4D^{s(\mathbf{p})})$ and $\mu(\mathcal{J}(\mathbf{p})) \leq 2^{100a^3+10a}\mu(J)$, because then by the doubling property (1.1.2),

$$\begin{aligned} \mu(B(c(\mathbf{p}), 4D^{s(\mathbf{p})})) &\leq 2^{4a}\mu(B(c(\mathbf{p}), D^{s(\mathbf{p})}/4)) \\ &\leq 2^{4a}\mu(\mathcal{J}(\mathbf{p})) \leq 2^{100a^3+14a}\mu(J). \end{aligned}$$

Suppose first that $s(J) = S$. Then $J = I_0$, so (2.0.9) and the fact that $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u})) \subseteq \mathcal{J}_0(\mathfrak{T}(\mathbf{u}))$ imply that $s(J) = -S$. Thus $S = 0$. It follows that J is the only

dyadic cube, so any $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ has $\mathcal{J}(\mathbf{p}) = J$, and therefore satisfies $J \subseteq B(c(\mathbf{p}), 4D^{s(\mathbf{p})})$ and $\mu(\mathcal{J}(\mathbf{p})) \leq 2^{100a^3+10a}\mu(J)$.

It remains to consider the case $s(J) < S$. Then, by (2.0.7) and (2.0.8), there exists some cube $J' \in \mathcal{D}$ with $s(J') = s(J) + 1$ and $J \subset J'$. By definition of $\mathcal{J}(\mathfrak{T}(\mathbf{u}))$ there exists some $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ such that $\mathcal{J}(\mathbf{p}) \subset B(c(J'), 100D^{s(J')+1})$.

Since $c(J) \in J \subset J' \subset B(c(J'), 4D^{s(J')})$, the triangle inequality, $s(J') = s(J) + 1$ and $D = 2^{100a^2}$ imply

$$B(c(J'), 204D^{s(J')+1}) \subset B(c(J), 204D^{s(J')+1} + 4D^{s(J')}) \subset B(c(J), 2^8 D^{s(J)+2}).$$

From the doubling property (1.1.2), $D = 2^{100a^2}$ and (2.0.10), we obtain

$$\mu(B(c(J'), 204D^{s(J')+1})) \leq 2^{200a^3+10a}\mu(J). \quad (7.3.8)$$

If $J \subset B(c(\mathbf{p}), 4D^{s(\mathbf{p})})$, then we need only check that $\mu(\mathcal{J}(\mathbf{p})) \leq 2^{100a^3+10a}\mu(J)$. This follows immediately from $\mathcal{J}(\mathbf{p}) \subset B(c(J'), 100D^{s(J')+1})$ and (7.3.8).

From now on we assume $J \not\subset B(c(\mathbf{p}), 4D^{s(\mathbf{p})})$. Since

$$c(\mathbf{p}) \in \mathcal{J}(\mathbf{p}) \subset B(c(J'), 100D^{s(J')+1}),$$

we have by (2.0.10) and the triangle inequality

$$J \subset J' \subset B(c(J'), 4D^{s(J')}) \subset B(c(\mathbf{p}), 104D^{s(J')+1}).$$

In particular this implies $104D^{s(J')+1} > 4D^{s(\mathbf{p})}$. By the triangle inequality we also have

$$B(c(\mathbf{p}), 104D^{s(J')+1}) \subset B(c(J), 204D^{s(J')+1}),$$

so from (7.3.8),

$$\mu(B(c(\mathbf{p}), 104D^{s(J')+1})) \leq 2^{200a^3+10a}\mu(J),$$

which proves \mathbf{p} satisfies the needed criteria with $r = 104D^{s(J')+1}$. □

7.4. Almost orthogonality of separated trees. The main result of this subsection is the almost orthogonality estimate for operators associated to distinct trees in a forest in Lemma 7.4.4 below. We will deduce it from Lemmas 7.4.5 and 7.4.6, which are proven in Subsections 7.5 and 7.6, respectively. Before stating it, we introduce some relevant notation.

The adjoint of the operator $T_{\mathbf{p}}$ defined in (2.0.21) is given by

$$T_{\mathbf{p}}^*g(x) = \int_{E(\mathbf{p})} \overline{K_{s(\mathbf{p})}(y, x)} e(-Q(y)(x) + Q(y)(y))g(y) d\mu(y). \quad (7.4.1)$$

Lemma 7.4.1 (adjoint tile support). *For each $\mathbf{p} \in \mathfrak{P}$, we have*

$$T_{\mathbf{p}}^*g = \mathbf{1}_{B(c(\mathbf{p}), 5D^{s(\mathbf{p})})} T_{\mathbf{p}}^* \mathbf{1}_{\mathcal{J}(\mathbf{p})} g.$$

For each $\mathbf{u} \in \mathfrak{U}$ and each $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$, we have

$$T_{\mathbf{p}}^*g = \mathbf{1}_{\mathcal{J}(\mathbf{u})} T_{\mathbf{p}}^* \mathbf{1}_{\mathcal{J}(\mathbf{u})} g.$$

Proof. By (2.0.32), $E(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{u})$. Thus by (7.4.1)

$$\begin{aligned} T_{\mathfrak{p}}^* g(x) &= T_{\mathfrak{p}}^*(\mathbf{1}_{\mathcal{J}(\mathfrak{p})} g)(x) \\ &= \int_{E(\mathfrak{p})} \overline{K_{s(\mathfrak{p})}(y, x)} e(-Q(y)(x) + Q(y)(y)) \mathbf{1}_{\mathcal{J}(\mathfrak{p})}(y) g(y) d\mu(y). \end{aligned}$$

If this integral is not 0, then there exists $y \in \mathcal{J}(\mathfrak{p})$ such that $K_{s(\mathfrak{p})}(y, x) \neq 0$. By (2.1.2), (2.0.10) and the triangle inequality, it follows that

$$x \in B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})}).$$

Thus

$$T_{\mathfrak{p}}^* g(x) = \mathbf{1}_{B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})}(x) T_{\mathfrak{p}}^*(\mathbf{1}_{\mathcal{J}(\mathfrak{p})} g)(x).$$

The second claimed equation follows now since $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathfrak{u})$ and by (2.0.37) we have $B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})}) \subset \mathcal{J}(\mathfrak{u})$. \square

Lemma 7.4.2 (adjoint tree estimate). *For all bounded g supported on G we have that*

$$\begin{aligned} \left\| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g \right\|_2 &\leq 2^{181a^3} \text{dens}_1(\mathfrak{T}(\mathfrak{u}))^{1/2} \|g\|_2, \\ \left\| \mathbf{1}_F \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g \right\|_2 &\leq 2^{282a^3} \text{dens}_1(\mathfrak{T}(\mathfrak{u}))^{1/2} \text{dens}_2(\mathfrak{T}(\mathfrak{u}))^{1/2} \|g\|_2. \end{aligned}$$

Proof. By Lemma 7.3.1, we have for all bounded f and g with $|g| \leq \mathbf{1}_G$ that

$$\begin{aligned} \left| \int_X \overline{\sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g} f d\mu \right| &= \left| \int_X \bar{g} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}} f d\mu \right| \\ &\leq 2^{181a^3} \text{dens}_1(\mathfrak{T}(\mathfrak{u}))^{1/2} \|g\|_2 \|f\|_2. \end{aligned} \quad (7.4.2)$$

Let $f = \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g$. Since $|g| \leq \mathbf{1}_G$, f is bounded and has bounded support. In particular $\|f\|_2 < \infty$. Dividing (7.4.2) by $\|f\|_2$ completes the proof.

The proof of the second part is similar with $f = \mathbf{1}_F \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g$. \square

We define

$$S_{2,\mathfrak{u}} g := \left| \sum_{\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})} T_{\mathfrak{p}}^* g \right| + M_{\mathcal{B},1} g + |g|.$$

Lemma 7.4.3 (adjoint tree control). *We have for all $\mathfrak{u} \in \mathfrak{U}$ and all bounded g supported on G*

$$\|S_{2,\mathfrak{u}} g\|_2 \leq 2^{182a^3} \|g\|_2.$$

Proof. This follows immediately from Minkowski's inequality, Proposition 2.0.6 and Lemma 7.4.2, using that $a \geq 4$. \square

Now we are ready to state the main result of this subsection.

Lemma 7.4.4 (correlation separated trees). *For any $\mathbf{u}_1 \neq \mathbf{u}_2 \in \mathfrak{U}$ and all bounded g_1, g_2 with bounded support, we have*

$$\left| \int_X \sum_{\mathbf{p}_1 \in \mathfrak{T}(\mathbf{u}_1)} \sum_{\mathbf{p}_2 \in \mathfrak{T}(\mathbf{u}_2)} T_{\mathbf{p}_1}^* g_1 \overline{T_{\mathbf{p}_2}^* g_2} d\mu \right| \quad (7.4.3)$$

$$\leq 2^{512a^3 - 4n} \prod_{j=1}^2 \|S_{2, \mathbf{u}_j} g_j\|_{L^2(\mathcal{J}(\mathbf{u}_1) \cap \mathcal{J}(\mathbf{u}_2))}. \quad (7.4.4)$$

Proof of Lemma 7.4.4. By Lemma 7.4.1 and (2.0.8), the left hand side (7.4.3) is 0 unless $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$ or $\mathcal{J}(\mathbf{u}_2) \subset \mathcal{J}(\mathbf{u}_1)$. Without loss of generality we assume that $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$.

Define

$$\mathfrak{S} := \{\mathbf{p} \in \mathfrak{T}(\mathbf{u}_1) \cup \mathfrak{T}(\mathbf{u}_2) : d_{\mathbf{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \geq 2^{Zn/2}\}. \quad (7.4.5)$$

Lemma 7.4.4 follows by combining the definition (2.0.3) of Z with the following two lemmas. \square

Lemma 7.4.5 (correlation distant tree parts). *We have for all $\mathbf{u}_1 \neq \mathbf{u}_2 \in \mathfrak{U}$ with $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$ and all bounded g_1, g_2 with bounded support*

$$\left| \int_X \sum_{\mathbf{p}_1 \in \mathfrak{T}(\mathbf{u}_1)} \sum_{\mathbf{p}_2 \in \mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}} T_{\mathbf{p}_1}^* g_1 \overline{T_{\mathbf{p}_2}^* g_2} d\mu \right| \quad (7.4.6)$$

$$\leq 2^{511a^3} 2^{-Zn/(4a^2+2a^3)} \prod_{j=1}^2 \|S_{2, \mathbf{u}_j} g_j\|_{L^2(\mathcal{J}(\mathbf{u}_1))}. \quad (7.4.7)$$

Lemma 7.4.6 (correlation near tree parts). *We have for all $\mathbf{u}_1 \neq \mathbf{u}_2 \in \mathfrak{U}$ with $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$ and all bounded g_1, g_2 with bounded support*

$$\left| \int_X \sum_{\mathbf{p}_1 \in \mathfrak{T}(\mathbf{u}_1)} \sum_{\mathbf{p}_2 \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}} T_{\mathbf{p}_1}^* g_1 \overline{T_{\mathbf{p}_2}^* g_2} d\mu \right| \quad (7.4.8)$$

$$\leq 2^{232a^3 + 21a + 5} 2^{-\frac{25}{161a} Zn \kappa} \prod_{j=1}^2 \|S_{2, \mathbf{u}_j} g_j\|_{L^2(\mathcal{J}(\mathbf{u}_1))}. \quad (7.4.9)$$

In the proofs of both lemmas, we will need the following observation.

Lemma 7.4.7 (overlap implies distance). *Let $\mathbf{u}_1 \neq \mathbf{u}_2 \in \mathfrak{U}$ with $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$. If $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_1) \cup \mathfrak{T}(\mathbf{u}_2)$ with $\mathcal{J}(\mathbf{p}) \cap \mathcal{J}(\mathbf{u}_1) \neq \emptyset$, then $\mathbf{p} \in \mathfrak{S}$. In particular, we have $\mathfrak{T}(\mathbf{u}_1) \subset \mathfrak{S}$.*

Proof. Suppose first that $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_1)$. Then $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$, by (2.0.32). Thus we have by the separation condition (2.0.36), (2.0.15), (2.0.32) and the triangle inequality

$$\begin{aligned} d_{\mathbf{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) &\geq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}_2)) - d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}_1)) \\ &\geq 2^{Z(n+1)} - 4 \\ &\geq 2^{Zn/2}, \end{aligned}$$

using that $Z = 2^{12a} \geq 4$. Hence $\mathfrak{p} \in \mathfrak{S}$.

Suppose now that $\mathfrak{p} \in \mathfrak{T}(\mathbf{u}_2)$. If $\mathcal{J}(\mathfrak{p}) \subset \mathcal{J}(\mathbf{u}_1)$, then the same argument as above with \mathbf{u}_1 and \mathbf{u}_2 swapped shows $\mathfrak{p} \in \mathfrak{S}$. If $\mathcal{J}(\mathfrak{p}) \not\subset \mathcal{J}(\mathbf{u}_1)$ then, by (2.0.8), $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathfrak{p})$. Pick $\mathfrak{p}' \in \mathfrak{T}(\mathbf{u}_1)$, we have $\mathcal{J}(\mathfrak{p}') \subset \mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathfrak{p})$. Hence, by Lemma 2.1.2 and the first paragraph

$$d_{\mathfrak{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \geq d_{\mathfrak{p}'}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \geq 2^{Zn},$$

so $\mathfrak{p} \in \mathfrak{S}$. □

To simplify the notation, we will write at various places throughout the proof of Lemmas 7.4.5 and 7.4.6 for a subset $\mathfrak{C} \subset \mathfrak{P}$

$$T_{\mathfrak{C}}f := \sum_{\mathfrak{p} \in \mathfrak{C}} T_{\mathfrak{p}}f, \quad T_{\mathfrak{C}}^*g := \sum_{\mathfrak{p} \in \mathfrak{C}} T_{\mathfrak{p}}^*g.$$

7.5. Proof of the Tiles with large separation Lemma. Lemma 7.4.5 follows from the van der Corput estimate in Proposition 2.0.5. We apply this proposition in Section 7.5.3. To prepare this application, we first, in Section 7.5.1, construct a suitable partition of unity, and show then, in Section 7.5.2 the Hölder estimates needed to apply Proposition 2.0.5.

7.5.1. *A partition of unity.* Define

$$\mathcal{J}' = \{J \in \mathcal{J}(\mathfrak{S}) : J \subset \mathcal{J}(\mathbf{u}_1)\}.$$

Lemma 7.5.1 (dyadic partition 1). *We have that*

$$\mathcal{J}(\mathbf{u}_1) = \dot{\bigcup}_{J \in \mathcal{J}'} J.$$

Proof. By Lemma 7.1.2, it remains only to show that each $J \in \mathcal{J}(\mathfrak{S})$ with $J \cap \mathcal{J}(\mathbf{u}_1) \neq \emptyset$ is in \mathcal{J}' . But if $J \notin \mathcal{J}'$, then by (2.0.8) $\mathcal{J}(\mathbf{u}_1) \subset J$. Pick $\mathfrak{p} \in \mathfrak{T}(\mathbf{u}_1) \subset \mathfrak{S}$. Then $\mathcal{J}(\mathfrak{p}) \subset J$. This contradicts the definition of $\mathcal{J}(\mathfrak{S})$. □

For cubes $J \in \mathcal{D}$, denote

$$B(J) := B(c(J), 8D^{s(J)}). \tag{7.5.1}$$

The main result of this subsection is the following.

Lemma 7.5.2 (Lipschitz partition unity). *There exists a family of functions χ_J , $J \in \mathcal{J}'$ such that*

$$\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} = \sum_{J \in \mathcal{J}'} \chi_J, \tag{7.5.2}$$

and for all $J \in \mathcal{J}'$ and all $y, y' \in \mathcal{J}(\mathbf{u}_1)$

$$0 \leq \chi_J(y) \leq \mathbf{1}_{B(J)}(y), \tag{7.5.3}$$

$$|\chi_J(y) - \chi_J(y')| \leq 2^{227a^3} \frac{\rho(y, y')}{D^{s(J)}}. \tag{7.5.4}$$

In the proof, we will use the following auxiliary lemma.

Lemma 7.5.3 (moderate scale change). *If $J, J' \in \mathcal{J}'$ with*

$$B(J) \cap B(J') \neq \emptyset,$$

then $|s(J) - s(J')| \leq 1$.

Proof of Lemma 7.5.2. For each cube $J \in \mathcal{J}$ let

$$\tilde{\chi}_J(y) = \mathbf{1}_{\mathcal{J}(\mathbf{u}_1)}(y) \max\{0, 8 - D^{-s(J)} \rho(y, c(J))\},$$

and set

$$a(y) = \sum_{J \in \mathcal{J}'} \tilde{\chi}_J(y).$$

We define

$$\chi_J(y) := \frac{\tilde{\chi}_J(y)}{a(y)}.$$

Then, due to (2.0.37) and (7.5.1), the properties (7.5.2) and (7.5.3) are clearly true. Estimate (7.5.4) follows from (7.5.3) if $y, y' \notin B(J)$. Thus we can assume that $y \in B(J)$. We have by the triangle inequality

$$|\chi_J(y) - \chi_J(y')| \leq \frac{|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')|}{a(y)} + \frac{\tilde{\chi}_J(y')|a(y) - a(y')|}{a(y)a(y')}$$

Since $\tilde{\chi}_J(z) \geq 4$ for all $z \in B(c(J), 4D^{s(J)}) \supset J$ and by Lemma 7.5.1, we have that $a(z) \geq 4$ for all $z \in \mathcal{J}(\mathbf{u}_1)$. So we can estimate the above further by

$$\leq 2^{-2}(|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')| + \tilde{\chi}_J(y')|a(y) - a(y')|).$$

If $y' \notin B(c(\mathbf{p}), 8D^{s(\mathbf{p})})$ then the second summand vanishes. Else, we can estimate the above, using also that $|\tilde{\chi}_J(y')| \leq 8$, by

$$\leq 2^{-2}(|\tilde{\chi}_J(y) - \tilde{\chi}_J(y')| + 2 \sum_{\substack{J' \in \mathcal{J}' \\ B(J') \cap B(J) \neq \emptyset}} |\tilde{\chi}_{J'}(y) - \tilde{\chi}_{J'}(y')|).$$

By the triangle inequality, we have for all dyadic cubes I

$$|\tilde{\chi}_I(y) - \tilde{\chi}_I(y')| \leq \rho(y, y') D^{-s(I)}.$$

Using this above, we obtain

$$|\chi_J(y) - \chi_J(y')| \leq \rho(y, y') \left(\frac{1}{4} D^{-s(J)} + 2 \sum_{\substack{J' \in \mathcal{J}' \\ B(J') \cap B(J) \neq \emptyset}} D^{-s(J')} \right).$$

By Lemma 7.5.3, this is at most

$$\frac{\rho(y, y')}{D^{s(J)}} \left(\frac{1}{4} + 2D |\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \right).$$

By (2.0.10) and Lemma 7.5.1, the balls $B(c(J'), \frac{1}{4}D^{s(J')})$ are pairwise disjoint. By the triangle inequality and Lemma 7.5.3, each such ball for J' in the set of the last display is contained in

$$B(c(J), 9D^{s(J)+1}).$$

By the doubling property (1.1.2), we further have

$$\mu(B(c(J), 9D^{s(J)+1})) \leq 2^{200a^3+7a} \mu\left(B(c(J'), \frac{1}{4}D^{s(J')})\right)$$

for each such ball. Thus

$$|\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \leq 2^{200a^3+7a}.$$

Recalling that $D = 2^{100a^2}$, we obtain

$$\frac{1}{4} + 2D|\{J' \in \mathcal{J}' : B(J') \cap B(J) \neq \emptyset\}| \leq 2^{200a^3+100a^2+7a+2}.$$

Since $a \geq 4$, (7.5.4) follows. \square

Proof of Lemma 7.5.3. Suppose that $s(J') < s(J) - 1$. Then $s(J) > -S$. Thus, by the definition of \mathcal{J}' there exists no $\mathfrak{p} \in \mathfrak{S}$ with

$$\mathcal{J}(\mathfrak{p}) \subset B(c(J), 100D^{s(J)+1}). \quad (7.5.5)$$

Since $s(J') < s(J)$, there exists a cube $J'' \in \mathcal{D}$ with $J \subset J''$ and $s(J'') = s(J') + 1$. By the definition of \mathcal{J}' , there exists a tile $\mathfrak{p} \in \mathfrak{S}$ with

$$\mathcal{J}(\mathfrak{p}) \subset B(c(J''), 100D^{s(J')+2}). \quad (7.5.6)$$

But by the triangle inequality and (2.0.1), we have

$$B(c(J''), 100D^{s(J')+2}) \subset B(c(J), 100D^{s(J)+1}),$$

which contradicts (7.5.5) and (7.5.6). \square

7.5.2. Hölder estimates for adjoint tree operators. Let $g_1, g_2 : X \rightarrow \mathbb{C}$ be bounded with bounded support. Define for $J \in \mathcal{J}'$

$$h_J(y) := \chi_J(y) \cdot (e(\mathcal{Q}(\mathfrak{u}_1)(y))T_{\mathfrak{T}(\mathfrak{u}_1)}^*g_1(y) \cdot \overline{(e(\mathcal{Q}(\mathfrak{u}_2)(y))T_{\mathfrak{T}(\mathfrak{u}_2) \cap \mathfrak{S}}^*g_2(y))}). \quad (7.5.7)$$

The main result of this subsection is the following τ -Hölder estimate for h_J , where $\tau = 1/a$.

Lemma 7.5.4 (Holder correlation tree). *We have for all $J \in \mathcal{J}'$ that*

$$\|h_J\|_{C^\tau(B(c(J), 16D^{s(J)}))} \leq 2^{485a^3} \prod_{j=1,2} \left(\inf_{B(c(J), \frac{1}{8}D^{s(J)})} |T_{\mathfrak{T}(\mathfrak{u}_j)}^*g_j| + \inf_J M_{\mathcal{B},1}|g_j| \right). \quad (7.5.8)$$

We will prove this lemma at the end of this section, after establishing several auxiliary results.

We begin with the following Hölder continuity estimate for adjoints of operators associated to tiles.

Lemma 7.5.5 (Holder correlation tile). *Let $\mathfrak{u} \in \mathfrak{U}$ and $\mathfrak{p} \in \mathfrak{T}(\mathfrak{u})$. Then for all $y, y' \in X$ and all bounded g with bounded support, we have*

$$\begin{aligned} & |e(\mathcal{Q}(\mathfrak{u})(y))T_{\mathfrak{p}}^*g(y) - e(\mathcal{Q}(\mathfrak{u})(y'))T_{\mathfrak{p}}^*g(y')| \\ & \leq \frac{2^{128a^3}}{\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \int_{E(\mathfrak{p})} |g(x)| d\mu(x). \end{aligned} \quad (7.5.9)$$

Proof. By (7.4.1), we have

$$\begin{aligned}
& |e(\mathcal{Q}(\mathbf{u})(y))T_{\mathbf{p}}^*g(y) - e(\mathcal{Q}(\mathbf{u})(y'))T_{\mathbf{p}}^*g(y')| \\
&= \left| \int_{E(\mathbf{p})} e(Q(x)(x) - Q(x)(y) + \mathcal{Q}(\mathbf{u})(y))\overline{K_{\mathbf{s}(\mathbf{p})}(x, y)}g(x) \right. \\
&\quad \left. - e(Q(x)(x) - Q(x)(y') + \mathcal{Q}(\mathbf{u})(y'))\overline{K_{\mathbf{s}(\mathbf{p})}(x, y')}g(x) \, d\mu(x) \right| \\
&\leq \int_{E(\mathbf{p})} |g(x)| |e(Q(x)(y) - Q(x)(y') - \mathcal{Q}(\mathbf{u})(y) + \mathcal{Q}(\mathbf{u})(y'))\overline{K_{\mathbf{s}(\mathbf{p})}(x, y)} \\
&\quad - \overline{K_{\mathbf{s}(\mathbf{p})}(x, y')}| \, d\mu(x) \\
&\leq \int_{E(\mathbf{p})} |g(x)| |e(-Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y')) - 1| \\
&\quad \times |\overline{K_{\mathbf{s}(\mathbf{p})}(x, y)}| \, d\mu(x) \quad (7.5.10)
\end{aligned}$$

$$+ \int_{E(\mathbf{p})} |g(x)| |\overline{K_{\mathbf{s}(\mathbf{p})}(x, y)} - \overline{K_{\mathbf{s}(\mathbf{p})}(x, y')}| \, d\mu(x). \quad (7.5.11)$$

By the oscillation estimate (1.1.4), we have

$$\begin{aligned}
& | -Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y') | \\
&\leq d_{B(y, 1.6\rho(y, y'))}(Q(x), \mathcal{Q}(\mathbf{u})). \quad (7.5.12)
\end{aligned}$$

Suppose that $y, y' \in B(c(\mathbf{p}), 5D^{\mathbf{s}(\mathbf{p})})$, so that $\rho(y, y') \leq 10D^{\mathbf{s}(\mathbf{p})}$. Let $k \in \mathbb{Z}$ be such that $2^{ak}\rho(y, y') \leq 10D^{\mathbf{s}(\mathbf{p})}$ but $2^{a(k+1)}\rho(y, y') > 10D^{\mathbf{s}(\mathbf{p})}$. In particular, $k \geq 0$. Then, using (1.1.7) followed by (1.1.5), we can bound (7.5.12) from above by

$$2^{-k}d_{B(c(\mathbf{p}), 16D^{\mathbf{s}(\mathbf{p})})}(Q(x), \mathcal{Q}(\mathbf{u})) \leq 2^{6a-k}d_{\mathbf{p}}(Q(x), \mathcal{Q}(\mathbf{u})).$$

Since $x \in E(\mathbf{p})$ we have $Q(x) \in \Omega(\mathbf{p}) \subset B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 1)$, and since $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ we have $\mathcal{Q}(\mathbf{u}) \in B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 4)$, so this is estimated by

$$\leq 5 \cdot 2^{6a-k}.$$

By definition of k , we have

$$-k < 1 - \frac{1}{a} \log_2 \left(\frac{10D^{\mathbf{s}(\mathbf{p})}}{\rho(y, y')} \right),$$

which gives

$$| -Q(x)(y) + Q(x)(y') + \mathcal{Q}(\mathbf{u})(y) - \mathcal{Q}(\mathbf{u})(y') | \leq 10 \cdot 2^{6a} \left(\frac{\rho(y, y')}{10D^{\mathbf{s}(\mathbf{p})}} \right)^{1/a}. \quad (7.5.13)$$

For all $x \in \mathcal{J}(\mathbf{p})$, we have by (1.1.2) that

$$\mu(B(x, D^{\mathbf{s}(\mathbf{p})})) \geq 2^{-3a}\mu(B(c(\mathbf{p}), 4D^{\mathbf{s}(\mathbf{p})})).$$

Combining the above with (2.1.3), (2.1.4) and (7.5.13), we obtain

$$(7.5.10) + (7.5.11) \leq \frac{2^{3a}}{\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \int_{E(\mathfrak{p})} |g(x)| d\mu(x) \times \\ \left(2^{102a^3} \cdot 10 \cdot 2^{6a} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} + 2^{127a^3} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \right)$$

Since $\rho(y, y') \leq 10D^{s(\mathfrak{p})}$, we conclude

$$(7.5.10) + (7.5.11) \leq \frac{2^{128a^3}}{\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \int_{E(\mathfrak{p})} |g(x)| d\mu(x).$$

Next, if $y, y' \notin B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})$, then $T_{\mathfrak{p}}^*g(y) = T_{\mathfrak{p}}^*g(y') = 0$, by Lemma 7.4.1. Then (7.5.9) holds.

Finally, if $y \in B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})$ and $y' \notin B(c(\mathfrak{p}), 5D^{s(\mathfrak{p})})$, then

$$|e(Q(\mathbf{u})(y))T_{\mathfrak{p}}^*g(y) - e(Q(\mathbf{u})(y'))T_{\mathfrak{p}}^*g(y')| = |T_{\mathfrak{p}}^*g(y)| \\ \leq \int_{E(\mathfrak{p})} |K_{s(\mathfrak{p})}(x, y)| |g(x)| d\mu(x).$$

By the same argument used to prove (2.1.6), this is bounded by

$$\leq 2^{102a^3} \int_{E(\mathfrak{p})} \frac{1}{\mu(B(x, D^s))} \psi(D^{-s}\rho(x, y)) |g(x)| d\mu(x). \quad (7.5.14)$$

It follows from the definition of ψ that

$$\psi(x) \leq \max\{0, (2 - 4x)^{1/a}\}.$$

Now for all $x \in E(\mathfrak{p})$, it follows by the triangle inequality and (2.0.10) that

$$2 - 4D^{-s(\mathfrak{p})}\rho(x, y) \leq 2 - 4D^{-s(\mathfrak{p})}\rho(y, c(\mathfrak{p})) + 4D^{-s(\mathfrak{p})}\rho(x, c(\mathfrak{p})) \\ \leq 18 - 4D^{-s(\mathfrak{p})}\rho(y, c(\mathfrak{p})) \leq 4D^{-s(\mathfrak{p})}\rho(y, y') - 2.$$

Combining the above with the previous estimate on ψ , we get

$$\psi(D^{-s(\mathfrak{p})}\rho(x, y)) \leq 4(D^{-s(\mathfrak{p})}\rho(y, y'))^{1/a}.$$

Further, we obtain from the doubling property (1.1.2) and (2.0.10) that

$$\mu(B(x, D^{s(\mathfrak{p})})) \geq 2^{-3a} \mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})})).$$

Plugging this into (7.5.14) and using $a \geq 4$, we get

$$|T_{\mathfrak{p}}^*g(y)| \leq \frac{2^{103a^3}}{\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \left(\frac{\rho(y, y')}{D^{s(\mathfrak{p})}} \right)^{1/a} \int_{E(\mathfrak{p})} |g(x)| d\mu(y),$$

which completes the proof of the lemma. \square

Recall that

$$B(J) := B(c(J), 8D^{s(J)}).$$

We also denote

$$B'(J) := B(c(J), 16D^{s(J)}),$$

$$B^\circ(J) := B(c(J), \frac{1}{8}D^{s(J)}).$$

Lemma 7.5.6 (limited scale impact). *Let $\mathbf{p} \in \mathfrak{T}(u_2) \setminus \mathfrak{S}$, $J \in \mathcal{J}'$ and suppose that*

$$B(\mathcal{J}(\mathbf{p})) \cap B^\circ(J) \neq \emptyset.$$

Then

$$s(J) \leq s(\mathbf{p}) \leq s(J) + 3.$$

Proof. For the first estimate, assume that $s(\mathbf{p}) < s(J)$, then in particular $s(\mathbf{p}) \leq s(u_1)$. Since $\mathbf{p} \notin \mathfrak{S}$, we have by Lemma 7.4.7 that $\mathcal{J}(\mathbf{p}) \cap \mathcal{J}(u_1) = \emptyset$. Since $B(c(J), \frac{1}{4}D^{s(J)}) \subset \mathcal{J}(J) \subset \mathcal{J}(u_1)$, this implies

$$\rho(c(J), c(\mathbf{p})) \geq \frac{1}{4}D^{s(J)}.$$

On the other hand

$$\rho(c(J), c(\mathbf{p})) \leq \frac{1}{8}D^{s(J)} + 8D^{s(\mathbf{p})},$$

by our assumption. Thus $D^{s(\mathbf{p})} \geq 64^{-1}D^{s(J)}$, which contradicts (2.0.1) and $a \geq 4$.

For the second estimate, assume that $s(\mathbf{p}) > s(J) + 3$. Since $J \in \mathcal{J}'$, we have $J \subsetneq \mathcal{J}(u_1)$. Thus there exists $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$, by (2.0.7) and (2.0.8). By definition of \mathcal{J}' , there exists some $\mathbf{p}' \in \mathfrak{S}$ such that $\mathcal{J}(\mathbf{p}') \subset B(c(J'), 100D^{s(J)+2})$. On the other hand, since $B(\mathcal{J}(\mathbf{p})) \cap B^\circ(J) \neq \emptyset$, by the triangle inequality it holds that

$$B(c(J'), 100D^{s(J)+3}) \subset B(c(\mathbf{p}), 10D^{s(\mathbf{p})}).$$

Using the definition of \mathfrak{S} , we have

$$2^{Zn/2} \leq d_{\mathbf{p}'}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) \leq d_{B(c(J'), 100D^{s(J)+2})}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)).$$

By (1.1.7), this is

$$\begin{aligned} &\leq 2^{-100a} d_{B(c(J'), 100D^{s(J)+3})}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) \\ &\leq 2^{-100a} d_{B(c(\mathbf{p}), 10D^{s(\mathbf{p})})}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)), \end{aligned}$$

and by (1.1.5) and the definition of \mathfrak{S}

$$\leq 2^{-94a} d_{\mathbf{p}}(\mathcal{Q}(u_1), \mathcal{Q}(u_2)) \leq 2^{-94a} 2^{Zn/2}.$$

This is a contradiction, the second estimate follows. \square

Lemma 7.5.7 (local tree control). *For all $J \in \mathcal{J}'$ and all bounded g with bounded support*

$$\sup_{B^\circ(J)} |T_{\mathfrak{T}(u_2) \setminus \mathfrak{S}}^* g| \leq 2^{104a^3} \inf_J M_{\mathcal{B},1} |g|$$

Proof. By the triangle inequality and since $T_{\mathbf{p}}^*g = \mathbf{1}_{B(c(\mathbf{p}), 5D^{s(\mathbf{p})})}T_{\mathbf{p}}^*g$, we have

$$\sup_{B^\circ(J)} |T_{\mathfrak{T}(u_2) \setminus \mathfrak{S}}^*g| \leq \sup_{B^\circ(J)} \sum_{\substack{\mathbf{p} \in \mathfrak{T}(u_2) \setminus \mathfrak{S} \\ B(J(\mathbf{p})) \cap B^\circ(J) \neq \emptyset}} |T_{\mathbf{p}}^*g|.$$

By Lemma 7.5.6, this is at most

$$\sum_{s=s(J)}^{s(J)+3} \sum_{\substack{\mathbf{p} \in \mathfrak{P}, s(\mathbf{p})=s \\ B(J(\mathbf{p})) \cap B^\circ(J) \neq \emptyset}} \sup_{B^\circ(J)} |T_{\mathbf{p}}^*g|. \quad (7.5.15)$$

If $x \in E(\mathbf{p})$ and $B(J(\mathbf{p})) \cap B^\circ(J) \neq \emptyset$, then

$$B(c(J), 16D^{s(\mathbf{p})}) \subset B(x, 32D^{s(\mathbf{p})}),$$

by (2.0.10) and the triangle inequality. Using the doubling property (1.1.2), it follows that

$$\mu(B(x, D^{s(\mathbf{p})})) \geq 2^{-5a} \mu(B(c(J), 16D^{s(\mathbf{p})})).$$

Using (7.4.1), (2.1.3) and that $a \geq 4$, we bound (7.5.15) by

$$2^{103a^3} \sum_{s=s(J)}^{s(J)+3} \sum_{\substack{\mathbf{p} \in \mathfrak{P}, s(\mathbf{p})=s \\ B(J(\mathbf{p})) \cap B^\circ(J) \neq \emptyset}} \frac{1}{\mu(B(c(J), 16D^s))} \int_{E(\mathbf{p})} |g| d\mu.$$

For each $I \in \mathcal{D}$, the sets $E(\mathbf{p})$ for $\mathbf{p} \in \mathfrak{P}$ with $J(\mathbf{p}) = I$ are pairwise disjoint by (2.0.20) and (2.0.13). Further, if $B(J(\mathbf{p})) \cap B^\circ(J) \neq \emptyset$ and $s(\mathbf{p}) \geq s(J)$, then $E(\mathbf{p}) \subset B(c(J), 16D^{s(\mathbf{p})})$. Thus the last display is bounded by

$$\begin{aligned} & 2^{103a^3} \sum_{s=s(J)}^{s(J)+3} \frac{1}{\mu(B(c(J), 16D^s))} \int_{B(c(J), 16D^s)} |g| d\mu \\ & \leq \inf_{x' \in J} 2^{103a^3+2} M_{\mathcal{B},1} |g|. \end{aligned}$$

The lemma follows since $a \geq 4$. \square

Lemma 7.5.8 (scales impacting interval). *Let $\mathfrak{C} = \mathfrak{T}(u_1)$ or $\mathfrak{C} = \mathfrak{T}(u_2) \cap \mathfrak{S}$. Then for each $J \in \mathcal{J}'$ and $\mathbf{p} \in \mathfrak{C}$ with $B(J(\mathbf{p})) \cap B'(J) \neq \emptyset$, we have $s(\mathbf{p}) \geq s(J)$.*

Proof. By Lemma 7.4.7, we have that in both cases, $\mathfrak{C} \subset \mathfrak{S}$. If $\mathbf{p} \in \mathfrak{C}$ with $B(J(\mathbf{p})) \cap B'(J) \neq \emptyset$ and $s(\mathbf{p}) < s(J)$, then $J(\mathbf{p}) \subset B(c(J), 100D^{s(J)+1})$. Since $\mathbf{p} \in \mathfrak{S}$, it follows from the definition of \mathcal{J}' that $s(J) = -S$, which contradicts $s(\mathbf{p}) < s(J)$. \square

Lemma 7.5.9 (global tree control 1). *Let $\mathfrak{C}_1 = \mathfrak{T}(u_1)$ and $\mathfrak{C}_2 = \mathfrak{T}(u_2) \cap \mathfrak{S}$. Then for $i = 1, 2$ and each $J \in \mathcal{J}'$ and all bounded g with bounded support, we have*

$$\sup_{B'(J)} |T_{\mathfrak{C}_i}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{C}_i}^*g| + 2^{128a^3+4a+3} \inf_J M_{\mathcal{B},1} |g| \quad (7.5.16)$$

and for all $y, y' \in B'(J)$

$$\begin{aligned} & |e(Q(u_i)(y))T_{\mathfrak{C}_i}^*g(y) - e(Q(u_i)(y'))T_{\mathfrak{C}_i}^*g(y')| \\ & \leq 2^{128a^3+4a+1} \left(\frac{\rho(y, y')}{D^{s(J)}} \right)^{1/a} \inf_J M_{\mathcal{B},1} |g|. \end{aligned} \quad (7.5.17)$$

Proof. Note that (7.5.16) follows from (7.5.17), since for $y' \in B^\circ(J)$, by the triangle inequality,

$$\left(\frac{\rho(y, y')}{D^{s(J)}}\right)^{1/a} \leq \left(16 + \frac{1}{8}\right)^{1/a} \leq 2^2.$$

By the triangle inequality, Lemma 7.4.1 and Lemma 7.5.5, we have for all $y, y' \in B'(J)$

$$\begin{aligned} & |e(\mathcal{Q}(\mathbf{u}_i)(y))T_{\mathfrak{C}_i}^*g(y) - e(\mathcal{Q}(\mathbf{u}_i)(y'))T_{\mathfrak{C}_i}^*g(y')| \tag{7.5.18} \\ & \leq \sum_{\substack{\mathfrak{p} \in \mathfrak{C}_i \\ B(\mathcal{J}(\mathfrak{p})) \cap B'(J) \neq \emptyset}} |e(\mathcal{Q}(\mathbf{u}_i)(y))T_{\mathfrak{p}}^*g(y) - e(\mathcal{Q}(\mathbf{u}_i)(y'))T_{\mathfrak{p}}^*g(y')| \\ & \leq 2^{128a^3} \rho(y, y')^{1/a} \sum_{\substack{\mathfrak{p} \in \mathfrak{C}_i \\ B(\mathcal{J}(\mathfrak{p})) \cap B'(J) \neq \emptyset}} \frac{D^{-s(\mathfrak{p})/a}}{\mu(B(c(\mathfrak{p}), 4D^{s(\mathfrak{p})}))} \int_{E(\mathfrak{p})} |g| d\mu. \end{aligned}$$

By Lemma 7.5.8, we have $s(\mathfrak{p}) \geq s(J)$ for all \mathfrak{p} occurring in the sum. Further, for each $s \geq s(J)$, the sets $E(\mathfrak{p})$ for $\mathfrak{p} \in \mathfrak{P}$ with $s(\mathfrak{p}) = s$ are pairwise disjoint by (2.0.20) and (2.0.13), and contained in $B(c(J), 32D^s)$ by (2.0.10) and the triangle inequality. Using also the doubling estimate (1.1.2), we obtain that the expression in the last display can be estimated by

$$\begin{aligned} & 2^{128a^3} \rho(y, y')^{1/a} \sum_{S \geq s \geq s(J)} D^{-s/a} \frac{2^{4a}}{\mu(B(c(J), 32D^s))} \int_{B(c(J), 32D^s)} |g| d\mu \\ & \leq 2^{128a^3+4a} \left(\frac{\rho(y, y')}{D^{s(J)}}\right)^{1/a} \sum_{S \geq s \geq s(J)} D^{(s(J)-s)/a} \inf_J M_{\mathcal{B},1} |g|. \end{aligned}$$

Since $D^{-1/a} \leq \frac{1}{2}$, we have

$$\sum_{S \geq s \geq s(J)} D^{(s(J)-s)/a} \leq 2.$$

Estimate (7.5.17), and therefore the lemma, follow. \square

Lemma 7.5.10 (global tree control 2). *We have for all $J \in \mathcal{J}'$ and all bounded g with bounded support*

$$\sup_{B'(J)} |T_{\mathfrak{T}(u_2) \cap \mathfrak{S}}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{T}(u_2)}^*g| + 2^{129a^3} \inf_J M_{\mathcal{B},1} |g|.$$

Proof. By Lemma 7.5.9

$$\begin{aligned} & \sup_{B'(J)} |T_{\mathfrak{T}(u_2) \cap \mathfrak{S}}^*g| \leq \inf_{B^\circ(J)} |T_{\mathfrak{T}(u_2) \cap \mathfrak{S}}^*g| + 2^{128a^3+4a+3} \inf_J M_{\mathcal{B},1} |g| \\ & \leq \inf_{B^\circ(J)} |T_{\mathfrak{T}(u_2)}^*g| + \sup_{B^\circ(J)} |T_{\mathfrak{T}(u_2) \setminus \mathfrak{S}}^*g| + 2^{128a^3+4a+3} \inf_J M_{\mathcal{B},1} |g|, \end{aligned}$$

and by Lemma 7.5.7

$$\leq \inf_{B^\circ(J)} |T_{\mathfrak{T}(u_2)}^*g| + (2^{104a^3} + 2^{128a^3+4a+3}) \inf_J M_{\mathcal{B},1} |g|.$$

This completes the proof. \square

Proof of Lemma 7.5.4. Let P be the product on the right hand side of (7.5.8), and h_J be as defined in (7.5.7). By (7.5.3) and Lemma 7.4.1, the function h_J is supported in $B'(J) \cap \mathcal{J}(\mathbf{u}_1)$. By (7.5.3), Lemma 7.5.9 and Lemma 7.5.10, we have for all $y \in B'(J)$:

$$|h_J(y)| \leq 2^{257a^3+4a+3} P.$$

We have by the triangle inequality

$$|h_J(y) - h_J(y')| \leq |\chi_J(y) - \chi_J(y')| |T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y)| |T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y)| \quad (7.5.19)$$

$$+ |\chi_J(y')| |e(\mathcal{Q}(\mathbf{u}_1)(y)) T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y) - e(\mathcal{Q}(\mathbf{u}_1)(y')) T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y')| |T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y)| \quad (7.5.20)$$

$$+ |\chi_J(y')| |T_{\mathfrak{T}(\mathbf{u}_1)}^* g_1(y')| |e(\mathcal{Q}(\mathbf{u}_2)(y)) T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y) - e(\mathcal{Q}(\mathbf{u}_2)(y')) T_{\mathfrak{T}(\mathbf{u}_2) \cap \mathfrak{S}}^* g_2(y')|. \quad (7.5.21)$$

As h_J is supported in $\mathcal{J}(\mathbf{u}_1)$, we can assume without loss of generality that $y' \in \mathcal{J}(\mathbf{u}_1)$. If $y \notin \mathcal{J}(\mathbf{u}_1)$, then (7.5.19) vanishes. If $y \in \mathcal{J}(\mathbf{u}_1)$ then we have by (7.5.4), Lemma 7.5.9 and Lemma 7.5.10

$$(7.5.19) \leq 2^{484a^3+4a+3} \frac{\rho(y, y')}{D^{s(J)}} P,$$

where P denotes the product on the right hand side of (7.5.8).

By (7.5.3), Lemma 7.5.9 and Lemma 7.5.10, we have

$$(7.5.20) \leq 2^{257a^3+4a+1} \left(\frac{\rho(y, y')}{D^{s(J)}} \right)^{1/a} P.$$

By (7.5.3), and twice Lemma 7.5.9, we have

$$(7.5.21) \leq 2^{256a^3+8a+5} \left(\frac{\rho(y, y')}{D^{s(J)}} \right)^{1/a} P.$$

Using that $\rho(y, y') \leq 32D^{s(J)}$ and $a \geq 4$, the lemma follows. \square

7.5.3. The van der Corput estimate.

Lemma 7.5.11 (lower oscillation bound). *For all $J \in \mathcal{J}'$, we have that*

$$d_{B(J)}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \geq 2^{-201a^3} 2^{Zn/2}.$$

Proof. Since $\emptyset \neq \mathfrak{T}(\mathbf{u}_1) \subset \mathfrak{S}$ by Lemma 7.4.7, there exists at least one tile $\mathbf{p} \in \mathcal{S}$ with $J(\mathbf{p}) \subsetneq \mathcal{J}(\mathbf{u}_1)$. Thus $\mathcal{J}(\mathbf{u}_1) \notin \mathcal{J}'$, so $J \subsetneq \mathcal{J}(\mathbf{u}_1)$. Thus there exists a cube $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$, by (2.0.7) and (2.0.8). By definition of \mathcal{J}' and the triangle inequality, there exists $\mathbf{p} \in \mathfrak{S}$ such that

$$J(\mathbf{p}) \subset B(c(J'), 100D^{s(J')+1}) \subset B(c(J), 128D^{s(J)+2}).$$

Thus, by definition of \mathfrak{S} :

$$2^{Zn/2} \leq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \leq d_{B(c(J), 128D^{s(J)+2})}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

By the doubling property (1.1.5), this is

$$\leq 2^{200a^3+4a} d_{B(J)}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)),$$

which gives the lemma using $a \geq 4$. \square

Now we are ready to prove Lemma 7.4.5.

Proof of Lemma 7.4.5. We have

$$(7.4.6) = \left| \int_X T_{\mathfrak{I}(\mathbf{u}_1)}^* g_1 \overline{T_{\mathfrak{I}(\mathbf{u}_2) \cap \mathfrak{S}} g_2} \right|.$$

By Lemma 7.4.1, the right hand side is supported in $\mathcal{J}(\mathbf{u}_1)$. Using (7.5.2) of Lemma 7.5.2 and the definition (7.5.7) of h_J , we thus have

$$\leq \sum_{J \in \mathcal{J}'} \left| \int_X e(\mathcal{Q}(\mathbf{u}_2)(y) - \mathcal{Q}(\mathbf{u}_1)(y)) h_J(y) d\mu(y) \right|.$$

Using Proposition 2.0.5 with the ball $B(J)$, we bound this by

$$\leq 2^{7a} \sum_{J \in \mathcal{J}'} \mu(B(J)) \|h_J\|_{C^\tau(B'(J))} (1 + d_{B(J)}(\mathcal{Q}(\mathbf{u}_2), \mathcal{Q}(\mathbf{u}_1)))^{-1/(2a^2+a^3)}.$$

Using Lemma 7.5.4, Lemma 7.5.11 and $a \geq 4$, we bound the above by

$$\begin{aligned} &\leq 2^{485a^3+7a+3a^3+3a} 2^{-Zn/(4a^2+2a^3)} \sum_{J \in \mathcal{J}'} \mu(B(J)) \\ &\quad \times \prod_{j=1}^2 \left(\inf_{B^\circ(J)} |T_{\mathfrak{I}(\mathbf{u}_j)}^* g_j| + \inf_J M_{\mathcal{B},1} g_j \right). \end{aligned} \quad (7.5.22)$$

By the doubling property (1.1.2)

$$\mu(B(J)) \leq 2^{6a} \mu(B^\circ(J)),$$

thus

$$\begin{aligned} &\mu(B(J)) \prod_{j=1}^2 \left(\inf_{B^\circ(J)} |T_{\mathfrak{I}(\mathbf{u}_j)}^* g_j| + \inf_J M_{\mathcal{B},1} g_j \right) \\ &\leq 2^{6a} \int_{B^\circ(J)} \prod_{j=1}^2 (|T_{\mathfrak{I}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x) \\ &\leq 2^{6a} \int_J \prod_{j=1}^2 (|T_{\mathfrak{I}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x). \end{aligned}$$

Summing over $J \in \mathcal{J}'$, we obtain

$$(7.5.22) \leq 2^{499a^3} 2^{-Zn/(4a^2+2a^3)} \int_X \prod_{j=1}^2 (|T_{\mathfrak{I}(\mathbf{u}_j)}^* g_j|(x) + M_{\mathcal{B},1} g_j(x)) d\mu(x).$$

Applying the Cauchy-Schwarz inequality, Lemma 7.4.5 follows. \square

7.6. Proof of The Remaining Tiles Lemma. We define

$$\mathcal{J}' = \{J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}_1)) : J \subset \mathcal{J}(\mathbf{u}_1)\},$$

note that this is different from the \mathcal{J}' defined in the previous subsection.

Lemma 7.6.1 (dyadic partition 2). *We have*

$$\mathcal{J}(\mathbf{u}_1) = \bigcup_{J \in \mathcal{J}'} J.$$

Proof. By Lemma 7.1.2, it remains only to show that each $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}_1))$ with $J \cap \mathcal{J}(\mathbf{u}_1) \neq \emptyset$ is in \mathcal{J}' . But if $J \notin \mathcal{J}'$, then by (2.0.8) $\mathcal{J}(\mathbf{u}_1) \subsetneq J$. Pick $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_1)$. Then $\mathcal{J}(\mathbf{p}) \subsetneq J$. This contradicts the definition of $\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))$. \square

Lemma 7.4.6 follows from the following key estimate.

Lemma 7.6.2 (bound for tree projection). *We have for all bounded f with bounded support*

$$\|P_{\mathcal{J}'}|T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2\|_2 \leq 2^{102a^3 + 21a + 5} 2^{-\frac{25}{101a} Zn\kappa} \|\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} M_{\mathcal{B},1} |g_2\|_2$$

We prove this lemma below. First, we deduce Lemma 7.4.6.

Proof of Lemma 7.4.6. By Lemma 7.2.1 and Lemma 7.4.1, we have

$$(7.4.8) \leq 2^{130a^3} \|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}_1))} |\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} g_1\|_2 \|P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))} |\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2\|_2.$$

It follows from the definition of the projection operator P and Jensen's inequality that

$$\|P_{\mathcal{L}(\mathfrak{T}(\mathbf{u}_1))} |g_1 \mathbf{1}_{\mathcal{J}(\mathbf{u}_1)}\|_2 \leq \|g_1 \mathbf{1}_{\mathcal{J}(\mathbf{u}_1)}\|_2.$$

By Lemma 7.6.1, a cube $J \in \mathcal{J}(\mathfrak{T}(\mathbf{u}_1))$ intersects $\mathcal{J}(\mathbf{u}_1)$ if and only if $J \in \mathcal{J}'$. Thus

$$P_{\mathcal{J}(\mathfrak{T}(\mathbf{u}_1))} |\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2| = P_{\mathcal{J}'} |T_{\mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}}^* g_2|.$$

Combining this with Lemma 7.6.2, the definition (2.0.2) and $a \geq 4$ proves the lemma. \square

We need two more auxiliary lemmas before we prove Lemma 7.6.2.

Lemma 7.6.3 (thin scale impact). *If $\mathbf{p} \in \mathfrak{T}(\mathbf{u}_2) \setminus \mathfrak{S}$ and $J \in \mathcal{J}'$ with $B(\mathcal{J}(\mathbf{p})) \cap B(J) \neq \emptyset$, then*

$$s(\mathbf{p}) \leq s(J) + 2 - \frac{Zn}{202a^3}.$$

Proof. Suppose that $s(\mathbf{p}) > s(J) + 2 - \frac{Zn}{202a^3} =: s(J) - s_1$. Then, we have $s_1 + 2 \geq 0$ so

$$\rho(c(\mathbf{p}), c(J)) \leq 8D^{s(J)} + 8D^{s(\mathbf{p})} \leq 16D^{s(\mathbf{p}) + s_1 + 2}.$$

There exists a tile $\mathbf{q} \in \mathfrak{T}(\mathbf{u}_1)$. By (2.0.32), it satisfies $\mathcal{J}(\mathbf{q}) \subsetneq \mathcal{J}(\mathbf{u}_1)$. Thus $\mathcal{J}(\mathbf{u}_1) \notin \mathcal{J}'$. It follows that $J \subsetneq \mathcal{J}(\mathbf{u}_1)$. By (2.0.7) and (2.0.8), there exists a cube $J' \in \mathcal{D}$ with $J \subset J'$ and $s(J') = s(J) + 1$. By definition of \mathcal{J}' , there exists a tile $\mathbf{p}' \in \mathfrak{T}(\mathbf{u}_1)$ with

$$\mathcal{J}(\mathbf{p}') \subset B(c(J'), 100D^{s(J')+1}).$$

By the triangle inequality, the definition (2.0.1) and $a \geq 4$, we have

$$B(c(J'), 100D^{s(J')+1}) \subset B(c(\mathbf{p}), 128D^{s(\mathbf{p}) + s_1 + 2}).$$

Since $\mathbf{p}' \in \mathfrak{X}(\mathbf{u}_1)$ and $\mathcal{J}(\mathbf{u}_1) \subset \mathcal{J}(\mathbf{u}_2)$, we have by (2.0.36)

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{u}_2)) > 2^{Z(n+1)}.$$

Hence, by (2.0.32), the triangle inequality and using that by (2.0.3) $Z(n+1) = 2^{12a}(n+1) \geq 3$

$$d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) > 2^{Z(n+1)} - 4 \geq 2^{Z(n+1)-1}.$$

It follows that

$$2^{Z(n+1)-1} \leq d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)) \leq d_{B(c(\mathbf{p}), 128D^{s(\mathbf{p})+s_1+2})}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

Using (1.1.5), we obtain

$$\leq 2^{9a+100a^3(s_1+3)} d_{\mathbf{p}}(\mathcal{Q}(\mathbf{u}_1), \mathcal{Q}(\mathbf{u}_2)).$$

Since $\mathbf{p}' \notin \mathfrak{S}$ this is bounded by

$$\leq 2^{9a+100a^3(s_1+3)} 2^{Zn/2}.$$

Thus

$$Zn/2 + Z - 1 \leq 9a + 100a^3(s_1 + 3),$$

contradicting the definition of s_1 . \square

Lemma 7.6.4 (square function count). *For each $J \in \mathcal{J}'$ and all s , we have*

$$\frac{1}{\mu(J)} \int_J \left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{J}(\mathbf{u}_1) = \emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)} \right)^2 d\mu \leq 2^{14a+1} (8D^{-s})^\kappa.$$

Proof. Since $J \in \mathcal{J}'$ we have $J \subset \mathcal{J}(\mathbf{u}_1)$. Thus, if $B(I) \cap J \neq \emptyset$ then

$$B(I) \cap J \subset \{x \in J : \rho(x, X \setminus J) \leq 8D^{s(I)}\}. \quad (7.6.1)$$

Furthermore, for each s the balls $B(I)$ with $s(I) = s$ have bounded overlap: Consider the collection $\mathcal{D}_{s,x}$ of all $I \in \mathcal{D}$ with $x \in B(I)$ and $s(I) = s$. By (2.0.10) and (2.0.8), the balls $B(c(I), \frac{1}{4}D^{s(I)})$, $I \in \mathcal{D}_{s,x}$ are disjoint, and by the triangle inequality, they are contained in $\bar{B}(x, 9D^s)$. By the doubling property (1.1.2), we have

$$\mu(B(x, 9D^s)) \leq 2^{7a} \mu(B(c(I), \frac{1}{4}D^{s(I)}))$$

for each $I \in \mathcal{D}_{s,x}$. Thus

$$\mu(B(x, 9D^s)) \geq \sum_{I \in \mathcal{D}_{s,x}} \mu(B(c(I), \frac{1}{4}D^{s(I)})) \geq 2^{-7a} |\mathcal{D}_{s,x}| \mu(B(x, 9D^s)).$$

Dividing by the positive $\mu(B(x, 9D^s))$, we obtain that for each x

$$\left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{J}(\mathbf{u}_1) = \emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)}(x) \right)^2 \leq |\mathcal{D}_{s(J)-s,x}|^2 \leq 2^{14a}. \quad (7.6.2)$$

Combining (7.6.1), (7.6.2) and the small boundary property (2.0.11), noting that $8D^{s(I)} = 8D^{-s}D^{s(J)}$, the lemma follows. \square

Proof of Lemma 7.6.2. Expanding the definition of $P_{\mathcal{J}'}$, we have

$$\begin{aligned} & \|P_{\mathcal{J}'}|T_{\mathfrak{I}(u_2)\setminus\mathfrak{S}}^*g_2\|_2 \\ &= \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left(\int_J \left| \sum_{\mathfrak{p} \in \mathfrak{I}(u_2)\setminus\mathfrak{S}} T_{\mathfrak{p}}^*g_2(y) \right| d\mu(y) \right)^2 \right)^{1/2}. \end{aligned}$$

By Lemma 7.4.1, the innermost sum in the last display is 0 if $J \cap B(\mathcal{J}(\mathfrak{p})) = \emptyset$. Then we split that sum according to the scale of \mathfrak{p} relative to the scale of J . By Lemma 7.6.3, $s_1 \leq s(J) - s(\mathfrak{p}) \leq 2S$ with $s_1 := \lfloor \frac{Zn}{202a^3} - 2 \rfloor$:

$$= \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left(\int_J \left| \sum_{\substack{s=s_1 \\ \mathfrak{p} \in \mathfrak{I}(u_2)\setminus\mathfrak{S} \\ s(\mathfrak{p})=s(J)-s \\ J \cap B(\mathcal{J}(\mathfrak{p})) \neq \emptyset}}^{2S} T_{\mathfrak{p}}^*g_2(y) \right| d\mu(y) \right)^2 \right)^{1/2}.$$

Then we apply the triangle inequality and Minkowski's inequality to get

$$\leq \sum_{s=s_1}^{2S} \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left(\int_J \sum_{\substack{\mathfrak{p} \in \mathfrak{I}(u_2)\setminus\mathfrak{S} \\ s(\mathfrak{p})=s(J)-s \\ J \cap B(\mathcal{J}(\mathfrak{p})) \neq \emptyset}} |T_{\mathfrak{p}}^*g_2(y)| d\mu(y) \right)^2 \right)^{1/2}. \quad (7.6.3)$$

We have by Lemma 7.4.1 and (2.1.3)

$$|T_{\mathfrak{p}}^*g_2(y)| \leq 2^{102a^3} \mathbf{1}_{B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))}(y)} \int_{E(\mathfrak{p})} \frac{1}{\mu(B(x, D^{s(\mathfrak{p}))})} |g_2(x)| d\mu(x).$$

Using the doubling property (1.1.2), it follows that

$$\mu(B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))}) \leq 2^{4a} \mu(B(x, D^{s(\mathfrak{p}))}).$$

Thus, using also $a \geq 4$

$$|T_{\mathfrak{p}}^*g_2(y)| \leq 2^{102a^3+4a} \mathbf{1}_{B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))}(y)} \frac{1}{\mu(B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))})} \int_{E(\mathfrak{p})} |g_2(x)| d\mu(x).$$

Since for each $I \in \mathcal{D}$ the sets $E(\mathfrak{p})$, $\mathfrak{p} \in \mathfrak{P}(I)$ are disjoint, it follows that

$$\begin{aligned} & \int_J \sum_{\substack{\mathfrak{p} \in \mathfrak{I}(u_2)\setminus\mathfrak{S} \\ \mathcal{J}(\mathfrak{p})=I \\ J \cap B(\mathcal{J}(\mathfrak{p})) \neq \emptyset}} |T_{\mathfrak{p}}^*g_2(y)| d\mu \\ & \leq 2^{102a^3+4a} \int_J \mathbf{1}_{B(I)} \frac{1}{\mu(B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))})} \int_{B(c(\mathfrak{p}), 8D^{s(\mathfrak{p}))}} |g_2(x)| d\mu(x) \\ & \leq 2^{102a^3+4a} \int_J M_{\mathcal{B},1} |g_2|(y) \mathbf{1}_{B(I)}(y) d\mu(y). \end{aligned}$$

By Lemma 7.4.7, we have $\mathcal{J}(\mathbf{p}) \cap \mathcal{J}(\mathbf{u}_1) = \emptyset$ for all $\mathbf{p} \in \mathfrak{I}(\mathbf{u}_2) \setminus \mathfrak{S}$. Thus we can estimate (7.6.3) by

$$2^{102a^3+4a} \sum_{s=s_1}^{2S} \left(\sum_{J \in \mathcal{J}'} \frac{1}{\mu(J)} \left(\int_J \sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{J}(\mathbf{u}_1) = \emptyset \\ J \cap B(I) \neq \emptyset}} M_{\mathcal{B},1} |g_2| \mathbf{1}_{B(I)} d\mu \right)^2 \right)^{1/2},$$

which is by Cauchy-Schwarz at most

$$2^{102a^3+4a} \sum_{s=s_1}^{2S} \left(\sum_{J \in \mathcal{J}'} \int_J (M_{\mathcal{B},1} |g_2|)^2 d\mu \times \frac{1}{\mu(J)} \int_J \left(\sum_{\substack{I \in \mathcal{D}, s(I)=s(J)-s \\ I \cap \mathcal{J}(\mathbf{u}_1) = \emptyset \\ J \cap B(I) \neq \emptyset}} \mathbf{1}_{B(I)} \right) d\mu \right)^{1/2}. \quad (7.6.4)$$

Using Lemma 7.6.4, we bound (7.6.4) by

$$2^{102a^3+4a} \sum_{s=s_1}^{2S} \left(2^{14a+1} (8D^{-s})^\kappa \sum_{J \in \mathcal{J}'} \int_J (M_{\mathcal{B},1} |g_2|)^2 \right)^{1/2},$$

and since dyadic cubes in \mathcal{J}' form a partition of $\mathcal{J}(\mathbf{u}_1)$ by Lemma 7.6.1, $\kappa \leq 1$ by (2.0.2), and $a \geq 4$

$$\begin{aligned} &\leq 2^{102a^3+11a+2} \sum_{s=s_1}^{2S} D^{-s\kappa/2} \|\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} M_{\mathcal{B},1} |g_2|\|_2 \\ &\leq 2^{102a^3+11a+2} D^{-s_1\kappa/2} \frac{1}{1-D^{-\kappa/2}} \|\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} M_{\mathcal{B},1} |g_2|\|_2. \end{aligned}$$

By convexity of $t \mapsto D^{-t}$ and since $D \geq 2$, we have for all $0 \leq t \leq 1$

$$D^{-t} \leq 1 - t(1 - 1/D) \leq 1 - \frac{1}{2}t.$$

Using this for $t = \kappa/2$ and using that $s_1 = \frac{Zn}{202a^3} - 2$ and the definitions (2.0.1) and (2.0.2) of κ and D

$$\begin{aligned} &\leq 2^{102a^3+11a+2} 2^{-100a^2 \left(\frac{Zn}{202a^3} - 3 \right) \frac{\kappa}{2}} \frac{2}{\kappa} \|\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} M_{\mathcal{B},1} |g_2|\|_2 \\ &= 2^{102a^3+21a+4} 2^{150a^2\kappa} 2^{-\frac{100}{404a} Zn\kappa} \|\mathbf{1}_{\mathcal{J}(\mathbf{u}_1)} M_{\mathcal{B},1} |g_2|\|_2. \end{aligned}$$

Using the definition (2.0.2) of κ and $a \geq 4$, the lemma follows. \square

7.7. Forests. In this subsection, we complete the proof of Proposition 2.0.4 from the results of the previous subsections.

Define an n -row to be an n -forest $(\mathfrak{U}, \mathfrak{T})$, i.e. satisfying conditions (2.0.32) - (2.0.37), such that in addition the sets $\mathcal{J}(\mathbf{u}), \mathbf{u} \in \mathfrak{U}$ are pairwise disjoint.

Lemma 7.7.1 (forest row decomposition). *Let $(\mathfrak{U}, \mathfrak{T})$ be an n -forest. Then there exists a decomposition*

$$\mathfrak{U} = \bigcup_{1 \leq j \leq 2^n} \mathfrak{U}_j$$

such that for all $j = 1, \dots, 2^n$ the pair $(\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$ is an n -row.

Proof. Define recursively \mathfrak{U}_j to be a maximal disjoint set of tiles \mathbf{u} in

$$\mathfrak{U} \setminus \bigcup_{j' < j} \mathfrak{U}_{j'}$$

with inclusion maximal $\mathcal{J}(\mathbf{u})$. Properties (2.0.32), -(2.0.37) for $(\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$ follow immediately from the corresponding properties for $(\mathfrak{U}, \mathfrak{T})$, and the cubes $\mathcal{J}(\mathbf{u}), \mathbf{u} \in \mathfrak{U}_j$ are disjoint by definition. The collections \mathfrak{U}_j are also disjoint by definition.

Now we show by induction on j that each point is contained in at most $2^n - j$ cubes $\mathcal{J}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j} \mathfrak{U}_{j'}$. This implies that $\bigcup_{j=1}^{2^n} \mathfrak{U}_j = \mathfrak{U}$, which completes the proof of the Lemma. For $j = 0$ each point is contained in at most 2^n cubes by (2.0.34). For larger j , if x is contained in any cube $\mathcal{J}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' < j} \mathfrak{U}_{j'}$, then it is contained in a maximal such cube. Thus it is contained in a cube in $\mathcal{J}(\mathbf{u})$ with $\mathbf{u} \in \mathfrak{U}_j$. Thus the number $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j} \mathfrak{U}_{j'}$ with $x \in \mathcal{J}(\mathbf{u})$ is zero, or is less than the number of $\mathbf{u} \in \mathfrak{U} \setminus \bigcup_{j' \leq j-1} \mathfrak{U}_{j'}$ with $x \in \mathcal{J}(\mathbf{u})$ by at least one. \square

We pick a decomposition of the forest $(\mathfrak{U}, \mathfrak{T})$ into 2^n n -rows

$$(\mathfrak{U}_j, \mathfrak{T}_j) := (\mathfrak{U}_j, \mathfrak{T}|_{\mathfrak{U}_j})$$

as in Lemma 7.7.1. To save some space in the proofs of the remaining lemmas in this section we will write

$$\begin{aligned} T_{\mathfrak{C}} &= \sum_{\mathbf{p} \in \mathfrak{C}} T_{\mathbf{p}}, & T_{\mathfrak{C}}^* &= \sum_{\mathbf{p} \in \mathfrak{C}} T_{\mathbf{p}}^*, \\ T_{\mathfrak{R}_j} &= \sum_{\mathbf{u} \in \mathfrak{U}_j} T_{\mathfrak{T}(\mathbf{u})}, & T_{\mathfrak{R}_j}^* &= \sum_{\mathbf{u} \in \mathfrak{U}_j} T_{\mathfrak{T}(\mathbf{u})}^*. \end{aligned}$$

Lemma 7.7.2 (row bound). *For each $1 \leq j \leq 2^n$ and each bounded g supported on G we have*

$$\|T_{\mathfrak{R}_j}^* g\|_2 \leq 2^{182a^3} 2^{-n/2} \|g\|_2 \quad (7.7.1)$$

and

$$\|\mathbf{1}_F T_{\mathfrak{R}_j}^* g\|_2 \leq 2^{283a^3} 2^{-n/2} \text{dens}_2 \left(\bigcup_{\mathbf{u} \in \mathfrak{U}} \mathfrak{T}(\mathbf{u}) \right)^{1/2} \|g\|_2. \quad (7.7.2)$$

Proof. Since for each j the top cubes $\mathcal{J}(\mathbf{u})$, $\mathbf{u} \in \mathfrak{U}_j$ are disjoint, we have for all bounded g supported on G by Lemma 7.4.1

$$\begin{aligned} & \left\| \mathbf{1}_F \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}}^* g \right\|_2^2 = \left\| \mathbf{1}_F \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} \mathbf{1}_{\mathcal{J}(\mathbf{u})} T_{\mathfrak{p}}^* \mathbf{1}_{\mathcal{J}(\mathbf{u})} g \right\|_2^2 \\ &= \sum_{\mathbf{u} \in \mathfrak{U}_j} \int_{\mathcal{J}(\mathbf{u})} \left| \mathbf{1}_F \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}}^* \mathbf{1}_{\mathcal{J}(\mathbf{u})} g \right|^2 d\mu \leq \sum_{\mathbf{u} \in \mathfrak{U}_j} \left\| \mathbf{1}_F \sum_{\mathfrak{p} \in \mathfrak{T}(\mathbf{u})} T_{\mathfrak{p}}^* \mathbf{1}_G \mathbf{1}_{\mathcal{J}(\mathbf{u})} g \right\|_2^2. \end{aligned}$$

Applying Lemma 7.4.2 and the density assumption (2.0.35), then taking square roots, we obtain

$$\left\| \mathbf{1}_F T_{\mathfrak{R}_j}^* g \right\|_2 \leq 2^{282a^3} 2^{(4a+1-n)/2} \text{dens}_2 \left(\bigcup_{\mathbf{u} \in \mathfrak{U}} \mathfrak{T}(\mathbf{u}) \right)^{1/2} \left(\sum_{\mathbf{u} \in \mathfrak{U}_j} \left\| \mathbf{1}_{\mathcal{J}(\mathbf{u})} g \right\|_2^2 \right)^{1/2}.$$

Again by disjointedness of the cubes $\mathcal{J}(\mathbf{u})$, this is estimated by

$$2^{282a^3} 2^{(4a+1-n)/2} \text{dens}_2 \left(\bigcup_{\mathbf{u} \in \mathfrak{U}} \mathfrak{T}(\mathbf{u}) \right)^{1/2} \|g\|_2.$$

Thus (7.7.2) follows, since $a \geq 4$. The proof of (7.7.1) is the same up to replacing F by X . \square

Lemma 7.7.3 (row correlation). *For all $1 \leq j, j' \leq 2^n$ with $j \neq j'$ and for all bounded g_1, g_2 supported on G , it holds that*

$$\left| \int T_{\mathfrak{R}_j}^* g_1 \overline{T_{\mathfrak{R}_{j'}}^* g_2} d\mu \right| \leq 2^{876a^3 - 4n} \|g_1\|_2 \|g_2\|_2.$$

Proof. We have by Lemma 7.4.1 and the triangle inequality that

$$\begin{aligned} & \left| \int T_{\mathfrak{R}_j}^* g_1 \overline{T_{\mathfrak{R}_{j'}}^* g_2} d\mu \right| \\ & \leq \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \left| \int T_{\mathfrak{T}_j(\mathbf{u})}^* (\mathbf{1}_{\mathcal{J}(\mathbf{u})} g_1) \overline{T_{\mathfrak{T}_{j'}(\mathbf{u}')}^* (\mathbf{1}_{\mathcal{J}(\mathbf{u}')} g_2)} d\mu \right|. \end{aligned}$$

By Lemma 7.4.4, this is bounded by

$$2^{512a^3 - 4n} \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \|S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})} g_1)\|_{L^2(\mathcal{J}(\mathbf{u}') \cap \mathcal{J}(\mathbf{u}))} \|S_{2,\mathbf{u}'}(\mathbf{1}_{\mathcal{J}(\mathbf{u}')} g_2)\|_{L^2(\mathcal{J}(\mathbf{u}') \cap \mathcal{J}(\mathbf{u}))}. \quad (7.7.3)$$

We apply the Cauchy-Schwarz inequality in the form

$$\sum_{i \in M} a_i b_i \leq \left(\sum_{i \in M} a_i^2 \right)^{1/2} \left(\sum_{i \in M} b_i^2 \right)^{1/2}$$

to the outer two sums:

$$\leq 2^{512a^3 - 4n} \left(\sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \|S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})} g_1)\|_{L^2(\mathcal{J}(\mathbf{u}') \cap \mathcal{J}(\mathbf{u}))}^2 \right)^{1/2}$$

$$\left(\sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \|S_{2,\mathbf{u}'}(\mathbf{1}_{\mathcal{J}(\mathbf{u}')}g_2)\|_{L^2(\mathcal{J}(\mathbf{u}') \cap \mathcal{J}(\mathbf{u}))}^2 \right)^{1/2}.$$

We can now estimate the factor involving g_1 as follows:

$$\begin{aligned} & \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \|S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})}g_1)\|_{L^2(\mathcal{J}(\mathbf{u}') \cap \mathcal{J}(\mathbf{u}))}^2 \\ &= \sum_{\mathbf{u} \in \mathfrak{U}_j} \sum_{\mathbf{u}' \in \mathfrak{U}_{j'}} \int_{\mathcal{J}(\mathbf{u}) \cap \mathcal{J}(\mathbf{u}')} |S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})}(y)g_1(y))|^2 d\mu(y) \end{aligned}$$

By pairwise disjointedness of the sets $\mathcal{J}(\mathbf{u}')$ for $\mathbf{u}' \in \mathfrak{U}_{j'}$, we have

$$\begin{aligned} & \leq \sum_{\mathbf{u} \in \mathfrak{U}_j} \int_{\mathcal{J}(\mathbf{u})} |S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})}(y)g_1(y))|^2 d\mu(y) \leq \sum_{\mathbf{u} \in \mathfrak{U}_j} \int_X |S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})}(y)g_1(y))|^2 d\mu(y) \\ &= \sum_{\mathbf{u} \in \mathfrak{U}_j} \|S_{2,\mathbf{u}}(\mathbf{1}_{\mathcal{J}(\mathbf{u})}g_1)\|_2^2 \end{aligned}$$

By Lemma 7.4.3 we now estimate:

$$\leq \sum_{\mathbf{u} \in \mathfrak{U}_j} (2^{182a^3})^2 \|\mathbf{1}_{\mathcal{J}(\mathbf{u})}g_1\|_2^2$$

By pairwise disjointedness of the sets $\mathcal{J}(\mathbf{u})$ for $\mathbf{u} \in \mathfrak{U}_j$ (and writing out the definition of L^2 -norms), we have

$$\leq (2^{182a^3})^2 \|g_1\|_2^2$$

Arguing similarly for g_2 , we obtain the desired inequality. \square

Define for $1 \leq j \leq 2^n$

$$E_j := \bigcup_{\mathbf{u} \in \mathfrak{U}_j} \bigcup_{\mathbf{p} \in \mathfrak{T}(\mathbf{u})} E(\mathbf{p}).$$

Lemma 7.7.4 (disjoint row support). *The sets E_j , $1 \leq j \leq 2^n$ are pairwise disjoint.*

Proof. Suppose that $\mathbf{p} \in \mathfrak{T}(\mathbf{u})$ and $\mathbf{p}' \in \mathfrak{T}(\mathbf{u}')$ with $\mathbf{u} \neq \mathbf{u}'$ and $x \in E(\mathbf{p}) \cap E(\mathbf{p}')$. Suppose without loss of generality that $s(\mathbf{p}) \leq s(\mathbf{p}')$. Then $x \in \mathcal{J}(\mathbf{p}) \cap \mathcal{J}(\mathbf{p}') \subset \mathcal{J}(\mathbf{u}')$. By (2.0.8) it follows that $\mathcal{J}(\mathbf{p}) \subset \mathcal{J}(\mathbf{u}')$. By (2.0.36), it follows that

$$d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) > 2^{Z(n+1)}.$$

By the triangle inequality. Lemma 2.1.2 and (2.0.32) it follows that

$$\begin{aligned} d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{p}')) &\geq d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), \mathcal{Q}(\mathbf{u}')) - d_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{u}')) \\ &> 2^{Z(n+1)} - d_{\mathbf{p}'}(\mathcal{Q}(\mathbf{p}'), \mathcal{Q}(\mathbf{u}')) \\ &\geq 2^{Z(n+1)} - 4. \end{aligned}$$

Since $Z \geq 3$ by (2.0.3), it follows that $\mathcal{Q}(\mathbf{p}') \notin B_{\mathbf{p}}(\mathcal{Q}(\mathbf{p}), 1)$, so $\Omega(\mathbf{p}') \not\subset \Omega(\mathbf{p})$ by (2.0.15). Hence, by (2.0.14), $\Omega(\mathbf{p}) \cap \Omega(\mathbf{p}') = \emptyset$. But if $x \in E(\mathbf{p}) \cap E(\mathbf{p}')$ then $\mathcal{Q}(x) \in \Omega(\mathbf{p}) \cap \Omega(\mathbf{p}')$. This is a contradiction, and the lemma follows. \square

Now we prove Proposition 2.0.4.

Proof of Proposition 2.0.4. By (7.4.1), we have for each j

$$T_{\mathfrak{R}_j}^* g = \sum_{u \in \mathfrak{U}_j} \sum_{p \in \mathfrak{T}(u)} T_p^* g = \sum_{u \in \mathfrak{U}_j} \sum_{p \in \mathfrak{T}(u)} T_p^* \mathbf{1}_{E_j} g = T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g.$$

Hence, by Lemma 7.7.1 and the triangle inequality,

$$\begin{aligned} \left\| \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p^* g \right\|_2^2 &= \left\| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* g \right\|_2^2 = \left\| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g \right\|_2^2 \\ &= \int_X \left| \sum_{j=1}^{2^n} T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g \right|^2 d\mu \\ &\leq \sum_{j=1}^{2^n} \|T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g\|_2^2 + \sum_{j=1}^{2^n} \sum_{j' \neq j}^{2^n} \left| \int_X \overline{T_{\mathfrak{R}_j}^* \mathbf{1}_{E_j} g} T_{\mathfrak{R}_{j'}}^* \mathbf{1}_{E_{j'}} g \right| d\mu. \end{aligned}$$

We use Lemma 7.7.2 to estimate each term in the first sum, and Lemma 7.7.3 to bound each term in the second sum:

$$\leq 2^{566a^3-n} \sum_{j=1}^{2^n} \|\mathbf{1}_{E_j} g\|_2^2 + 2^{876a^3-4n} \sum_{j=1}^{2^n} \sum_{j'=1}^{2^n} \|\mathbf{1}_{E_j} g\|_2 \|\mathbf{1}_{E_{j'}} g\|_2.$$

By Cauchy-Schwarz in the second two sums, this is at most

$$2^{876a^3} (2^{-n} + 2^n 2^{-4n}) \sum_{j=1}^n \|\mathbf{1}_{E_j} g\|_2^2,$$

and by disjointedness of the sets E_j , this is at most

$$2^{877a^3-n} \|g\|_2^2.$$

Taking square roots, it follows that for all g

$$\left\| \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p^* g \right\|_2 \leq 2^{439a^3 - \frac{n}{2}} \|g\|_2. \quad (7.7.4)$$

On the other hand, we have by disjointedness of the sets E_j from Lemma 7.7.4 and the triangle inequality

$$\left\| \mathbf{1}_G \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p f \right\|_2^2 = \left\| \sum_{j=1}^{2^n} \mathbf{1}_{E_j} \mathbf{1}_G T_{\mathfrak{R}_j} f \right\|_2^2 \leq \sum_{j=1}^{2^n} \|\mathbf{1}_{E_j} \mathbf{1}_G T_{\mathfrak{R}_j} f\|_2^2 \leq \sum_{j=1}^{2^n} \|\mathbf{1}_G T_{\mathfrak{R}_j} f\|_2^2.$$

Now with $|f| \leq \mathbf{1}_F$ and Lemma 7.7.2 we obtain

$$\begin{aligned} \|\mathbf{1}_G T_{\mathfrak{R}_j} f\|_2^2 &= \left| \int_X \overline{\mathbf{1}_G T_{\mathfrak{R}_j} f} T_{\mathfrak{R}_j} f \right| = \left| \int_X \overline{T_{\mathfrak{R}_j}^* \mathbf{1}_G T_{\mathfrak{R}_j} f} \mathbf{1}_F f \right| \\ &\leq \|f\|_2 \|\mathbf{1}_F T_{\mathfrak{R}_j}^* \mathbf{1}_G T_{\mathfrak{R}_j} f\|_2 \leq 2^{283a^3-n/2} \text{dens}_2 \left(\bigcup_{u \in \mathfrak{U}} \mathfrak{T}(u) \right)^{1/2} \|\mathbf{1}_G T_{\mathfrak{R}_j} f\|_2 \|f\|_2. \end{aligned}$$

Dividing this last inequality by the finite $\|\mathbf{1}_G T_{\mathfrak{A}_j} f\|_2$, substituting back and taking square roots we get

$$\begin{aligned} \left\| \mathbf{1}_G \sum_{u \in \mathfrak{U}} \sum_{p \in \mathfrak{T}(u)} T_p f \right\|_2 &\leq 2^{283a^3} \text{dens}_2 \left(\bigcup_{u \in \mathfrak{U}} \mathfrak{T}(u) \right)^{\frac{1}{2}} 2^{-\frac{n}{2}} \left(\sum_{j=1}^{2^n} \|f\|_2^2 \right)^{\frac{1}{2}} \\ &= 2^{283a^3} \text{dens}_2 \left(\bigcup_{u \in \mathfrak{U}} \mathfrak{T}(u) \right)^{\frac{1}{2}} \|f\|_2. \end{aligned} \quad (7.7.5)$$

Proposition 2.0.4 follows by taking the product of the $(2 - \frac{2}{q})$ -th power of (7.7.4) and the $(\frac{2}{q} - 1)$ -st power of (7.7.5). \square

8. PROOF OF THE HÖLDER CANCELLATIVE CONDITION

We need the following auxiliary lemma. Recall that $\tau = 1/a$.

Lemma 8.0.1 (Lipschitz Holder approximation). *Let $z \in X$ and $R > 0$. Let $\varphi : X \rightarrow \mathbb{C}$ be a function supported in the ball $B := B(z, R)$ with finite norm $\|\varphi\|_{C^\tau(B(z, 2R))}$. Let $0 < t \leq 1$. There exists a function $\tilde{\varphi} : X \rightarrow \mathbb{C}$, supported in $B(z, 2R)$, such that for every $x \in X$*

$$|\varphi(x) - \tilde{\varphi}(x)| \leq (t/2)^\tau \|\varphi\|_{C^\tau(B(z, 2R))} \quad (8.0.1)$$

and

$$\|\tilde{\varphi}\|_{\text{Lip}(B(z, 2R))} \leq 2^{4a} t^{-1-a} \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.2)$$

Proof. Define for $x, y \in X$ the Lipschitz and thus measurable function

$$L(x, y) := \max\{0, 1 - \frac{\rho(x, y)}{tR}\}. \quad (8.0.3)$$

We have that $L(x, y) \neq 0$ implies

$$y \in B(x, tR). \quad (8.0.4)$$

We have for $y \in B(x, 2^{-1}tR)$ that

$$|L(x, y)| \geq 2^{-1}. \quad (8.0.5)$$

Hence

$$\int L(x, y) d\mu(y) \geq 2^{-1} \mu(B(x, 2^{-1}tR)). \quad (8.0.6)$$

Let n be the smallest integer so that

$$2^n t \geq 1. \quad (8.0.7)$$

Iterating $n + 2$ times the doubling condition (1.1.2), we obtain

$$\int L(x, y) d\mu(y) \geq 2^{-1-a(n+2)} \mu(B(x, 2R)). \quad (8.0.8)$$

Now define

$$\tilde{\varphi}(x) := \left(\int L(x, y) d\mu(y) \right)^{-1} \int L(x, y) \varphi(y) d\mu(y).$$

Using that φ is supported in $B(z, R)$ and (8.0.4), we have that $\tilde{\varphi}$ is supported in $B(z, 2R)$.

We prove (8.0.1). For any $x \in X$, using that L is nonnegative,

$$\left(\int L(x, y) \, d\mu(y) \right) |\varphi(x) - \tilde{\varphi}(x)| \quad (8.0.9)$$

$$= \left| \int L(x, y) (\varphi(x) - \varphi(y)) \, d\mu(y) \right|. \quad (8.0.10)$$

Using (8.0.4), we estimate the last display by

$$\leq \int_{B(x, tR)} L(x, y) |\varphi(x) - \varphi(y)| \, d\mu(y). \quad (8.0.11)$$

We claim that in this integral, $|\varphi(x) - \varphi(y)| \leq \rho(x, y)^\tau \|\varphi\|_{C^\tau(B(z, 2R))} (2R)^{-\tau}$. Indeed, if x or y does not belong to $B(z, 2R)$, then the other point is not in $B(z, R)$ as $\rho(x, y) \leq tR \leq R$. Therefore, $\varphi(x) = \varphi(y) = 0$ since φ is supported in $B(z, R)$. Otherwise, both points are in $B(z, 2R)$, and the inequality follows from the definition of $\|\varphi\|_{C^\tau(B(z, 2R))}$.

Therefore, we can estimate the last display further by

$$\leq \left(\int_{B(x, tR)} L(x, y) \rho(x, y)^\tau \, d\mu(y) \right) \|\varphi\|_{C^\tau(B(z, 2R))} (2R)^{-\tau}. \quad (8.0.12)$$

Using the condition on the domain of integration to estimate $\rho(x, y)$ by tR and then expanding the domain by positivity of the integrand, we estimate this further by

$$\leq \left(\int L(x, y) \, d\mu(y) \right) \|\varphi\|_{C^\tau(B(z, 2R))} (t/2)^\tau. \quad (8.0.13)$$

Dividing the string of inequalities from (8.0.9) to (8.0.13) by the positive integral of L proves (8.0.1).

We turn to (8.0.2). For every $x \in X$, we have

$$\left| \int L(x, y) \, d\mu(y) \right| |\tilde{\varphi}(x)| = \left| \int L(x, y) \varphi(y) \, d\mu(y) \right| \quad (8.0.14)$$

$$\leq \left| \int L(x, y) \, d\mu(y) \right| \sup_{x' \in X} |\varphi(x')|. \quad (8.0.15)$$

As φ is supported on B , dividing by the integral of L , we obtain

$$|\tilde{\varphi}(x)| \leq \sup_{x' \in B} |\varphi(x')| \leq \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.16)$$

If $\rho(x, x') \geq R$, then we have by the triangle inequality

$$R \frac{|\tilde{\varphi}(x') - \tilde{\varphi}(x)|}{\rho(x, x')} \leq 2 \sup_{x'' \in X} |\tilde{\varphi}(x'')| \leq 2 \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.17)$$

Now assume $\rho(x, x') < R$. For $y \in X$ we have by the triangle inequality and a two fold case distinction for the maximum in the definition of L ,

$$|L(x, y) - L(x', y)| \leq \frac{\rho(x, x')}{tR}. \quad (8.0.18)$$

We compute with (8.0.18), first adding and subtracting a term in the integral,

$$\left(\int L(x, y) \, d\mu(y) \right) |\tilde{\varphi}(x') - \tilde{\varphi}(x)| = \quad (8.0.19)$$

$$\left| \int L(x, y) \tilde{\varphi}(x') - L(x, y) \tilde{\varphi}(x) + L(x', y) \tilde{\varphi}(x') - L(x', y) \tilde{\varphi}(x) \, d\mu(y) \right|. \quad (8.0.20)$$

Grouping the second and third and the first and fourth term, we obtain using the definition of $\tilde{\varphi}$ and Fubini,

$$\leq \left| \int (L(x', y) - L(x, y)) \varphi(y) \, d\mu(y) \right| \quad (8.0.21)$$

$$+ \left| \int L(x, y) \, d\mu(y) - \int L(x', y) \, d\mu(y) \right| |\tilde{\varphi}(x')| \quad (8.0.22)$$

$$\leq 2 \int |L(x, y) - L(x', y)| \, d\mu(y) \|\varphi\|_{C^\tau(B(z, 2R))}, \quad (8.0.23)$$

where in the last inequality we have used (8.0.16). Using further (8.0.18) and the support of L , we estimate the last display by

$$\leq 2 \frac{\rho(x, x')}{tR} \mu(B(x, tR) \cup B(x', tR)) \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.24)$$

Using $\rho(x, x') < R$ and the triangle inequality, we estimate the last display by

$$\leq 2 \frac{\rho(x, x')}{tR} \mu(B(x, 2R)) \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.25)$$

Dividing by the integral over L and using (8.0.8) and (8.0.7), we obtain

$$\frac{R |\tilde{\varphi}(x') - \tilde{\varphi}(x)|}{\rho(x, x')} \leq 2^{2+a(n+2)} t^{-1} \|\varphi\|_{C^\tau(B(z, 2R))} \leq 2^{2+3a} t^{-1-a} \|\varphi\|_{C^\tau(B(z, 2R))}. \quad (8.0.26)$$

Combining (8.0.17) and (8.0.26) using $a \geq 4$ and $t \leq 1$ and adding (8.0.16) proves (8.0.2) and completes the proof of Lemma 8.0.1. \square

We turn to the proof of Proposition 2.0.5.

Proof of Proposition 2.0.5. Let $z \in X$ and $R > 0$ and set $B = B(z, R)$. Let φ be given as in Proposition 2.0.5. Set

$$t := (1 + d_B(\vartheta, \theta))^{-\frac{\tau}{2+a}} \quad (8.0.27)$$

and define $\tilde{\varphi}$ as in Lemma 8.0.1. Let ϑ and θ be in Θ . Then

$$\left| \int e(\vartheta(x) - \theta(x)) \varphi(x) \, d\mu(x) \right| \quad (8.0.28)$$

$$\leq \left| \int e(\vartheta(x) - \theta(x)) \tilde{\varphi}(x) \, d\mu(x) \right| \quad (8.0.29)$$

$$+ \left| \int e(\vartheta(x) - \theta(x)) (\varphi(x) - \tilde{\varphi}(x)) \, d\mu(x) \right| \quad (8.0.30)$$

Using the cancellative condition (1.1.9) of Θ on the ball $B(z, 2R)$, the term (8.0.29) is bounded above by

$$2^a \mu(B(z, 2R)) \|\tilde{\varphi}\|_{\text{Lip}(B(z, 2R))} (1 + d_{B(z, 2R)}(\vartheta, \theta))^{-\tau}. \quad (8.0.31)$$

Using the doubling condition (1.1.2), the inequality (8.0.2), and the estimate $d_B \leq d_{B(z, 2R)}$ from the definition, we estimate (8.0.31) from above by

$$2^{6a} t^{-1-a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\tau}. \quad (8.0.32)$$

The term (8.0.30) we estimate using (8.0.1) and that ϑ and θ are real and thus $e(\vartheta)$ and $e(\theta)$ bounded in absolute value by 1. We obtain for (8.0.30) with (1.1.2) the upper bound

$$\mu(B(z, 2R))(t/2)^\tau \|\varphi\|_{C^\tau(B)} \leq 2^a \mu(B) t^\tau \|\varphi\|_{C^\tau(B)}. \quad (8.0.33)$$

Using the definition (8.0.27) of t and adding (8.0.32) and (8.0.33) estimates (8.0.28) from above by

$$2^{6a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau}{2+a}} \quad (8.0.34)$$

$$+ 2^a \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau^2}{2+a}}. \quad (8.0.35)$$

$$\leq 2^{1+6a} \mu(B) \|\varphi\|_{C^\tau(B)} (1 + d_B(\vartheta, \theta))^{-\frac{\tau^2}{2+a}}, \quad (8.0.36)$$

where we used $\tau \leq 1$. This completes the proof of Proposition 2.0.5. \square

9. PROOF OF VITALI COVERING AND HARDY–LITTLEWOOD

We begin with a classical representation of the Lebesgue norm.

Lemma 9.0.1 (layer cake representation). *Let $1 \leq p < \infty$. Then for any measurable function $u : X \rightarrow [0, \infty)$ on the measure space X relative to the measure μ we have*

$$\|u\|_p^p = p \int_0^\infty \lambda^{p-1} \mu(\{x : u(x) \geq \lambda\}) d\lambda. \quad (9.0.1)$$

Proof. The left-hand side of (9.0.1) is by definition

$$\int_X u(x)^p d\mu(x). \quad (9.0.2)$$

Writing $u(x)$ as an elementary integral in λ and then using Fubini, we write for the last display

$$= \int_X \int_0^{u(x)} p \lambda^{p-1} d\lambda d\mu(x) \quad (9.0.3)$$

$$= p \int_0^\infty \lambda^{p-1} \mu(\{x : u(x) \geq \lambda\}) d\lambda. \quad (9.0.4)$$

This proves the lemma. \square

The following lemma will be used to define M in the proof of Proposition 2.0.6.

Lemma 9.0.2 (covering separable space). *For each $r > 0$, there exists a countable collection $C(r) \subset X$ of points such that*

$$X \subset \bigcup_{c \in C(r)} B(c, r).$$

Proof. It clearly suffices to construct finite collections $C(r, k)$ such that

$$B(o, r2^k) \subset \bigcup_{c \in C(r, k)} B(c, r),$$

since then the collection $C(r) = \bigcup_{k \in \mathbb{N}} C(r, k)$ has the desired property.

Suppose that $Y \subset B(o, r2^k)$ is a collection of points such that for all $y, y' \in Y$ with $y \neq y'$, we have $\rho(y, y') \geq r$. Then the balls $B(y, r/2)$ are pairwise disjoint and contained in $B(o, r2^{k+1})$. If $y \in B(o, r)$, then $B(o, r2^{k+1}) \subset B(y, r2^{k+2})$. Thus, by the doubling property (1.1.2),

$$\mu(B(y, \frac{r}{2})) \geq 2^{-(k+2)a} \mu(B(o, r2^{k+1})).$$

Thus, we have

$$\mu(B(o, r2^{k+1})) \geq \sum_{y \in Y} \mu(B(y, \frac{r}{2})) \geq |Y| 2^{-(k+2)a} \mu(B(o, r2^{k+1})).$$

We conclude that $|Y| \leq 2^{(k+2)a}$. In particular, there exists a set Y of maximal cardinality. Define $C(r, k)$ to be such a set.

If $x \in B(o, r2^k)$ and $x \notin C(r, k)$, then there must exist $y \in C(r, k)$ with $\rho(x, y) < r$. Thus $C(r, k)$ has the desired property. \square

We turn to the proof of Proposition 2.0.6.

Proof of Proposition 2.0.6. Let the collection \mathcal{B} be given. We first show (2.0.43).

We recursively choose a finite sequence $B_i \in \mathcal{B}$ for $i \geq 0$ as follows. Assume $B_{i'}$ is already chosen for $0 \leq i' < i$. If there exists a ball $B_i \in \mathcal{B}$ so that B_i is disjoint from all $B_{i'}$ with $0 \leq i' < i$, then choose such a ball $B_i = B(x_i, r_i)$ with maximal r_i .

If there is no such ball, stop the selection and set $i'' := i$.

By disjointedness of the chosen balls and since $0 \leq u$, we have

$$\sum_{0 \leq i < i''} \int_{B_i} u(x) d\mu(x) \leq \int_X u(x) d\mu(x). \quad (9.0.5)$$

By (2.0.42), we conclude

$$\lambda \sum_{0 \leq i < i''} \mu(B_i) \leq \int_X u(x) d\mu(x). \quad (9.0.6)$$

Let $x \in \bigcup \mathcal{B}$. Choose a ball $B' = B(x', r') \in \mathcal{B}$ such that $x \in B'$. If B' is one of the selected balls, then

$$x \in \bigcup_{0 \leq i < i''} B_i \subset \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.0.7)$$

If B' is not one of the selected balls, then as it is not selected at time i'' , there is a selected ball B_i with $B' \cap B_i \neq \emptyset$. Choose such B_i with minimal index i . As B' is therefore disjoint from all balls $B_{i'}$ with $i' < i$ and as it was not selected in place of B_i , we have $r_i \geq r'$.

Using a point y in the intersection of B_i and B' , we conclude by the triangle inequality

$$\rho(x_i, x') \leq \rho(x_i, y) + \rho(x', y) \leq r_i + r' \leq 2r_i. \quad (9.0.8)$$

By the triangle inequality again, we further conclude

$$\rho(x_i, x) \leq \rho(x_i, x') + \rho(x', x) \leq 2r_i + r' \leq 3r_i. \quad (9.0.9)$$

It follows that

$$x \in \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.0.10)$$

With (9.0.7) and (9.0.10), we conclude

$$\bigcup \mathcal{B} \subset \bigcup_{0 \leq i < i''} B(x_i, 3r_i). \quad (9.0.11)$$

With the doubling property (1.1.2) applied twice, we conclude

$$\mu(\bigcup \mathcal{B}) \leq \sum_{0 \leq i < i''} \mu(B(x_i, 3r_i)) \leq 2^{2a} \sum_{0 \leq i < i''} \mu(B_i). \quad (9.0.12)$$

With (9.0.6) and (9.0.12) we conclude (2.0.43).

We turn to the proof of (2.0.44). We first consider the case $p_1 = 1$ and recall $M_{\mathcal{B}} = M_{\mathcal{B},1}$. We write for the p_2 -th power of left-hand side of (2.0.44) with Lemma 9.0.1 and a change of variables

$$\|M_{\mathcal{B}}u(x)\|_{p_2}^{p_2} = p_2 \int_0^\infty \lambda^{p_2-1} \mu(\{x : M_{\mathcal{B}}u(x) \geq \lambda\}) d\lambda \quad (9.0.13)$$

$$= 2^{p_2} p_2 \int_0^\infty \lambda^{p_2-1} \mu(\{x : M_{\mathcal{B}}u(x) \geq 2\lambda\}) d\lambda. \quad (9.0.14)$$

Fix $\lambda \geq 0$ and let $x \in X$ satisfy $M_{\mathcal{B}}u(x) \geq 2\lambda$. By definition of $M_{\mathcal{B}}$, there is a ball $B' \in \mathcal{B}$ such that $x \in B'$ and

$$\int_{B'} u(y) d\mu(y) \geq 2\lambda \mu(B'). \quad (9.0.15)$$

Define $u_\lambda(y) := 0$ if $|u(y)| < \lambda$ and $u_\lambda(y) := u(y)$ if $|u(y)| \geq \lambda$. Then with (9.0.15)

$$\int_{B'} u_\lambda(y) d\mu(y) = \int_{B'} u(y) d\mu(y) - \int_{B'} (u - u_\lambda)(y) d\mu(y) \quad (9.0.16)$$

$$\geq 2\lambda \mu(B') - \int_{B'} (u - u_\lambda)(y) d\mu(y). \quad (9.0.17)$$

As $(u - u_\lambda)(y) \leq \lambda$ by definition, we can estimate the last display by

$$\geq 2\lambda \mu(B') - \int_{B'} \lambda d\mu(y) = \lambda \mu(B'). \quad (9.0.18)$$

Hence x is contained in $\bigcup (\mathcal{B}_\lambda)$, where \mathcal{B}_λ is the collection of balls B'' in \mathcal{B} such that

$$\int_{B''} u_\lambda(y) d\mu(y) \geq \lambda \mu(B''). \quad (9.0.19)$$

We have thus seen

$$\{x : M_{\mathcal{B}}u(x) \geq 2\lambda\} \subset \bigcup \mathcal{B}_\lambda. \quad (9.0.20)$$

Applying (2.0.43) to the collection \mathcal{B}_λ gives

$$\lambda \mu(\{x : M_{\mathcal{B}}u(x) \geq 2\lambda\}) \leq 2^{2a} \int u_\lambda(x) dx. \quad (9.0.21)$$

With Lemma 9.0.1,

$$\lambda \mu(\{x : M_{\mathcal{B}}u(x) \geq 2\lambda\}) \leq 2^{2a} \int_0^\infty \mu(\{x : |u_\lambda(x)| \geq \lambda'\}) d\lambda'. \quad (9.0.22)$$

By definition of u_λ , making a case distinction between $\lambda \geq \lambda'$ and $\lambda < \lambda'$, we see that

$$\mu(\{x : |u_\lambda(x)| \geq \lambda'\}) \leq \mu(\{x : |u(x)| \geq \max(\lambda, \lambda')\}). \quad (9.0.23)$$

We obtain with (9.0.14), (9.0.22), and (9.0.23)

$$\|M_{\mathcal{B}}u(x)\|_{p_2}^{p_2} \quad (9.0.24)$$

$$\leq 2^{p_2+2a} p_2 \int_0^\infty \lambda^{p_2-2} \int_0^\infty \mu(\{x : |u(x)| \geq \max(\lambda, \lambda')\}) d\lambda' d\lambda. \quad (9.0.25)$$

We split the integral into $\lambda \geq \lambda'$ and $\lambda < \lambda'$ and resolve the maximum correspondingly. We have for $\lambda \geq \lambda'$ with Lemma 9.0.1

$$\int_0^\infty \lambda^{p_2-2} \int_0^\lambda \mu(\{x : |u(x)| \geq \lambda\}) d\lambda' d\lambda \quad (9.0.26)$$

$$= \int_0^\infty \lambda^{p_2-1} \mu(\{x : |u(x)| \geq \lambda\}) d\lambda. \quad (9.0.27)$$

$$= p_2^{-1} \|u\|_{p_2}^{p_2}. \quad (9.0.28)$$

We have for $\lambda < \lambda'$ with Fubini and Lemma 9.0.1

$$\int_0^\infty \lambda^{p_2-2} \int_\lambda^\infty \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda' d\lambda. \quad (9.0.29)$$

$$= \int_0^\infty \int_0^{\lambda'} \lambda^{p_2-2} \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda d\lambda'. \quad (9.0.30)$$

$$= (p_2 - 1)^{-1} \int_0^\infty (\lambda')^{p_2-1} \mu(\{x : |u(x)| \geq \lambda'\}) d\lambda'. \quad (9.0.31)$$

$$= (p_2 - 1)^{-1} p_2^{-1} \|u\|_{p_2}^{p_2}. \quad (9.0.32)$$

Adding the two estimates (9.0.28) and (9.0.32) gives

$$\|M_{\mathcal{B}}u(x)\|_{p_2}^{p_2} \leq 2^{p_2+2a} (1 + (p_2 - 1)^{-1}) \|u\|_{p_2}^{p_2} = 2^{p_2+2a} p_2 (p_2 - 1)^{-1} \|u\|_{p_2}^{p_2}. \quad (9.0.33)$$

With $a \geq 1$ and $p_2 > 1$, taking the p_2 -th root, we obtain (2.0.44). We turn to the case of general $1 \leq p_1 < p_2$. We have

$$M_{\mathcal{B}, p_1} u = (M_{\mathcal{B}}(|u|^{p_1}))^{\frac{1}{p_1}}. \quad (9.0.34)$$

Applying the special case of (2.0.44) for $M_{\mathcal{B}}$ gives

$$\|M_{\mathcal{B}, p_1} u\|_{p_2} = \|M_{\mathcal{B}}(|u|^{p_1})\|_{p_2/p_1}^{\frac{1}{p_1}} \quad (9.0.35)$$

$$\leq 2^{2a} (p_2/p_1) (p_2/p_1 - 1)^{-1} \|(|u|^{p_1})\|_{p_2/p_1}^{\frac{1}{p_1}} = 2^{2a} p_2 (p_2 - p_1)^{-1} \|u\|_{p_2}. \quad (9.0.36)$$

This proves (2.0.44) in general.

Now we construct the operator M satisfying (2.0.45) and (2.0.46). For each $k \in \mathbb{Z}$ we choose a countable set $C(2^k)$ as in Lemma 9.0.2. Define

$$\mathcal{B}_\infty = \{B(c, 2^k) : c \in C(2^k), k \in \mathbb{Z}\}.$$

By Lemma 9.0.2, this is a countable collection of balls. We choose an enumeration $\mathcal{B}_\infty = \{B_1, \dots\}$ and define

$$\mathcal{B}_n = \{B_1, \dots, B_n\}.$$

We define

$$Mw := 2^{2a} \sup_{n \in \mathbb{N}} M_{\mathcal{B}_n} w.$$

This function is measurable for each measurable w , since it is a countable supremum of measurable functions. Estimate (2.0.46) follows immediately from (2.0.44) and the monotone convergence theorem.

It remains to show (2.0.45). Let $B = B(x, r) \subset X$. Let k be the smallest integer such that $2^k \geq r$, in particular we have $2^k < 2r$. By definition of $C(2^k)$, there exists $c \in C(2^k)$ with $x \in B(c, 2^k)$. By the triangle inequality, we have $B(c, 2^k) \subset B(x, 4r)$, and hence by the doubling property (1.1.2)

$$\mu(B(c, 2^k)) \leq 2^{2a} \mu(B(x, r)).$$

It follows that for each $z \in B(x, r)$

$$\begin{aligned} \frac{1}{\mu(B(x, r))} \int_{B(x, r)} |w(y)| d\mu(y) &\leq \frac{2^{2a}}{\mu(B(c, 2^k))} \int_{B(c, 2^k)} |w(y)| d\mu(y) \\ &\leq Mw(z). \end{aligned}$$

This completes the proof. \square

10. TWO-SIDED METRIC SPACE CARLESON

We prove a variant of Theorem 1.1.1 for a two-sided Calderón–Zygmund kernel on the doubling metric measure space (X, ρ, μ, a) , i.e. a one-sided Calderón–Zygmund kernel K which additionally satisfies for all $x, x', y \in X$ with $x \neq y$ and $2\rho(x, x') \leq \rho(x, y)$,

$$|K(x, y) - K(x', y)| \leq \left(\frac{\rho(x, x')}{\rho(x, y)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x, y)}. \quad (10.0.1)$$

By the additional regularity, we can weaken the assumption (1.1.15) to a family of operators that is easier to work with in applications. Namely, for $r > 0$, $x \in X$, and a bounded, measurable function $f : X \rightarrow \mathbb{C}$ supported on a set of finite measure, we define

$$T_r f(x) := \int_{r \leq \rho(x, y)} K(x, y) f(y) d\mu(y) = \int_{X \setminus B(x, r)} K(x, y) f(y) d\mu(y). \quad (10.0.2)$$

Theorem 10.0.1 (two-sided metric space Carleson). *For all integers $a \geq 4$ and real numbers $1 < q \leq 2$ the following holds. Let (X, ρ, μ, a) be a doubling metric measure space. Let Θ be a cancellative compatible collection of functions and let K be a two-sided Calderón–Zygmund kernel on (X, ρ, μ, a) . Assume that for every*

bounded measurable function g on X supported on a set of finite measure and all $r > 0$ we have

$$\|T_r g\|_2 \leq 2^{a^3} \|g\|_2. \quad (10.0.3)$$

Then for all Borel sets F and G in X and all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$, we have, with T defined in (1.1.14),

$$\left| \int_G T f \, d\mu \right| \leq \frac{2^{474a^3}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}. \quad (10.0.4)$$

For the remainder of this chapter, fix an integer $a \geq 4$, a doubling metric measure space (X, ρ, μ, a) and a two-sided Calderón–Zygmund kernel K as in Theorem 10.0.1.

The following lemma is proved in Section 10.1.

Lemma 10.0.2 (nontangential-from-simple). *Assume (10.0.3) holds. Then, for every bounded measurable function $g : X \rightarrow \mathbb{C}$ supported on a set of finite measure we have*

$$\|T_* g\|_2 \leq 2^{3a^3} \|g\|_2. \quad (10.0.5)$$

Proof of Theorem 10.0.1. Let $1 < q \leq 2$ be a real number. Let Θ be a cancellative compatible collection of functions. By the assumption (10.0.3), we can apply Lemma 10.0.2 to obtain for every bounded measurable $g : X \rightarrow \mathbb{C}$ supported on a set of finite measure,

$$\|T_* g\|_2 \leq 2^{3a^3} \|g\|_2. \quad (10.0.6)$$

Define

$$K'(x, y) := 2^{-2a^3} K(x, y).$$

Then K' is a two-sided Calderón–Zygmund kernel on (X, ρ, μ, a) . Denote the corresponding maximally truncated non-tangential singular operator by T'_* and the corresponding generalized Carleson operator by T' . With (10.0.6), we obtain for g as above,

$$\|T'_* g\|_2 \leq 2^{a^3} \|g\|_2. \quad (10.0.7)$$

Applying Theorem 1.1.1 for K' yields that for all Borel sets F and G in X and all Borel functions $f : X \rightarrow \mathbb{C}$ with $|f| \leq \mathbf{1}_F$, we have

$$\left| \int_G T' f \, d\mu \right| \leq \frac{2^{450a^3}}{(q-1)^6} \mu(G)^{1-\frac{1}{q}} \mu(F)^{\frac{1}{q}}.$$

This finishes the proof since for all $x \in X$,

$$T' f(x) = 2^{-2a^3} T f(x).$$

□

The proof of Lemma 10.0.2 relies on the following auxiliary lemma which is proved in Section 10.2.

Lemma 10.0.3 (Calderon-Zygmund Weak (1, 1)). *Let $f : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure and assume for some $r > 0$ that for every bounded measurable function $g : X \rightarrow \mathbb{C}$ supported on a set of finite measure,*

$$\|T_r g\|_2 \leq 2^{a^3} \|g\|_2. \quad (10.0.8)$$

Then for all $\alpha > 0$, we have

$$\mu(\{x \in X : |T_r f(x)| > \alpha\}) \leq \frac{2^{a^3+19a}}{\alpha} \int |f(y)| d\mu(y). \quad (10.0.9)$$

Throughout Section 10.2 and Section 10.1, for any measurable bounded function $w : X \rightarrow \mathbb{C}$, let $Mw : X \rightarrow [0, \infty)$ denote the corresponding Hardy–Littlewood maximal function defined in Proposition 2.0.6. Note: In some cases, a stronger formalization of Proposition 2.0.6 is used, where w is not necessarily bounded. In particular it is applied to $T_r g$. Apart from Proposition 2.0.6, Section 10.2 and Section 10.1 have no dependencies in the previous chapters.

10.1. Proof of Cotlar’s Inequality.

Lemma 10.1.1 (geometric series estimate). *For all real numbers $x \geq 4$,*

$$\sum_{n=0}^{\infty} 2^{-\frac{n}{x}} \leq 2^x.$$

Proof. By convexity, for all $0 \leq \lambda \leq 1$

$$2^{\lambda(-\frac{1}{4})} \leq \lambda 2^{-\frac{1}{4}} + (1-\lambda)2^0.$$

For $\lambda := \frac{4}{x}$, we obtain

$$2^{-\frac{1}{x}} \leq 1 - (1 - 2^{-\frac{1}{4}}) \frac{4}{x}.$$

We conclude

$$\sum_{n=0}^{\infty} 2^{-\frac{n}{x}} = \frac{1}{1 - 2^{-\frac{1}{x}}} \leq \frac{1}{4(1 - 2^{-\frac{1}{4}})} x \leq 2^x.$$

□

Lemma 10.1.2 (estimate x shift). *Let $0 < r$ and $x \in X$. Let $g : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. Then for all x' with $\rho(x, x') \leq r$.*

$$|T_r g(x) - T_r g(x')| \leq 2^{a^3+2a+2} M g(x).$$

Proof. By definition,

$$|T_r g(x) - T_r g(x')| = \left| \int_{r \leq \rho(x, y)} K(x, y) g(y) d\mu(y) - \int_{r \leq \rho(x', y)} K(x', y) g(y) d\mu(y) \right|. \quad (10.1.1)$$

We split the first integral in (10.1.1) into the domains $r \leq \rho(x, y) < 2r$ and $2r \leq \rho(x, y)$. The integral over the first domain we estimate by (10.1.2) below. For the second domain, we observe with $\rho(x, x') \leq r$ and the triangle inequality that $r \leq \rho(x', y)$. We therefore combine on this domain with the corresponding part of the second integral in (10.1.1) and estimate that by (10.1.3) below. The remaining part of the second integral in (10.1.1) we estimate by (10.1.4). Overall, we have estimated (10.1.1) by

$$\int_{r \leq \rho(x, y) < 2r} |K(x, y)| |g(y)| d\mu(y) \quad (10.1.2)$$

$$+ \left| \int_{2r \leq \rho(x,y)} (K(x,y) - K(x',y))g(y) d\mu(y) \right| \quad (10.1.3)$$

$$+ \int_{r \leq \rho(x',y), r \leq \rho(x,y) < 2r} |K(x',y)||g(y)| d\mu(y). \quad (10.1.4)$$

Using the bound on K in (1.1.11) and the doubling condition (1.1.2), we estimate (10.1.2) by

$$\int_{r \leq \rho(x,y) < 2r} \frac{2^{a^3}}{V(x,y)} |g(y)| d\mu(y) \leq \frac{2^{a^3}}{\mu(B(x,r))} \int_{r \leq \rho(x,y) < 2r} |g(y)| d\mu(y) \quad (10.1.5)$$

$$\leq \frac{2^{a^3} \cdot 2^a}{\mu(B(x,2r))} \int_{\rho(x,y) < 2r} |g(y)| d\mu(y). \quad (10.1.6)$$

Using the definition of Mg , we estimate (10.1.6) by

$$\leq 2^{a^3+a} Mg(x). \quad (10.1.7)$$

Similarly, in the domain of (10.1.4) we note by the triangle inequality and assumption on x' that $\rho(x',y) < 3r$ and thus we estimate (10.1.4) by

$$\frac{2^{a^3}}{\mu(B(x',r))} \int_{\rho(x',y) < 4r} |g(y)| d\mu(y) \leq 2^{a^3+2a} Mg(x) \quad (10.1.8)$$

We turn to the remaining term. Using (10.0.1), we estimate (10.1.3) by

$$\int_{2r \leq \rho(x,y)} \left(\frac{\rho(x,x')}{\rho(x,y)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x,y)} |g(y)| d\mu(y) \quad (10.1.9)$$

We decompose and estimate (10.1.3) with the triangle inequality by

$$\sum_{j=1}^{\infty} \int_{2^j r \leq \rho(x,y) < 2^{j+1} r} \left(\frac{\rho(x,x')}{\rho(x,y)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x,y)} |g(y)| d\mu(y) \quad (10.1.10)$$

$$\leq \sum_{j=1}^{\infty} (2^{-j})^{\frac{1}{a}} \int_{2^j r \leq \rho(x,y) < 2^{j+1} r} \frac{2^{a^3}}{\mu(B(x,2^j r))} |g(y)| d\mu(y) \quad (10.1.11)$$

$$\leq \sum_{j=1}^{\infty} 2^{-\frac{j}{a}} \frac{2^{a^3+a}}{\mu(B(x,2^{j+1} r))} \int_{\rho(x,y) < 2^{j+1} r} |g(y)| d\mu(y) \quad (10.1.12)$$

$$\leq 2^{a^3+a} \sum_{j=1}^{\infty} 2^{-\frac{j}{a}} Mg(x). \quad (10.1.13)$$

Using Lemma 10.1.1, we estimate (10.1.13) by

$$\leq 2^{a^3+2a} Mg(x). \quad (10.1.14)$$

Summing the estimates for (10.1.2), (10.1.3), and (10.1.4) proves the lemma. \square

Lemma 10.1.3 (Cotlar control). *Let $0 < r \leq R$ and $x \in X$. Let $g : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. Then for all $x' \in X$ with $\rho(x, x') \leq \frac{R}{4}$ we have*

$$|T_R g(x)| \leq |T_r(g - g\mathbf{1}_{B(x, \frac{R}{2})})(x')| + 2^{a^3+4a+1} M g(x). \quad (10.1.15)$$

Proof. Let x and x' be given with $\rho(x, x') \leq \frac{R}{4}$. By Lemma 10.1.2, we estimate the left-hand-side of (10.1.15) by

$$|T_R(g)(x')| + 2^{a^3+2a+2} M g(x). \quad (10.1.16)$$

We have

$$T_R(g)(x') = \int_{R \leq \rho(x', y)} K(x', y) g(y) d\mu(y). \quad (10.1.17)$$

On the domain $R \leq \rho(x', y)$, we have $\frac{R}{2} \leq \rho(x, y)$. Hence we may write for (10.1.17)

$$\begin{aligned} T_R(g)(x') &= \int_{R \leq \rho(x', y)} K(x', y) (g - g\mathbf{1}_{B(x, \frac{R}{2})})(y) d\mu(y) \\ &= T_R(g - g\mathbf{1}_{B(x, \frac{R}{2})})(x'). \end{aligned} \quad (10.1.18)$$

Combining the estimate (10.1.16) with the identification (10.1.18), we obtain

$$|T_R g(x)| \leq |T_R(g - g\mathbf{1}_{B(x, \frac{R}{2})})(x')| + 2^{a^3+2a+2} M g(x). \quad (10.1.19)$$

We have

$$\begin{aligned} &(T_r - T_R)(g - g\mathbf{1}_{B(x, \frac{R}{2})})(x') \\ &= \int_{B(x', R) \setminus B(x', r)} K(x', y) (g - g\mathbf{1}_{B(x, \frac{R}{2})})(y) d\mu(y) \\ &= \int_{B(x', R) \setminus (B(x', r) \cup B(x, \frac{R}{2}))} K(x', y) g(y) d\mu(y) \end{aligned} \quad (10.1.20)$$

As $\frac{R}{2} \leq \rho(x, y)$ together with $\rho(x, x') \leq \frac{R}{4}$ implies $\frac{R}{4} \leq \rho(x', y)$, we can estimate the absolute value of (10.1.20) with (1.1.11) by

$$\begin{aligned} &\leq \frac{2^{a^3}}{\mu(B(x', \frac{R}{4}))} \int_{B(x, 2R) \setminus B(x', \frac{R}{4})} |g(y)| d\mu(y) \\ &\leq \frac{2^{a^3+a}}{\mu(B(x', \frac{R}{2}))} \int_{B(x, 2R)} |g(y)| d\mu(y) \\ &\leq 2^{a^3+a} \frac{\mu(B(x, 2R))}{\mu(B(x, \frac{R}{4}))} M g(x) \leq 2^{a^3+4a} M g(x). \end{aligned}$$

By the triangle inequality, (10.1.15) follows now from (10.1.19) and the estimate for (10.1.20). \square

Lemma 10.1.4 (Cotlar sets). *Assume that (10.0.3) holds. Let $0 < r \leq R$ and $x \in X$. Let $g : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. Then the measure $|F_1|$ of the set F_1 of all $x' \in B(x, \frac{R}{4})$ such that*

$$|T_r g(x')| > 4M(T_r g)(x) \quad (10.1.21)$$

is less than or equal to $\mu(B(x, \frac{R}{4}))/4$. Moreover, the measure $|F_2|$ of the set F_2 of all $x' \in B(x, \frac{R}{4})$ such that

$$|T_r(g\mathbf{1}_{B(x, \frac{R}{2})})(x')| > 2^{a^3+20a+2}Mg(x) \quad (10.1.22)$$

is less than or equal to $\mu(B(x, \frac{R}{4}))/4$.

Proof. Let r , R , x and g be given. If $M(T_r g)(x) = 0$, then $T_r g$ is zero almost everywhere and the estimate on $|F_1|$ is trivial. Assume $M(T_r g)(x) > 0$. We have with (10.1.21)

$$M(T_r g)(x) \geq \frac{1}{\mu(B(x, \frac{R}{4}))} \int_{B(x, \frac{R}{4})} |T_r g(x')| dx' \quad (10.1.23)$$

$$\geq \frac{1}{\mu(B(x, \frac{R}{4}))} \int_{F_1} 4M(T_r g)(x) dx'. \quad (10.1.24)$$

Dividing by $M(T_r g)(x)$ gives

$$1 \geq \frac{4}{\mu(B(x, \frac{R}{4}))} |F_1|. \quad (10.1.25)$$

This gives the desired bound for the measure of F_1 . We turn to the set F_2 . Similarly as above we may assume $Mg(x) > 0$. The set F_2 is then estimated with Lemma 10.0.3 by

$$\frac{2^{a^3+19a}}{2^{a^3+20a+2}Mg(x)} \int |g\mathbf{1}_{B(x, \frac{R}{2})}|(y) d\mu(y) \quad (10.1.26)$$

$$\leq \frac{1}{2^{a+2}Mg(x)} \mu(B(x, \frac{R}{2}))Mg(x) \leq \frac{\mu(B(x, \frac{R}{4}))}{4}. \quad (10.1.27)$$

This gives the desired bound for the measure of F_2 . \square

Lemma 10.1.5 (Cotlar estimate). *Assume that (10.0.3) holds. Let $0 < r \leq R$ and $x \in X$. Let $g : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. Then*

$$|T_R g(x)| \leq 2^2 M(T_r g)(x) + 2^{a^3+20a+3} Mg(x). \quad (10.1.28)$$

Proof. By Lemma 10.1.4, the set of all $x' \in B(x, \frac{R}{4})$ such that at least one of the conditions (10.1.21) and (10.1.22) is satisfied has measure less than or equal to $\mu(B(x, \frac{R}{4}))/2$ and hence is not all of $B(x, \frac{R}{4})$. Pick an $x' \in B(x, \frac{R}{4})$ such that both conditions are not satisfied. Applying Lemma 10.1.3 for this x' and using the triangle inequality estimates the left-hand side of (10.1.28) by

$$4M(T_r g)(x) + 2^{a^3+20a+2}Mg(x) + 2^{a^3+4a+1}Mg(x). \quad (10.1.29)$$

This proves the lemma. \square

For the next Lemma, we define the following operation.

$$T_*^r g(x) := \sup_{r < R} \sup_{x' \in B(x, R)} |T_R(g)(x')|. \quad (10.1.30)$$

Lemma 10.1.6 (simple nontangential operator). *Assume that (10.0.3) holds. For every $r > 0$ and every bounded measurable function g supported on a set of finite measure the function $T_*^r g$ is lower-semicontinuous and we have*

$$\|T_*^r g\|_2 \leq 2^{a^3+24a+6} \|g\|_2, \quad (10.1.31)$$

Proof. For a fixed λ we note that (rewriting $x' \in B(x, R)$ as $x \in B(x', R)$):

$$\{x : T_*^r g(x) > \lambda\} = \bigcup_{r < R} \bigcup_{\{x' : |T_R(g)(x')| > \lambda\}} B(x', R) \quad (10.1.32)$$

The right-hand side is open, hence $T_*^r g$ is lower-semicontinuous by definition.

For the second part, with Lemma 10.1.2 and the triangle inequality, we estimate for every $x \in X$

$$T_*^r g(x) \leq 2^{a^3+2a+2} M g(x) + \sup_{r < R} |T_R(g)(x)|. \quad (10.1.33)$$

Using further Lemma 10.1.5, we estimate

$$T_*^r g(x) \leq 2^{a^3+2a+2} M g(x) + 2^{a^3+20a+3} M g(x) + 2^2 M(T_r g)(x). \quad (10.1.34)$$

Taking the L^2 norm and using Proposition 2.0.6 with $a = 4$ and $p_2 = 2$ and $p_1 = 1$, we obtain

$$\|T_*^r g\|_2 \leq 2^{a^3+20a+4} \|M g\|_2 + 2^2 \|M(T_r g)\|_2 \quad (10.1.35)$$

$$\leq 2^{a^3+24a+5} \|g\|_2 + 2^{4a+3} \|T_r g\|_2. \quad (10.1.36)$$

Applying (10.0.3) gives

$$\|T_*^r g\|_2 \leq 2^{a^3+24a+5} \|g\|_2 + 2^{a^3+4a+3} \|g\|_2. \quad (10.1.37)$$

This shows (10.1.31) and completes the proof of the inequality. \square

In order to pass from the one-sided truncation in T_r and T_*^r to the two-sided truncation in T_* , we show in the following two lemmas that the integral in (1.1.13) can be exchanged for an integral over the difference of two balls.

Lemma 10.1.7 (small annulus). *Let $f : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. Let $x \in X$ and $R > 0$. Then, for all $\epsilon > 0$, there exists some $\delta > 0$ such that*

$$\left| \int_{R < \rho(x, y) < R + \delta} K(x, y) f(y) d\mu(y) \right| \leq \epsilon \quad (10.1.38)$$

and

$$\left| \int_{R - \delta < \rho(x, y) < R} K(x, y) f(y) d\mu(y) \right| \leq \epsilon. \quad (10.1.39)$$

Proof. We only prove the second inequality, the first one is analogous. Note that the integrand is bounded in $X \setminus B(x, \frac{R}{2})$. So for $0 < \delta \leq \frac{R}{2}$,

$$\begin{aligned} & \left| \int_{R - \delta < \rho(x, y) < R} K(x, y) f(y) d\mu(y) \right| \\ & \leq \frac{2^{a^3}}{\mu(B(x, \frac{R}{2}))} \sup_{y \in X} |f(y)| \cdot \mu(\{y \in X : R - \delta < \rho(x, y) < R\}). \end{aligned}$$

By continuity from above of μ , the right factor becomes arbitrarily small as $\delta \rightarrow 0$. Thus, for small enough δ , the whole expression is $\leq \epsilon$. \square

Lemma 10.1.8 (nontangential operator boundary). *Let $f : X \rightarrow \mathbb{C}$ be a bounded measurable function supported on a set of finite measure. For all $x \in X$,*

$$T_*f(x) = \sup_{R_1 < R_2} \sup_{x' \in B(x, R_1)} \left| \int_{B(x', R_2) \setminus B(x', R_1)} K(x', y) f(y) \, d\mu(y) \right| \quad (10.1.40)$$

Proof. We show two inequalities. Let $\epsilon > 0$. Let $R_1 < R_2$ and $x' \in B(x, R_1)$. Then for small enough $\delta > 0$,

$$\left| \int_{R_1 < \rho(x', y) < R_2} K(x', y) f(y) \, d\mu(y) \right| \quad (10.1.41)$$

$$\leq \left| \int_{R_1 < \rho(x', y) < R_1 + \delta} K(x', y) f(y) \, d\mu(y) \right| \quad (10.1.42)$$

$$+ \left| \int_{R_1 + \delta \leq \rho(x', y) < R_2} K(x', y) f(y) \, d\mu(y) \right|. \quad (10.1.43)$$

By Lemma 10.1.7, we can choose δ such that (10.1.42) is bounded by ϵ . Without loss of generality, we can assume $R_1 + \delta < R_2$. Then (10.1.43) is bounded by the right hand side of (10.1.40) and we obtain

$$\leq \epsilon + \sup_{R_1 < R_2} \sup_{x' \in B(x, R_1)} \left| \int_{B(x', R_2) \setminus B(x', R_1)} K(x', y) f(y) \, d\mu(y) \right|.$$

The inequality still holds when taking the suprema over $R_1 < R_2$ and $\rho(x, x') < R_1$ in (10.1.41). Since $\epsilon > 0$ was arbitrary, this proves the first inequality.

The other direction is similar. Let $\epsilon > 0$. Let $R_1 < R_2$ and $x' \in B(x, R_1)$. Then for $\delta > 0$,

$$\left| \int_{B(x', R_2) \setminus B(x', R_1)} K(x', y) f(y) \, d\mu(y) \right| \quad (10.1.44)$$

$$\leq \left| \int_{R_1 - \delta < \rho(x', y) < R_1} K(x', y) f(y) \, d\mu(y) \right| \quad (10.1.45)$$

$$+ \left| \int_{R_1 - \delta < \rho(x', y) < R_2} K(x', y) f(y) \, d\mu(y) \right|. \quad (10.1.46)$$

By Lemma 10.1.7, we can choose δ such that (10.1.45) is bounded by ϵ . Without loss of generality, we can assume $\rho(x, x') < R_1 - \delta$. Then (10.1.46) is bounded by the left hand side of (10.1.40) and we obtain

$$\leq \epsilon + \sup_{R_1 < R_2} \sup_{x' \in B(x, R_1)} \left| \int_{R_1 < \rho(x', y) < R_2} K(x', y) f(y) \, d\mu(y) \right|.$$

The inequality still holds when taking the suprema over $R_1 < R_2$ and $\rho(x, x') < R_1$ in (10.1.41). Since $\epsilon > 0$ was arbitrary, this proves the second inequality. \square

Proof of Lemma 10.0.2. Fix g as in the Lemma. Applying Lemma 10.1.6 with a sequence of r tending to 0 and using Lebesgue monotone convergence shows

$$\|T_*^0 g\|_2 \leq 2^{a^3+24a+6} \|g\|_2, \quad (10.1.47)$$

where

$$T_*^0 g(x) := \sup_{0 < R} \sup_{x' \in B(x, R)} \left| \int_{X \setminus B(x', R)} K(x', y) g(y) d\mu(y) \right|. \quad (10.1.48)$$

We now write using Lemma 10.1.8 and the triangle inequality,

$$\begin{aligned} T_* g(x) &\leq \sup_{0 < R_1 < R_2} \sup_{x' \in B(x, R_1)} \left| \int_{X \setminus B(x', R_1)} K(x', y) g(y) d\mu(y) \right| \\ &\quad + \sup_{0 < R_1 < R_2} \sup_{x' \in B(x, R_1)} \left| \int_{X \setminus B(x', R_2)} K(x', y) g(y) d\mu(y) \right|. \end{aligned}$$

Noting that the first integral does not depend on R_2 and estimating the second summand by the larger supremum over all $x' \in B(x, R_2)$, at which time the integral does not depend on R_1 , we estimate further

$$\begin{aligned} &\leq \sup_{0 < R_1} \sup_{x' \in B(x, R_1)} \left| \int_{X \setminus B(x', R_1)} K(x', y) g(y) d\mu(y) \right| \\ &\quad + \sup_{0 < R_2} \sup_{x' \in B(x, R_2)} \left| \int_{X \setminus B(x', R_2)} K(x', y) g(y) d\mu(y) \right|. \end{aligned}$$

Applying the triangle inequality on the left-hand side of (10.0.5) and applying (10.1.47) twice proves (10.0.5). This completes the proof of Lemma 10.0.2. \square

10.2. Calderón-Zygmund Decomposition. Calderón-Zygmund decomposition is a tool to extend L^2 bounds to L^p bounds with $p < 2$ or to the so-called weak $(1, 1)$ type endpoint bound. It is classical and can be found in [Ste93].

The following lemma is Theorem 3.1(b) in [Ste93]. The proof uses Proposition 2.0.6.

Lemma 10.2.1 (Maximal theorem). *Let $f : X \rightarrow \mathbb{C}$ be bounded, measurable, supported on a set of finite measure, and let $\alpha > 0$. Then*

$$\mu(\{x \in X : Mf(x) > \alpha\}) \leq \frac{2^{2a}}{\alpha} \int |f(y)| d\mu(y). \quad (10.2.1)$$

Proof. By definition, for each $x \in X$ with $Mf(x) > \alpha$, there exists a ball B_x such that $x \in B_x$ and

$$\alpha \mu(B_x) < \int_{B_x} |f(y)| d\mu(y). \quad (10.2.2)$$

Since $\{x \in X : Mf(x) > \alpha\}$ is open and μ is inner regular on open sets, it suffices to show that

$$\mu(E) \leq \frac{2^{2a}}{\alpha} \int |f(y)| d\mu(y)$$

for every compact $E \subset \{x \in X : Mf(x) > \alpha\}$. For such an E , by compactness, we can select a finite subcollection $\mathcal{B} \subset \{B_x : x \in E\}$ that covers E . By (2.0.43) applied to (10.2.2),

$$\alpha\mu(\bigcup \mathcal{B}) \leq 2^{2a} \int |f(y)| d\mu(y) \quad (10.2.3)$$

and hence

$$\mu(E) \leq \mu(\bigcup \mathcal{B}) \leq \frac{2^{2a}}{\alpha} \int |f(y)| d\mu(y).$$

□

Lemma 10.2.2 (Lebesgue differentiation). *Let f be a bounded measurable function supported on a set of finite measure. Then for μ almost every x , we have*

$$\lim_{n \rightarrow \infty} \frac{1}{\mu(B_n)} \int_{B_n} f(y) dy = f(x),$$

where $\{B_n\}_{n \geq 1}$ is a sequence of balls with radii $r_n > 0$ such that $x \in B_n$ for each $n \geq 1$ and

$$\lim_{n \rightarrow \infty} r_n = 0.$$

Proof. This follows from the Lebesgue differentiation theorem, which is already formalized in Lean. □

Lemma 10.2.3 (Disjoint family countable). *In a doubling metric measure space (X, ρ, μ, a) , every disjoint family of balls $B_j = B(x_j, r_j)$, $j \in J$, is countable.*

Proof. Choose an arbitrary $x \in X$ as reference point. For $q, Q \in \mathbb{Q}_+$, let $J_{q,Q}$ denote the set of all $j \in J$ such that $B_j \subset B(x, Q)$ and $r_j \geq q$. It suffices to show that all the $J_{q,Q}$ are finite. Indeed, for all $j \in J_{q,Q}$,

$$\mu(B(x, Q)) \leq \mu(B(x_j, 2Q)) = \mu(B(x_j, \frac{2Q}{r_j} r_j)) \leq 2^{a \log_2 \lceil \frac{2Q}{r_j} \rceil} \mu(B_j).$$

Since the B_j are disjoint,

$$|J_{q,Q}| \mu(B(x, Q)) \leq 2^{a \log_2 \lceil \frac{2Q}{q} \rceil} \sum_{j \in J_{q,Q}} \mu(B_j) \leq 2^{a \log_2 \lceil \frac{2Q}{q} \rceil} \mu(B(x, Q)) \quad (10.2.4)$$

and hence $|J_{q,Q}| \leq 2^{a \log_2 \lceil \frac{2Q}{q} \rceil}$. □

The following lemma corresponds to Lemma 3.2 in [Ste93] with additional proof of the bounded intersection property taken from the proof of Proposition 7.1. It uses the following notation, which will be used throughout the rest of this section: whenever B_j denotes a ball $B_j = B(x_j, r_j)$, define

$$B_{n,j} = B(x_j, nr_j).$$

Lemma 10.2.4 (Ball covering). *Given an open set $O \neq X$, there exists a countable family of balls $B_j = B(x_j, r_j)$ such that*

$$B_j \cap B_{j'} = \emptyset \quad \text{for } j \neq j', \quad (10.2.5)$$

and

$$\bigcup_j B_{3,j} = O, \quad (10.2.6)$$

and

$$B_{\tau,j} \cap (X \setminus O) \neq \emptyset \quad \text{for all } j \quad (10.2.7)$$

and we have the bounded intersection property that each $x \in O$ is contained in at most 2^{6a} of the $B_{3,j}$.

Proof. Define for $x \in O$,

$$\delta(x) := \sup\{\delta \in \mathbb{R} : B(x, \delta) \subset O\}. \quad (10.2.8)$$

Since O is open, and $O \neq X$, we have

$$0 < \delta(x) < \infty. \quad (10.2.9)$$

Using Zorn's Lemma, we select a maximal disjoint subfamily of $\{B(x, \frac{\delta(x)}{6}) : x \in O\}$. We obtain a (by Lemma 10.2.3 countable) family of balls $B_j = B(x_j, \frac{\delta(x_j)}{6}), j \in J$ such that (10.2.5), (10.2.7), and $\bigcup_j B_{3,j} \subset O$ are also immediate. For the other inclusion, first observe that for $x, y \in X$, if $B(x, \frac{\delta(x)}{6}) \cap B(y, \frac{\delta(y)}{6}) \neq \emptyset$, then

$$\delta(x) \leq \rho(x, y) + \delta(y) \leq \left(\frac{\delta(x)}{6} + \frac{\delta(y)}{6}\right) + \delta(y) = \frac{\delta(x)}{6} + \frac{7\delta(y)}{6},$$

so

$$\delta(x) \leq 2\delta(y). \quad (10.2.10)$$

Now let $z \in O$. By maximality, there exists some $j \in J$ with $B(z, \frac{\delta(z)}{6}) \cap B_j \neq \emptyset$. By (10.2.10),

$$\rho(z, x_j) < \frac{\delta(z)}{6} + \frac{\delta(x_j)}{6} \leq \frac{3\delta(x_j)}{6} = 3r_j$$

and thus $z \in B_{3,j}$.

We now turn to the bounded intersection property. Assume that for some j_1, \dots, j_N ,

$$z \in \bigcap_{k=1}^N B_{3,j_k}. \quad (10.2.11)$$

Similarly as above, observe for $1 \leq k \leq N$,

$$\delta(z) \leq \rho(z, x_{j_k}) + \delta(x_{j_k}) \leq \frac{\delta(x_{j_k})}{2} + \delta(x_{j_k}) = \frac{3\delta(x_{j_k})}{2} \quad (10.2.12)$$

and

$$\delta(x_{j_k}) \leq \rho(x_{j_k}, z) + \delta(z) \leq \frac{\delta(x_{j_k})}{2} + \delta(z),$$

so

$$\delta(x_{j_k}) \leq 2\delta(z). \quad (10.2.13)$$

By (10.2.12) and (10.2.13), for all $1 \leq k \leq N$, $B(z, \frac{\delta(z)}{6}) \subset B(x_{j_k}, 5r_{j_k})$ and $B_{j_k} \subset B(z, \frac{8\delta(z)}{6})$. Using this and (10.2.5), we obtain

$$N\mu(B(z, \frac{\delta(z)}{6})) \leq \sum_{k=1}^N \mu(B(x_{j_k}, 5r_{j_k})) \leq 2^{3a} \sum_{k=1}^N \mu(B_{j_k}) \quad (10.2.14)$$

$$= 2^{3a} \mu(\bigcup_{k=1}^N B_{j_k}) \leq 2^{3a} \mu(B(z, \frac{8\delta(z)}{6})) \leq 2^{6a} \mu(B(z, \frac{\delta(z)}{6})) \quad (10.2.15)$$

and conclude $N \leq 2^{6a}$. \square

Most of the next lemma and its proof is taken from Theorem 4.2 in [Ste93].

Lemma 10.2.5 (Calderon Zygmund decomposition). *Let f be a bounded, a.e. measurable function supported on a set of finite measure and let $\alpha > \frac{1}{\mu(X)} \int |f| d\mu$. Then there exists a bounded a.e. measurable function g supported on a set of finite measure, a countable family of balls $B_{3,j}$ (where we allow $B_{3,1} = X$ in the special case that $\mu(X) < \infty$) such that each $x \in X$ is contained in at most 2^{6a} of the $B_{3,j}$, and a countable family of a.e. measurable functions $\{b_j\}_{j \in J}$ such that for all $x \in X$*

$$f(x) = g(x) + \sum_j b_j(x) \quad (10.2.16)$$

and such that the following holds. For almost every $x \in X$,

$$|g(x)| \leq 2^{3a} \alpha. \quad (10.2.17)$$

We have

$$\int |g(y)| d\mu(y) \leq \int |f(y)| d\mu(y). \quad (10.2.18)$$

For every j

$$\text{supp } b_j \subset B_{3,j}. \quad (10.2.19)$$

For every j

$$\int_{B_{3,j}} b_j(x) d\mu(x) = 0, \quad (10.2.20)$$

and

$$\int_{B_{3,j}} |b_j(x)| d\mu(x) \leq 2^{2a+1} \alpha \mu(B_{3,j}). \quad (10.2.21)$$

We have

$$\sum_j \mu(B_{3,j}) \leq \frac{2^{4a}}{\alpha} \int |f(y)| d\mu(y) \quad (10.2.22)$$

and

$$\sum_j \int_{B_{3,j}} |b_j(y)| d\mu(y) \leq 2 \int |f(y)| d\mu(y). \quad (10.2.23)$$

Proof. Let $E_\alpha := \{x \in X : Mf(x) > \alpha\}$. Then E_α is open. Assume first that $E_\alpha \neq X$. We apply Lemma 10.2.4 with $O = E_\alpha$ to obtain the family $B_j, j \in J$. Without loss of generality, we can assume $J = \mathbb{N}$. Define inductively

$$Q_j := B_{3,j} \setminus \left(\bigcup_{i < j} Q_i \cup \bigcup_{i > j} B_i \right). \quad (10.2.24)$$

Then $B_j \subset Q_j \subset B_{3,j}$, the Q_j are pairwise disjoint and $\bigcup_j Q_j = E_\alpha$. Define

$$g(x) := \begin{cases} f(x), & x \in X \setminus E_\alpha, \\ \frac{1}{\mu(Q_j)} \int_{Q_j} f(y) d\mu(y), & x \in Q_j, \end{cases} \quad (10.2.25)$$

and, for each j ,

$$b_j(x) := \mathbf{1}_{Q_j}(x) \left(f(x) - \frac{1}{\mu(Q_j)} \int_{Q_j} f(y) d\mu(y) \right). \quad (10.2.26)$$

Then (10.2.16), (10.2.19) and (10.2.20) are true by construction. Boundedness of g is immediate from the definition, as is the fact that $\text{supp } g$ is contained within $\text{supp } f \cup \bigcup_j Q_j$. Now the fact that $\text{supp } g$ has finite measure follows from $\bigcup_j Q_j = E_\alpha$ and (2.0.46).

For (10.2.17), we first do the case $x \in X \setminus E_\alpha$. By definition of Mf ,

$$\frac{1}{\mu(B)} \int_B |f(y)| d\mu(y) \leq \alpha \quad (10.2.27)$$

for every ball $B \subset X$ with $x \in B$. It follows by Lemma 10.2.2 that for almost every $x \in X \setminus E_\alpha$, $|f(x)| \leq \alpha$. In the case $x \in E_\alpha$, there exists some $j \in J$ with $x \in Q_j$ and we have that

$$\frac{1}{\mu(B_{7,j})} \int_{B_{7,j}} |f(y)| d\mu(y) \leq \alpha \quad (10.2.28)$$

because $B_{7,j} \cap (X \setminus E_\alpha) \neq \emptyset$. We get

$$|g(x)| \leq \frac{1}{\mu(Q_j)} \int_{Q_j} |f(y)| d\mu(y) \leq \frac{1}{\mu(B_j)} \int_{B_{7,j}} |f(y)| d\mu(y) \leq 2^{3\alpha} \alpha. \quad (10.2.29)$$

To prove (10.2.18), we estimate

$$\begin{aligned} \int |g(z)| d\mu(z) &\leq \int_{X \setminus E_\alpha} |f(z)| d\mu(z) + \sum_j \int_{Q_j} \frac{1}{\mu(Q_j)} \int_{Q_j} |f(y)| d\mu(y) d\mu(z) \\ &= \int |f(z)| d\mu(z). \end{aligned}$$

Using the triangle inequality, we have that

$$\int_{B_{3,j}} |b_j(y)| dy \leq \int_{Q_j} |f(y)| d\mu(y) + \int_{Q_j} \frac{1}{\mu(Q_j)} \int_{Q_j} |f(x)| d\mu(x) d\mu(y) \quad (10.2.30)$$

$$= 2 \int_{Q_j} |f(y)| dy. \quad (10.2.31)$$

With (10.2.28), we estimate further

$$\leq 2 \int_{B_{7,j}} |f(y)| dy \leq 2\mu(B_{7,j})\alpha \leq 2^{2a+1}\alpha\mu(B_{3,j}) \quad (10.2.32)$$

to obtain (10.2.21). Further, summing up (10.2.30) in j yields (10.2.23). At last, we estimate with Lemma 10.2.1

$$\sum_j \mu(B_{3,j}) \leq 2^{2a} \sum_j \mu(B_j) \leq 2^{2a} \mu(E_\alpha) \leq \frac{2^{4a}}{\alpha} \int |f(y)| d\mu(y), \quad (10.2.33)$$

proving (10.2.22).

Assume now that $E_\alpha = X$. It follows from Lemma 10.2.1 that then $\mu(X) < \infty$. Define

$$g := \frac{1}{\mu(X)} \int |f(y)| d\mu(y)$$

and

$$b_1 := f - g.$$

Then $f = g + b_1$ and $\text{supp } b_1 \subset B_{3,1} := X$ and (10.2.16), (10.2.18), (10.2.19), (10.2.20) all hold immediately. By assumption, $\alpha > \frac{1}{\mu(X)} \int |f| d\mu = g$, so (10.2.17) holds. We also have, using the definitions and the same assumption,

$$\int |b_1(y)| d\mu(y) \leq 2 \int |f(y)| d\mu(y) \leq 2\alpha\mu(X), \quad (10.2.34)$$

which verifies both (10.2.23) and (10.2.21). Finally, by Lemma 10.2.1,

$$\mu(X) \leq \frac{2^{2a}}{\alpha} \int |f(y)| d\mu(y),$$

which shows (10.2.22). \square

We use Lemma 10.2.5 to prove Lemma 10.0.3. For the remainder of this section, let $f : X \rightarrow \mathbb{C}$, $r > 0$ and $\alpha > 0$ as in Lemma 10.0.3. We define the constant

$$c := 2^{-a^3-12a-4} \quad (10.2.35)$$

and $\alpha' := c\alpha$. If $\alpha' \leq \frac{1}{\mu(X)} \int |f| d\mu$, then we directly have

$$\mu(\{x \in X : |T_r f(x)| > \alpha\}) \leq \mu(X) \leq \frac{1}{\alpha'} \int |f(y)| d\mu(y) \leq \frac{2^{a^3+19a}}{\alpha} \int |f(y)| d\mu(y),$$

which proves (10.0.9). So assume from now on that $\alpha' > \frac{1}{\mu(X)} \int |f| d\mu$. Using Lemma 10.2.5 for f and α' , we obtain the decomposition

$$f = g + b = g + \sum_j b_j$$

such that the properties (10.2.16)-(10.2.23) are satisfied (with α' replacing α). Let

$$B_{3,j} = B(x_j, 3r_j) \quad (10.2.36)$$

as in the lemma. Then

$$B_{6,j} = B(x_j, 6r_j) \quad (10.2.37)$$

is a ball with the same center as $B_{3,j}$ but with

$$\mu(B_{6,j}) \leq 2^a \mu(B_{3,j}). \quad (10.2.38)$$

Let

$$\Omega := \bigcup_j B_{6,j}. \quad (10.2.39)$$

We deal with $T_r g$ and $T_r b$ separately in the following lemmas.

Lemma 10.2.6 (Estimate good).

$$\mu(\{x \in X : |T_r g(x)| > \alpha/2\}) \leq \frac{2^{2a^3+3a+2} c}{\alpha} \int |f(y)| d\mu(y).$$

Proof. We estimate using monotonicity of the integral

$$\mu(\{x \in X : |T_r g(x)| > \alpha/2\}) \leq \frac{4}{\alpha^2} \int |T_r g(y)|^2 d\mu(y).$$

Using (10.0.8) followed by (10.2.17) and (10.2.18), we estimate the right hand side above by

$$\leq \frac{4 \cdot 2^{2a^3}}{\alpha^2} \int |g(y)|^2 d\mu(y) \leq \frac{2^{2a^3+3a+2} c}{\alpha} \int |g(y)| dy \leq \frac{2^{2a^3+3a+2} c}{\alpha} \int |f(y)| d\mu(y). \quad (10.2.40)$$

□

Lemma 10.2.7 (Estimate bad partial). *Let $x \in X \setminus \Omega$. Then*

$$|T_r b(x)| \leq 3F(x) + \alpha/8,$$

where

$$F(x) := 2^{a^3+2a+1} c \alpha \sum_{j \in J} \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{\mu(B_{3,j})}{V(x, x_j)}.$$

Proof. We decompose the index set J into the following disjoint sets:

$$\begin{aligned} \mathcal{J}_1(x) &:= \{j : r + 3r_j \leq \rho(x, x_j)\}, \\ \mathcal{J}_2(x) &:= \{j : r - 3r_j \leq \rho(x, x_j) < r + 3r_j\}, \\ \mathcal{J}_3(x) &:= \{j : \rho(x, x_j) < r - 3r_j\}. \end{aligned}$$

Then

$$|T_r b(x)| \leq \sum_{j \in \mathcal{J}_1(x)} |T_r b_j(x)| \quad (10.2.41)$$

$$+ \sum_{j \in \mathcal{J}_2(x)} |T_r b_j(x)| \quad (10.2.42)$$

$$+ \sum_{j \in \mathcal{J}_3(x)} |T_r b_j(x)|. \quad (10.2.43)$$

For all $j \in \mathcal{J}_3(x)$, $\text{supp } b_j \subset B_{3,j} \subset B(x, r)$, and thus $T_r b_j(x) = 0$, so (10.2.43) = 0.

Next, for $j \in \mathcal{J}_1(x)$, $\text{supp } b_j \subset B_{3,j} \subset X \setminus B(x, r)$, and we have

$$T_r b_j(x) = \int_{X \setminus B(x, r)} K(x, y) b_j(y) d\mu(y) = \int_{B_{3,j}} K(x, y) b_j(y) d\mu(y).$$

Using (10.2.20), the above is equal to

$$\int_{B_{3,j}} (K(x, y) - K(x, x_j)) b_j(y) d\mu(y).$$

Since $x \in X \setminus \Omega$, we have for each $y \in B_{3,j}$ that

$$\rho(x, x_j) \geq 6r_j > 2\rho(x_j, y), \quad (10.2.44)$$

so we can apply (1.1.12) to estimate

$$\begin{aligned} (10.2.41) &\leq \sum_{j \in \mathcal{J}_1(x)} \int_{B_{3,j}} \left(\frac{\rho(x_j, y)}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{2^{a^3}}{V(x, x_j)} |b_j(y)| d\mu(y) \\ &\leq 2^{a^3} \sum_j \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{1}{V(x, x_j)} \int_{B_{3,j}} |b_j(y)| dy \end{aligned} \quad (10.2.45)$$

and by (10.2.21),

$$\leq 2^{a^3+2a+1} c\alpha \sum_j \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{\mu(B_{3,j})}{V(x, x_j)} = F(x). \quad (10.2.46)$$

Next, we estimate (10.2.42). For each $j \in \mathcal{J}_2(x)$, set

$$d_j := \frac{1}{\mu(B_{3,j})} \int_{B_{3,j}} \mathbf{1}_{X \setminus B(x, r)}(y) b_j(y) dy.$$

Then by (10.2.21)

$$|d_j| \leq 2^{2a+1} c\alpha. \quad (10.2.47)$$

For each $j \in \mathcal{J}_2(x)$, we have

$$\begin{aligned} T_r b_j(x) &= \int_{B_{3,j}} K(x, y) (\mathbf{1}_{X \setminus B(x, r)}(y) b_j(y) - d_j) dy + \int_{B_{3,j}} d_j K(x, y) dy \\ &= \int_{B_{3,j}} (K(x, y) - K(x, x_j)) (\mathbf{1}_{X \setminus B(x, r)}(y) b_j(y) - d_j) dy + \int_{B_{3,j}} d_j K(x, y) dy. \end{aligned}$$

Thus, using the triangle inequality, the equation above and (10.2.47), we obtain

$$\begin{aligned} |T_r b_j(x)| &\leq \\ &\int_{B_{3,j}} |K(x, y) - K(x, x_j)| (|b_j(y)| + 2^{2a+1} c\alpha) dy + 2^{2a+1} c\alpha \int_{B_{3,j}} |K(x, y)| dy. \end{aligned} \quad (10.2.48)$$

By (10.2.44), we can apply (1.1.12) and arguing as in (10.2.46), we get that

$$(10.2.42) \leq 2F(x) + 2^{2a+1} c\alpha \sum_{j \in \mathcal{J}_2(x)} \int_{B_{3,j}} |K(x, y)| d\mu(y), \quad (10.2.49)$$

with F as in (10.2.46). Define

$$A := \bigcup_{j \in \mathcal{J}_2(x)} B_{3,j}.$$

We claim that

$$A \subset B(x, 3r) \setminus B(x, \frac{r}{3}). \quad (10.2.50)$$

Indeed, for each $j \in \mathcal{J}_2(x)$ and $y \in B_{3,j}$, using again (10.2.44),

$$\rho(x, x_j) < r + r_{3,j} \leq r + \frac{1}{2}\rho(x, x_j) \implies \rho(x, x_j) < 2r$$

and hence

$$\rho(x, y) \leq \rho(x, x_j) + \rho(x_j, y) < 2r + 3r_j \leq 2r + \frac{1}{2}\rho(x, x_j) < 3r.$$

For the lower bound, we observe

$$\rho(x, x_j) \geq r - 3r_j \geq r - \frac{1}{2}\rho(x, x_j) \implies \rho(x, x_j) \geq \frac{2}{3}r,$$

and conclude

$$\rho(x, y) \geq \rho(x, x_j) - \rho(y, x_j) \geq \rho(x, x_j) - 3r_j \geq \rho(x, x_j) - \frac{1}{2}\rho(x, x_j) \geq \frac{1}{3}r.$$

Using the bounded intersection property of the $B_{3,j}$, (10.2.50) and (1.1.11), we get

$$\sum_{j \in \mathcal{J}_2(x)} \int_{B_{3,j}} |K(x, y)| d\mu(y) \leq 2^{6a} \int_A |K(x, y)| d\mu(y) \quad (10.2.51)$$

$$\leq 2^{6a} \int_{B(x, 3r) \setminus B(x, \frac{r}{3})} |K(x, y)| d\mu(y) \quad (10.2.52)$$

$$\leq 2^{6a} \int_{B(x, 3r) \setminus B(x, \frac{r}{3})} \frac{2^{a^3}}{V(x, y)} d\mu(y) \quad (10.2.53)$$

$$\leq 2^{a^3+6a} \int_{B(x, 3r) \setminus B(x, \frac{r}{3})} \frac{1}{\mu(B(x, \frac{r}{3}))} d\mu(y) \quad (10.2.54)$$

$$\leq 2^{a^3+6a} \frac{\mu(B(x, 3r))}{\mu(B(x, \frac{r}{3}))} \quad (10.2.55)$$

$$\leq 2^{a^3+10a}. \quad (10.2.56)$$

Combining the estimates (10.2.46) for (10.2.41), (10.2.49) for (10.2.42), and (10.2.56), we get

$$|T_r b(x)| \leq 3F(x) + 2^{a^3+12a+1} c\alpha.$$

By the definition (10.2.35) of c , this equals

$$3F(x) + \alpha/8.$$

□

Lemma 10.2.8 (Estimate F set). *For F as defined in Lemma 10.2.7, we have*

$$\mu(\{x \in X \setminus \Omega : F(x) > \alpha/8\}) \leq \frac{2^{a^3+9a+4}}{\alpha} \int |f(y)| d\mu(y). \quad (10.2.57)$$

Proof. We estimate

$$\mu(\{x \in X \setminus \Omega : F(x) > \alpha/8\}) \leq \frac{8}{\alpha} \int_{X \setminus \Omega} F(x) d\mu(x) \quad (10.2.58)$$

$$\leq \frac{8}{\alpha} \int_{X \setminus \Omega} 2^{a^3+2a+1} c \alpha \sum_j \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{\mu(B_{3,j})}{V(x, x_j)} d\mu(x) \quad (10.2.59)$$

$$\leq 2^{a^3+2a+4} c \sum_j \mu(B_{3,j}) \int_{X \setminus B_{6,j}} \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{1}{V(x, x_j)} d\mu(x) \quad (10.2.60)$$

Using

$$V(x, x_j) = \mu(B(x, \rho(x, x_j))) \geq 2^{-a} \mu(B(x, 2\rho(x, x_j))) \geq 2^{-a} \mu(B(x_j, \rho(x_j, x))),$$

we have for all $j \in J$,

$$\begin{aligned} & \int_{X \setminus B_{6,j}} \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{1}{V(x, x_j)} d\mu(x) \\ & \leq 2^a \int_{X \setminus B_{6,j}} \left(\frac{3r_j}{\rho(x, x_j)} \right)^{\frac{1}{a}} \frac{1}{\mu(B(x_j, \rho(x_j, x)))} d\mu(x) \\ & \leq 2^a \sum_{n=1}^{\infty} \int_{B(x_j, 2^{n+1} \cdot 3r_j) \setminus B(x_j, 2^n \cdot 3r_j)} \left(\frac{3r_j}{2^n \cdot 3r_j} \right)^{\frac{1}{a}} \frac{1}{\mu(B(x_j, 2^n \cdot 3r_j))} d\mu(x) \\ & \leq 2^a \sum_{n=1}^{\infty} 2^{-\frac{n}{a}} \frac{\mu(B(x_j, 2^{n+1} \cdot 3r_j))}{\mu(B(x_j, 2^n \cdot 3r_j))} \\ & \leq 2^{3a}, \end{aligned}$$

where we used Lemma 10.1.1 in the last step. Plugging this into (10.2.60) and using (10.2.22), we conclude that

$$\mu(\{x \in X \setminus \Omega : F(x) > \alpha/8\}) \leq \frac{2^{a^3+9a+4}}{\alpha} \int |f(y)| d\mu(y).$$

□

Lemma 10.2.9 (Estimate bad). *We have*

$$\mu(\{x \in X : |T_r b(x)| > \alpha/2\}) \leq \frac{\frac{2^{5a}}{c} + 2^{a^3+9a+4}}{\alpha} \int |f(y)| d\mu(y).$$

Proof. We estimate

$$\begin{aligned} & \mu(\{x \in X : |T_r b(x)| > \alpha/2\}) \\ & \leq \mu(\Omega) + \mu(\{x \in X \setminus \Omega : |T_r b(x)| > \alpha/2\}). \end{aligned} \quad (10.2.61)$$

Using (10.2.38) and (10.2.22), we conclude that

$$\mu(\Omega) \leq \sum_j \mu(B_{6,j}) \leq 2^a \sum_j \mu(B_{3,j}) \leq \frac{2^{5a}}{c\alpha} \int |f(y)| d\mu(y). \quad (10.2.62)$$

It follows from Lemma 10.2.7 and the triangle inequality that

$$\mu(\{x \in X \setminus \Omega : |T_r b(x)| > \alpha/2\}) \leq \mu(\{x \in X \setminus \Omega : F(x) > \alpha/8\}). \quad (10.2.63)$$

The claim now follows from Lemma 10.2.8. \square

Proof of Lemma 10.0.3. It follows by the triangle inequality and subadditivity of μ that

$$\mu(\{x \in X : |T_r f(x)| > \alpha\})$$

$$\leq \mu(\{x \in X : |T_r g(x)| > \alpha/2\}) + \mu(\{x \in X : |T_r b(x)| > \alpha/2\}).$$

Using Lemma 10.2.6, Lemma 10.2.9 and the definition (10.2.35) of c , we get

$$\begin{aligned} &\leq \frac{2^{2a^3+3a+2}c + \frac{2^{5a}}{c} + 2^{a^3+9a+4}}{\alpha} \int |f(y)| d\mu(y) \\ &= \frac{2^{a^3-9a-2} + 2^{a^3+17a+4} + 2^{a^3+9a+4}}{\alpha} \int |f(y)| d\mu(y) \\ &\leq \frac{2^{a^3+19a}}{\alpha} \int |f(y)| d\mu(y). \end{aligned}$$

\square

11. PROOF OF THE CLASSICAL CARLESON THEOREM

The convergence of partial Fourier sums is proved in Section 11.1 in two steps. In the first step, we establish convergence on a suitable dense subclass of functions. We choose smooth functions as subclass, the convergence is stated in Lemma 11.1.2 and proved in Section 11.2. In the second step, one controls the relevant error of approximating a general function by a function in the subclass. This is stated in Lemma 11.1.3 and proved in Section 11.6. The proof relies on a bound on the real Carleson maximal operator stated in Lemma 11.1.5 and proved in Section 11.7, which involves showing that the real line fits into the setting of Section 2. This latter proof refers to the two-sided variant of the Carleson Theorem 10.0.1. Two assumptions in Theorem 1.1.1 require more work. The boundedness of the operator T_r defined in (10.0.2) is established in 11.1.6. This lemma is proved in Section 11.3. The cancellative property is verified by Lemma 11.1.7, which is proved in Section 11.4. Several further auxiliary lemmas are stated and proved in Section 11.1, the proof of one of these auxiliary lemmas, Lemma 11.1.10, is done in Section 11.5.

All subsections past Section 11.1 are mutually independent.

11.1. The classical Carleson theorem. Let a uniformly continuous 2π -periodic function $f : \mathbb{R} \rightarrow \mathbb{C}$ and $\epsilon > 0$ be given. Let

$$C_{a,q} := \frac{2^{474a^3}}{(q-1)^6} \quad (11.1.1)$$

denote the constant from Theorem 10.0.1. Define

$$\epsilon' := \frac{\epsilon}{4C_\epsilon}, \quad (11.1.2)$$

where

$$C_\epsilon = \left(\frac{8}{\pi\epsilon}\right)^{\frac{1}{2}} C_{4,2} + \pi.$$

Since f is continuous and periodic, f is uniformly continuous. Thus, there is a $0 < \delta < \pi$ such that for all $x, x' \in \mathbb{R}$ with $|x - x'| \leq \delta$ we have

$$|f(x) - f(x')| \leq \epsilon'. \quad (11.1.3)$$

Define

$$f_0 := f * \phi_\delta, \quad (11.1.4)$$

where ϕ_δ is a nonnegative smooth bump function with $\text{supp}(\phi_\delta) \subset (-\delta, \delta)$ and $\int_{\mathbb{R}} \phi_\delta(x) dx = 1$.

Lemma 11.1.1 (smooth approximation). *The function f_0 is 2π -periodic. The function f_0 is smooth (and therefore measurable). The function f_0 satisfies for all $x \in \mathbb{R}$:*

$$|f(x) - f_0(x)| \leq \epsilon', \quad (11.1.5)$$

Proof. Periodicity follows directly from the definitions. The other properties are part of the Lean library. \square

We prove in Section 11.2:

Lemma 11.1.2 (convergence for smooth). *There exists some $N_0 \in \mathbb{N}$ such that for all $N > N_0$ and $x \in [0, 2\pi]$ we have*

$$|S_N f_0(x) - f_0(x)| \leq \frac{\epsilon}{4}. \quad (11.1.6)$$

We prove in Section 11.6:

Lemma 11.1.3 (control approximation effect). *There is a set $E \subset \mathbb{R}$ with Lebesgue measure $|E| \leq \epsilon$ such that for all*

$$x \in [0, 2\pi] \setminus E \quad (11.1.7)$$

we have

$$\sup_{N \geq 0} |S_N f(x) - S_N f_0(x)| \leq \frac{\epsilon}{4}. \quad (11.1.8)$$

We are now ready to prove Theorem 1.0.1. We first prove a version with explicit exceptional sets.

Theorem 11.1.4 (classical Carleson with exceptional sets). *Let f be a 2π -periodic complex-valued continuous function on \mathbb{R} . For all $\epsilon > 0$, there exists a Borel set $E \subset [0, 2\pi]$ with Lebesgue measure $|E| \leq \epsilon$ and a positive integer N_0 such that for all $x \in [0, 2\pi] \setminus E$ and all integers $N > N_0$, we have*

$$|f(x) - S_N f(x)| \leq \epsilon. \quad (11.1.9)$$

Proof. Let N_0 be as in Lemma 11.1.2. For every

$$x \in [0, 2\pi] \setminus E, \quad (11.1.10)$$

and every $N > N_0$ we have by the triangle inequality

$$\begin{aligned} & |f(x) - S_N f(x)| \\ & \leq |f(x) - f_0(x)| + |f_0(x) - S_N f_0(x)| + |S_N f_0(x) - S_N f(x)|. \end{aligned} \quad (11.1.11)$$

Using Lemmas 11.1.1 to 11.1.3, we estimate (11.1.11) by

$$\leq \epsilon' + \frac{\epsilon}{4} + \frac{\epsilon}{4} \leq \epsilon. \quad (11.1.12)$$

This shows (11.1.9) for the given E and N_0 . \square

Now we turn to the proof of the statement of Carleson theorem given in the introduction.

Proof of Theorem 1.0.1. By applying Theorem 11.1.4 with a sequence of $\epsilon_n := 2^{-n}\delta$ for $n \geq 1$ and taking the union of corresponding exceptional sets E_n , we see that outside a set of measure δ , the partial Fourier sums converge pointwise for $N \rightarrow \infty$. Applying this with a sequence of δ shrinking to zero and taking the intersection of the corresponding exceptional sets, which has measure zero, we see that the Fourier series converges outside a set of measure zero. \square

Let $\kappa : \mathbb{R} \rightarrow \mathbb{C}$ be the function defined by $\kappa(0) = 0$ and for $0 < |x| < 1$

$$\kappa(x) = \frac{1 - |x|}{1 - e^{ix}} \quad (11.1.13)$$

and for $|x| \geq 1$,

$$\kappa(x) = 0. \quad (11.1.14)$$

Note that this function is continuous at every point x with $|x| > 0$.

The proof of Lemma 11.1.3 will use the following Lemma 11.1.5, which itself is proven in Section 11.7 as an application of Theorem 1.1.1.

Lemma 11.1.5 (real Carleson). *Let F, G be Borel subsets of \mathbb{R} with finite measure. Let f be a bounded measurable function on \mathbb{R} with $|f| \leq \mathbf{1}_F$. Then*

$$\left| \int_G T f(x) dx \right| \leq C_{4,2} |F|^{\frac{1}{2}} |G|^{\frac{1}{2}}, \quad (11.1.15)$$

where

$$T f(x) = \sup_{n \in \mathbb{Z}} \sup_{r > 0} \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right|. \quad (11.1.16)$$

One of the main assumptions of Theorem 10.0.1, concerning the operator T_r defined in (10.0.2), is verified by the following lemma, which is proved in Section 11.3.

Lemma 11.1.6 (Hilbert strong 2 2). *Let $0 < r$. Let f be a bounded, measurable function on \mathbb{R} . Then*

$$\|H_r f\|_2 \leq 2^9 \|f\|_2, \quad (11.1.17)$$

where

$$H_r f(x) := T_r f(x) = \int_{r \leq |x-y|} \kappa(x-y) f(y) dy \quad (11.1.18)$$

The next lemma will be used to verify that the collection Θ of modulation functions in our application of Theorem 1.1.1 satisfies the condition (1.1.9). It is proved in Section 11.4.

Lemma 11.1.7 (van der Corput). *Let $\alpha \leq \beta$ be real numbers. Let $g : \mathbb{R} \rightarrow \mathbb{C}$ be a measurable function and assume*

$$\|g\|_{Lip(\alpha, \beta)} := \sup_{\alpha \leq x \leq \beta} |g(x)| + \frac{|\beta - \alpha|}{2} \sup_{\alpha \leq x < y \leq \beta} \frac{|g(y) - g(x)|}{|y - x|} < \infty. \quad (11.1.19)$$

Then for any $\alpha \leq \beta$ and $n \in \mathbb{Z}$ we have

$$\int_{\alpha}^{\beta} g(x) e^{inx} dx \leq 2\pi |\beta - \alpha| \|g\|_{Lip(\alpha, \beta)} (1 + |n| |\beta - \alpha|)^{-1}. \quad (11.1.20)$$

We close this section with five lemmas that are used across the following subsections.

Lemma 11.1.8 (Dirichlet kernel). *We have for every 2π -periodic bounded measurable f and every $N \geq 0$*

$$S_N f(x) = \frac{1}{2\pi} \int_0^{2\pi} f(y) K_N(x-y) dy \quad (11.1.21)$$

where K_N is the 2π -periodic continuous function of \mathbb{R} given by

$$\sum_{n=-N}^N e^{inx'}. \quad (11.1.22)$$

We have for $e^{ix'} \neq 1$ that

$$K_N(x') = \frac{e^{iNx'}}{1 - e^{-ix'}} + \frac{e^{-iNx'}}{1 - e^{ix'}}. \quad (11.1.23)$$

Proof. We have by definitions and interchanging sum and integral

$$\begin{aligned} S_N f(x) &= \sum_{n=-N}^N \hat{f}_n e^{inx} \\ &= \sum_{n=-N}^N \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{in(x-y)} dy \end{aligned}$$

$$= \frac{1}{2\pi} \int_0^{2\pi} f(y) \sum_{n=-N}^N e^{in(x-y)} dy. \quad (11.1.24)$$

This proves the first statement of the lemma. By a telescoping sum, we have for every $x' \in \mathbb{R}$

$$\left(e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'} \right) \sum_{n=-N}^N e^{inx'} = e^{(N+\frac{1}{2})ix'} - e^{-(N+\frac{1}{2})ix'}. \quad (11.1.25)$$

If $e^{ix'} \neq 1$, the first factor on the left-hand side is not 0 and we may divide by this factor to obtain

$$\sum_{n=-N}^N e^{inx'} = \frac{e^{i(N+\frac{1}{2})x'}}{e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'}} - \frac{e^{-i(N+\frac{1}{2})x'}}{e^{\frac{1}{2}ix'} - e^{-\frac{1}{2}ix'}} = \frac{e^{iNx'}}{1 - e^{-ix'}} + \frac{e^{-iNx'}}{1 - e^{ix'}}. \quad (11.1.26)$$

This proves the second part of the lemma. \square

Lemma 11.1.9 (lower secant bound). *Let $\eta > 0$ and $-2\pi + \eta \leq x \leq 2\pi - \eta$ with $|x| \geq \eta$. Then*

$$|1 - e^{ix}| \geq \frac{2}{\pi}\eta \quad (11.1.27)$$

Proof. We have

$$|1 - e^{ix}| = \sqrt{(1 - \cos(x))^2 + \sin^2(x)} \geq |\sin(x)|.$$

If $0 \leq x \leq \frac{\pi}{2}$, then we have from concavity of \sin on $[0, \pi]$ and $\sin(0) = 0$ and $\sin(\frac{\pi}{2}) = 1$

$$|\sin(x)| \geq \frac{2}{\pi}x \geq \frac{2}{\pi}\eta.$$

When $x \in \frac{m\pi}{2} + [0, \frac{\pi}{2}]$ for $m \in \{-4, -3, -2, -1, 1, 2, 3\}$ one can argue similarly. \square

The following lemma will be proved in Section 11.5.

Lemma 11.1.10 (spectral projection bound). *Let f be a bounded 2π -periodic measurable function. Then, for all $N \geq 0$*

$$\|S_N f\|_{L^2[0, 2\pi]} \leq \|f\|_{L^2[0, 2\pi]}. \quad (11.1.28)$$

Lemma 11.1.11 (Hilbert kernel bound). *For $x, y \in \mathbb{R}$ with $x \neq y$ we have*

$$|\kappa(x - y)| \leq 2^2(2|x - y|)^{-1}. \quad (11.1.29)$$

Proof. Fix $x \neq y$. If $\kappa(x - y)$ is zero, then (11.1.29) is evident. Assume $\kappa(x - y)$ is not zero, then $0 < |x - y| < 1$. We have

$$|\kappa(x - y)| = \left| \frac{1 - |x - y|}{1 - e^{i(x-y)}} \right|. \quad (11.1.30)$$

We estimate with Lemma 11.1.9

$$|\kappa(x - y)| \leq \frac{1}{|1 - e^{i(x-y)}|} \leq \frac{2}{|x - y|}. \quad (11.1.31)$$

This proves (11.1.29) in the given case and completes the proof of the lemma. \square

Lemma 11.1.12 (Hilbert kernel regularity). *For $x, y, y' \in \mathbb{R}$ with $x \neq y, y'$ and*

$$2|y - y'| \leq |x - y|, \quad (11.1.32)$$

we have

$$|\kappa(x - y) - \kappa(x - y')| \leq 2^8 \frac{1}{|x - y|} \frac{|y - y'|}{|x - y|}. \quad (11.1.33)$$

Proof. Upon replacing y by $y - x$ and y' by $y' - x$ on the left-hand side of (11.1.32), we can assume that $x = 0$. Then the assumption (11.1.32) implies that y and y' have the same sign. Since $\kappa(y) = \bar{\kappa}(-y)$ we can assume that they are both positive. Then it follows from (11.1.32) that

$$\frac{y}{2} \leq y'.$$

We distinguish four cases. If $y, y' \leq 1$, then we have

$$|\kappa(-y) - \kappa(-y')| = \left| \frac{1 - y}{1 - e^{-iy}} - \frac{1 - y'}{1 - e^{-iy'}} \right|$$

and by the fundamental theorem of calculus

$$= \left| \int_{y'}^y \frac{-1 + e^{-it} + i(1 - t)e^{it}}{(1 - e^{-it})^2} dt \right|.$$

Using $y' \geq \frac{y}{2}$ and Lemma 11.1.9, we bound this by

$$\leq |y - y'| \sup_{\frac{y}{2} \leq t \leq 1} \frac{3}{|1 - e^{-it}|^2} \leq 3|y - y'| \left(2\frac{2}{y}\right)^2 \leq 2^6 \frac{|y - y'|}{|y|^2}.$$

If $y \leq 1$ and $y' > 1$, then $\kappa(-y') = 0$ and we have from the first case

$$|\kappa(-y) - \kappa(-y')| = |\kappa(-y) - \kappa(-1)| \leq 2^6 \frac{|y - 1|}{|y|^2} \leq 2^6 \frac{|y - y'|}{|y|^2}.$$

Similarly, if $y > 1$ and $y' \leq 1$, then $\kappa(-y) = 0$ and we have from the first case

$$|\kappa(-y) - \kappa(-y')| = |\kappa(-y') - \kappa(-1)| \leq 2^6 \frac{|y' - 1|}{|y'|^2} \leq 2^6 \frac{|y - y'|}{|y'|^2}.$$

Using again $y' \geq \frac{y}{2}$, we bound this by

$$\leq 2^6 \frac{|y - y'|}{|y/2|^2} = 2^8 \frac{|y - y'|}{|y|^2}$$

Finally, if $y, y' > 1$ then

$$|\kappa(-y) - \kappa(-y')| = 0 \leq 2^8 \frac{|y - y'|}{|y|^2}.$$

□

11.2. Smooth functions.

Lemma 11.2.1. *Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be 2π -periodic and continuously differentiable, and let $n \in \mathbb{Z} \setminus \{0\}$. Then*

$$\widehat{f}_n = \frac{1}{in} \widehat{f}'_n. \quad (11.2.1)$$

Proof. This is part of the Lean library. \square

Lemma 11.2.2. *Let $f : \mathbb{R} \rightarrow \mathbb{C}$ such that*

$$\sum_{n \in \mathbb{Z}} |\widehat{f}_n| < \infty. \quad (11.2.2)$$

Then

$$\sup_{x \in [0, 2\pi]} |f(x) - S_N f(x)| \rightarrow 0 \quad (11.2.3)$$

as $N \rightarrow \infty$.

Proof. This is part of the Lean library. \square

Lemma 11.2.3. *Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be 2π -periodic and twice continuously differentiable. Then*

$$\sup_{x \in [0, 2\pi]} |f(x) - S_N f(x)| \rightarrow 0 \quad (11.2.4)$$

as $N \rightarrow \infty$.

Proof. By Lemma 11.2.2, it suffices to show that the Fourier coefficients \widehat{f}_n are summable. Applying Lemma 11.2.1 twice and using the fact that f'' is continuous and thus bounded on $[0, 2\pi]$, we compute

$$\sum_{n \in \mathbb{Z}} |\widehat{f}_n| = |\widehat{f}_0| + \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{n^2} |\widehat{f}''_n| \leq |\widehat{f}_0| + \left(\sup_{x \in [0, 2\pi]} |f(x)| \right) \cdot \sum_{n \in \mathbb{Z} \setminus \{0\}} \frac{1}{n^2} < \infty.$$

\square

Proof of Lemma 11.1.2. Lemma 11.1.2 now follows directly from the previous Lemma 11.2.3. \square

11.3. The truncated Hilbert transform. Let M_n be the modulation operator acting on measurable 2π -periodic functions defined by

$$M_n g(x) = g(x) e^{inx}. \quad (11.3.1)$$

Define the approximate Hilbert transform by

$$L_N g = \frac{1}{N} \sum_{n=0}^{N-1} M_{n+N} S_{N+n} M_{-N-n} g. \quad (11.3.2)$$

Lemma 11.3.1 (modulated averaged projection). *We have for every bounded measurable 2π -periodic function g*

$$\|L_N g\|_{L^2[0, 2\pi]} \leq \|g\|_{L^2[0, 2\pi]}. \quad (11.3.3)$$

Proof. We have

$$\|M_n g\|_{L^2[0,2\pi]}^2 = \int_0^{2\pi} |e^{inx} g(x)|^2 dx = \int_0^{2\pi} |g(x)|^2 dx = \|g\|_{L^2[0,2\pi]}^2. \quad (11.3.4)$$

We have by the triangle inequality, the square root of the identity in (11.3.4), and Lemma 11.1.10

$$\begin{aligned} \|L_n g\|_{L^2[0,2\pi]} &= \left\| \frac{1}{N} \sum_{n=0}^{N-1} M_{n+N} S_{N+n} M_{-N-n} g \right\|_{L^2[0,2\pi]} \\ &\leq \frac{1}{N} \sum_{n=0}^{N-1} \|M_{n+N} S_{N+n} M_{-N-n} g\|_{L^2[0,2\pi]} = \frac{1}{N} \sum_{n=0}^{N-1} \|S_{N+n} M_{-N-n} g\|_{L^2[0,2\pi]} \\ &\leq \frac{1}{N} \sum_{n=0}^{N-1} \|M_{-N-n} g\|_{L^2[0,2\pi]} = \frac{1}{N} \sum_{n=0}^{N-1} \|g\|_{L^2[0,2\pi]} = \|g\|_{L^2[0,2\pi]}. \end{aligned} \quad (11.3.5)$$

This proves (11.3.3) and completes the proof of the lemma. \square

Lemma 11.3.2 (periodic domain shift). *Let f be a bounded 2π -periodic function. We have for any $0 \leq x \leq 2\pi$ that*

$$\int_0^{2\pi} f(y) dy = \int_{-x}^{2\pi-x} f(y) dy = \int_{-\pi}^{\pi} f(y-x) dy. \quad (11.3.6)$$

Proof. We have by periodicity and change of variables

$$\int_{-x}^0 f(y) dy = \int_{-x}^0 f(y+2\pi) dy = \int_{2\pi-x}^{2\pi} f(y) dy. \quad (11.3.7)$$

We then have by breaking up the domain of integration and using (11.3.7)

$$\begin{aligned} \int_0^{2\pi} f(y) dy &= \int_0^{2\pi-x} f(y) dy + \int_{2\pi-x}^{2\pi} f(y) dy \\ &= \int_0^{2\pi-x} f(y) dy + \int_{-x}^0 f(y) dy = \int_{-x}^{2\pi-x} f(y) dy. \end{aligned} \quad (11.3.8)$$

This proves the first identity of the lemma. The second identity follows by substitution of y by $y-x$. \square

Lemma 11.3.3 (Young convolution). *Let f and g be two bounded non-negative measurable 2π -periodic functions on \mathbb{R} . Then*

$$\left(\int_0^{2\pi} \left(\int_0^{2\pi} f(y) g(x-y) dy \right)^2 dx \right)^{\frac{1}{2}} \leq \|f\|_{L^2[0,2\pi]} \|g\|_{L^1[0,2\pi]}. \quad (11.3.9)$$

Proof. Using Fubini and Lemma 11.3.2, we observe

$$\begin{aligned} \int_0^{2\pi} \int_0^{2\pi} f(y)^2 g(x-y) dy dx &= \int_0^{2\pi} f(y)^2 \int_0^{2\pi} g(x-y) dx dy \\ &= \int_0^{2\pi} f(y)^2 \int_0^{2\pi} g(x) dx dy = \|f\|_{L^2[0,2\pi]}^2 \|g\|_{L^1[0,2\pi]}. \end{aligned} \quad (11.3.10)$$

Let h be the nonnegative square root of g , then h is bounded and 2π -periodic with $h^2 = g$. We estimate the square of the left-hand side of (11.3.9) with Cauchy-Schwarz and then with (11.3.10) by

$$\begin{aligned} & \int_0^{2\pi} \left(\int_0^{2\pi} f(y)h(x-y)h(x-y) dy \right)^2 dx \\ & \leq \int_0^{2\pi} \left(\int_0^{2\pi} f(y)^2 g(x-y) dy \right) \left(\int_0^{2\pi} g(x-y) dy \right) dx \\ & = \|f\|_{L^2[0,2\pi]}^2 \|g\|_{L^1[0,2\pi]}^2. \end{aligned}$$

Taking square roots, this proves the lemma. \square

For $0 < r < 1$, Define the kernel k_r to be the 2π -periodic function

$$k_r(x) := \min \left(r^{-1}, 1 + \frac{r}{|1 - e^{ix}|^2} \right), \quad (11.3.11)$$

where the minimum is understood to be r^{-1} in case $1 = e^{ix}$.

Lemma 11.3.4 (integrable bump convolution). *Let g, f be bounded measurable 2π -periodic functions. Let $0 < r < \pi$. Assume we have for all x*

$$|g(x)| \leq k_r(x). \quad (11.3.12)$$

Let

$$h(x) = \int_0^{2\pi} f(y)g(x-y) dy. \quad (11.3.13)$$

Then

$$\|h\|_{L^2[0,2\pi]} \leq 17\|f\|_{L^2[-\pi,\pi]}. \quad (11.3.14)$$

Proof. From monotonicity of the integral and (11.3.12),

$$\|g\|_{L^1[0,2\pi]} \leq \int_0^{2\pi} k_r(x) dx. \quad (11.3.15)$$

Using the symmetry $k_r(x) = k_r(-x)$, the assumption, and Lemma 11.1.9, the last display is equal to

$$\begin{aligned} & = 2 \int_0^\pi \min \left(\frac{1}{r}, 1 + \frac{r}{|1 - e^{ix}|^2} \right) dx \\ & \leq 2 \int_0^r \frac{1}{r} dx + 2 \int_r^\pi \left(1 + \frac{64r}{x^2} \right) dx \\ & \leq 2 + 2\pi + 2 \left(\frac{4r}{r} - \frac{4r}{\pi} \right) \leq 17. \end{aligned} \quad (11.3.16)$$

Together with Lemma 11.3.3, this proves the lemma. \square

Lemma 11.3.5 (Dirichlet approximation). *Let $0 < r < 1$. Let N be the smallest integer larger than $\frac{1}{r}$. There is a 2π -periodic continuous function L' on \mathbb{R} that satisfies for all $0 \leq x \leq 2\pi$ and all 2π -periodic bounded measurable functions f on \mathbb{R}*

$$L_N f(x) = \frac{1}{2\pi} \int_0^{2\pi} f(y) L'(x-y) dy. \quad (11.3.17)$$

Moreover, for all $-\pi \leq x \leq \pi$,

$$\left| L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}} \kappa(x) \right| \leq 12k_r(x). \quad (11.3.18)$$

Proof. We have by definition and Lemma 11.1.8

$$L_N g(x) = \frac{1}{N} \sum_{n=0}^{N-1} \int_0^{2\pi} e^{-i(N+n)x} K_{N+n}(x-y) e^{i(N+n)y} g(y) dy. \quad (11.3.19)$$

This is of the form (11.3.17) with the continuous function

$$L'(x) = \frac{1}{N} \sum_{n=0}^{N-1} K_{N+n}(x) e^{-i(N+n)x}. \quad (11.3.20)$$

With (11.1.22) of Lemma 11.1.8 we have $|K_N(x)| \leq 2N + 1$ for every x and thus

$$|L'(x)| \leq \frac{1}{N} \sum_{n=0}^{N-1} (2N + 2n + 1) \leq 4N \leq 2^3 r^{-1}. \quad (11.3.21)$$

Therefore, for $|x| \in [0, r) \cup (1, \pi]$, we have

$$\left| L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}}(x) \kappa(x) \right| = |L'(x)| \leq 2^3 r^{-1}. \quad (11.3.22)$$

This proves (11.3.18) for $|x| \in [0, r)$ since $k_r(x) = r^{-1}$ in this case.

For $e^{ix} \neq 1$ one may use the expression (11.1.23) for K_N in Lemma 11.1.8 to obtain

$$\begin{aligned} L'(x) &= \frac{1}{N} \sum_{n=0}^{N-1} \left(\frac{e^{i(N+n)x}}{1 - e^{-ix}} + \frac{e^{-i(N+n)x}}{1 - e^{ix}} \right) e^{i(N+n)x} \\ &= \frac{1}{N} \sum_{n=0}^{N-1} \left(\frac{e^{i2(N+n)x}}{1 - e^{-ix}} + \frac{1}{1 - e^{ix}} \right) \\ &= \frac{1}{1 - e^{ix}} + \frac{1}{N} \frac{e^{i2Nx}}{1 - e^{-ix}} \sum_{n=0}^{N-1} e^{i2nx} \end{aligned} \quad (11.3.23)$$

and thus

$$L'(x) - \mathbf{1}_{\{y: r < |y| < 1\}} \kappa(x) = L''(x) + \frac{1 - \mathbf{1}_{\{y: r < |y| < 1\}}(x)(1 - |x|)}{1 - e^{ix}}, \quad (11.3.24)$$

where

$$L''(x) := \frac{1}{N} \frac{e^{i2Nx}}{1 - e^{-ix}} \sum_{n=0}^{N-1} e^{i2nx}.$$

For $x \in [-\pi, r] \cup [r, \pi]$, we have using Lemma 11.1.9 that

$$\left| \frac{1 - \mathbf{1}_{\{y: r < |y| < 1\}}(x)(1 - |x|)}{1 - e^{ix}} \right| = \left| \frac{\min(|x|, 1)}{1 - e^{ix}} \right| \leq \frac{2 \min(|x|, 1)}{|x|}$$

$$\leq 2 \cdot 1 \leq 2k_r(x). \quad (11.3.25)$$

Next, we need to estimate $L''(x)$. If the real part of e^{ix} is negative, we have

$$1 \leq |1 - e^{-ix}| \leq 2. \quad (11.3.26)$$

and hence

$$|L''(x)| \leq \frac{1}{N} \sum_{n=0}^{N-1} 1 = 1 \leq 1 + \frac{r}{|1 - e^{ix}|^2}. \quad (11.3.27)$$

If the real part of e^{ix} is positive and in particular while still $e^{ix} \neq \pm 1$, then we have by telescoping

$$(1 - e^{2ix}) \sum_{n=0}^{N-1} e^{i2nx} = 1 - e^{i2Nx}. \quad (11.3.28)$$

As $e^{2ix} \neq 1$, we may divide by $1 - e^{2ix}$ and insert this into (11.3.23) to obtain

$$L''(x) = \frac{1}{N} \frac{e^{i2Nx}}{1 - e^{-ix}} \frac{1 - e^{i2Nx}}{1 - e^{2ix}}. \quad (11.3.29)$$

Hence, using the nonnegativity of the real part of e^{ix}

$$\begin{aligned} |L''(x)| &\leq \frac{2}{N} \frac{1}{|1 - e^{ix}|} \frac{1}{|1 - e^{2ix}|} \\ &= \frac{2}{N} \frac{1}{|1 - e^{ix}|^2} \frac{1}{|1 + e^{ix}|} \leq \frac{2r}{|1 - e^{ix}|^2} \leq 2 \left(1 + \frac{r}{|1 - e^{ix}|^2} \right) \end{aligned} \quad (11.3.30)$$

Moreover, for $|x| \in [r, \pi]$, one checks easily that

$$1 + \frac{r}{|1 - e^{ix}|^2} \leq 5k_r(x).$$

Therefore, (11.3.27) and (11.3.30) show that, in this range of x ,

$$|L''(x)| \leq 10k_r(x).$$

Together with (11.3.25), this prove (11.3.18) for $|x| \in [r, \pi]$. As the range $|x| \in [0, r]$ is covered by (11.3.22), this completes the proof of the lemma. \square

We now prove Lemma 11.1.6.

Proof of Lemma 11.1.6. We first show that if f is supported in $[1, 4]$, then

$$\|H_r f\|_{L^2[2,3]} \leq 2^8 \|f\|_{L^2(\mathbb{R})}. \quad (11.3.31)$$

Let \tilde{f} be the 2π -periodic extension of f to \mathbb{R} . Let N be the smallest integer larger than $\frac{1}{r}$. Then, Equation (11.3.18) shows that the kernels of H_r and $2\pi L_N$ differ by at most $12k_r$ on $[-\pi, \pi]$. Consider $x \in [2, 3]$. When computing $H_r f(x)$ and $2\pi L_N f(x)$, the kernels are computed at points of the form $x - y$ with $f(y) \neq 0$, i.e., $y \in [1, 4]$. As $x \in [2, 3]$, the difference $x - y$ is bounded in absolute value by 2, and therefore belongs to $[-\pi, \pi]$, where the above bound holds.

It follows that, for $x \in [2, 3]$,

$$|H_r \tilde{f}(x)| \leq 2\pi |L_N \tilde{f}(x)| + 12 \left| \int_0^{2\pi} k_r(x - y) \tilde{f}(y) dy \right|.$$

Taking L^2 norm and using its sub-additivity, we get

$$\|H_r \tilde{f}\|_{L^2[2,3]} \leq 2\pi \|L_N \tilde{f}\|_{L^2[0,2\pi]} + 12 \left(\int_0^{2\pi} \left| \int_0^{2\pi} k_r(x-y) \tilde{f}(y) dy \right|^2 dx \right)^{\frac{1}{2}}.$$

Since κ is supported in $[-1, 1]$, we have that $H_r \tilde{f}$ agrees with $H_r f$ on $[2, 3]$. Using Lemma 11.3.1 and Lemma 11.3.4, we conclude

$$\|H_r f\|_{L^2[2,3]} \leq 2\pi \|f\|_{L^2[0,2\pi]} + 12 \cdot 17 \|f\|_{L^2[0,2\pi]}, \quad (11.3.32)$$

which gives (11.3.31).

Suppose now that f is supported in $[c, c+3]$ for some $c \in \mathbb{R}$. Then the function $g(x) = f(x-c+1)$ is supported in $[1, 4]$. By a change of variables in (11.1.18), we have $H_r g(x) = H_r f(x-c+1)$. Thus, by (11.3.31)

$$\|H_r f\|_{L^2[c+1, c+2]} = \|H_r g\|_{L^2[2,3]} \leq 2^8 \|g\|_2 = 2^8 \|f\|_2. \quad (11.3.33)$$

Let now f be arbitrary. Since $\kappa(x) = 0$ for $|x| > 1$, we have for all $x \in [c+1, c+2]$

$$H_r f(x) = H_r (f \mathbf{1}_{[c, c+3]})(x).$$

Thus

$$\int_{c+1}^{c+2} |H_r f(x)|^2 dx = \int_{c+1}^{c+2} |H_r (f \mathbf{1}_{[c, c+3]})(x)|^2 dx.$$

Applying the bound (11.3.33), this is

$$\leq 2^{16} \int_c^{c+3} |f(x)|^2 dx.$$

Summing over all $c \in \mathbb{Z}$, we obtain

$$\int_{\mathbb{R}} |H_r f(x)|^2 dx \leq 3 \cdot 2^{16} \int_{\mathbb{R}} |f(x)|^2 dx.$$

This completes the proof. \square

11.4. The proof of the van der Corput Lemma.

Proof of Lemma 11.1.7. Let g be a Lipschitz continuous function as in the lemma. Assume first that $n = 0$. Then

$$\int_{\alpha}^{\beta} g(x) dx \leq |\beta - \alpha| \sup_{\alpha \leq x \leq \beta} |g(x)| \leq |\beta - \alpha| \|g\|_{Lip(\alpha, \beta)} (1 + |n| |\beta - \alpha|)^{-1}$$

Assume now $n \neq 0$. Without loss of generality, we may assume $n > 0$. We distinguish two cases. If $\beta - \alpha < \frac{\pi}{n}$, we have by the triangle inequality

$$\left| \int_{\alpha}^{\beta} g(x) e^{inx} dx \right| \leq |\beta - \alpha| \sup_{x \in [\alpha, \beta]} |g(x)| \leq 2\pi |\beta - \alpha| \|g\|_{Lip(\alpha, \beta)} (1 + |n| |\beta - \alpha|)^{-1}.$$

We turn to the case $\frac{\pi}{n} \leq \beta - \alpha$. We have

$$e^{in(x+\pi/n)} = -e^{inx}.$$

Using this, we write

$$\int_{\alpha}^{\beta} g(x)e^{inx} dx = \frac{1}{2} \int_{\alpha}^{\beta} g(x)e^{inx} dx - \frac{1}{2} \int_{\alpha}^{\beta} g(x)e^{in(x+\pi/n)} dx.$$

We split the first integral at $\alpha + \frac{\pi}{n}$ and the second one at $\beta - \frac{\pi}{n}$, and make a change of variables in the second part of the first integral to obtain

$$\begin{aligned} &= \frac{1}{2} \int_{\alpha}^{\alpha+\frac{\pi}{n}} g(x)e^{inx} dx - \frac{1}{2} \int_{\beta-\frac{\pi}{n}}^{\beta} g(x)e^{in(x+\pi/n)} dx \\ &\quad + \frac{1}{2} \int_{\alpha+\frac{\pi}{n}}^{\beta} (g(x) - g(x - \frac{\pi}{n}))e^{inx} dx. \end{aligned}$$

The sum of the first two terms is by the triangle inequality bounded by

$$\frac{\pi}{n} \sup_{x \in [\alpha, \beta]} |g(x)|.$$

The third term is by the triangle inequality at most

$$\begin{aligned} &\frac{1}{2} \int_{\alpha+\frac{\pi}{n}}^{\beta} |g(x) - g(x - \frac{\pi}{n})| dx \\ &\leq \frac{|\beta - \alpha|}{2} \frac{\pi}{n} \sup_{\alpha \leq x < y \leq \beta} \frac{|g(x) - g(y)|}{|x - y|}. \end{aligned}$$

Adding the two terms, we obtain

$$\left| \int_{\alpha}^{\beta} g(x)e^{-inx} dx \right| \leq \frac{\pi}{n} \|g\|_{\text{Lip}(\alpha, \beta)}.$$

This completes the proof of the lemma, using that with $\frac{\pi}{n} \leq \beta - \alpha$,

$$\frac{\pi}{n} = \frac{2\pi|\beta - \alpha|}{2n|\beta - \alpha|} \leq 2\pi|\beta - \alpha|(1 + n|\beta - \alpha|)^{-1}.$$

□

11.5. Partial sums as orthogonal projections. This subsection proves Lemma 11.1.10

Proof of Lemma 11.1.10. The functions $e_n : x \mapsto e^{inx}$ form an orthonormal basis in $L^2[-\pi, \pi]$ (this is already in Mathlib). Therefore we have

$$\begin{aligned} \|S_N f\|_{L^2[-\pi, \pi]}^2 &= \left\| \sum_{n=-N}^N \langle \hat{f}_n, e_n \rangle_{L^2[-\pi, \pi]} \right\|_{L^2[-\pi, \pi]}^2 \\ &= \sum_{n=-N}^N |\hat{f}_n|^2 \\ &\leq \sum_{n \in \mathbb{Z}} |\hat{f}_n|^2 \\ &= \|f\|_{L^2[-\pi, \pi]}^2. \end{aligned}$$

This completes the proof of the lemma. □

11.6. The error bound.

Lemma 11.6.1 (Dirichlet kernel - Hilbert kernel relation). *For all $N \in \mathbb{Z}$ and $x \in [-\pi, \pi] \setminus \{0\}$,*

$$\left| K_N(x) - (e^{-iNx} \kappa(x) + \overline{e^{-iNx} \kappa(x)}) \right| \leq \pi.$$

Proof. Let $N \in \mathbb{Z}$ and $x \in [-\pi, \pi] \setminus \{0\}$. With Lemma 11.1.8, we obtain

$$K_N(x) - (e^{-iNx} \kappa(x) + \overline{e^{-iNx} \kappa(x)}) = e^{-iNx} \frac{\min(|x|, 1)}{1 - e^{ix}} + e^{iNx} \frac{\min(|x|, 1)}{1 - e^{-ix}}.$$

Using Lemma 11.1.9 with $\eta = \min(|x|, 1)$, we bound

$$\left| K_N(x) - (e^{-iNx} \kappa(x) + \overline{e^{-iNx} \kappa(x)}) \right| \leq \frac{\min(|x|, 1)}{|1 - e^{ix}|} + \frac{\min(|x|, 1)}{|1 - e^{-ix}|} \leq \frac{\pi}{2} + \frac{\pi}{2} = \pi.$$

□

Lemma 11.6.2 (partial Fourier sum bound). *Let $g : \mathbb{R} \rightarrow \mathbb{C}$ be a measurable 2π -periodic function such that for some $\delta > 0$ and every $x \in \mathbb{R}$,*

$$|g(x)| \leq \delta. \quad (11.6.1)$$

Then for every $x \in [0, 2\pi]$ and $N > 0$,

$$|S_N g(x)| \leq \frac{1}{2\pi} (Tg(x) + T\bar{g}(x)) + \pi\delta.$$

Proof. Let $x \in [0, 2\pi]$ and $N > 0$. We have with Lemma 11.1.8

$$|S_N g(x)| = \frac{1}{2\pi} \left| \int_0^{2\pi} g(y) K_N(x - y) dy \right|.$$

We use 2π -periodicity of g and K_N to shift the domain of integration to obtain

$$= \frac{1}{2\pi} \left| \int_{x-\pi}^{x+\pi} g(y) K_N(x - y) dy \right|.$$

Using the triangle inequality, we split this as

$$\leq \frac{1}{2\pi} \left| \int_{x-\pi}^{x+\pi} g(y) (K_N(x - y) - \max(|x - y|, 0) K_N(x - y)) dy \right| \quad (11.6.2)$$

$$+ \frac{1}{2\pi} \left| \int_{x-\pi}^{x+\pi} g(y) \max(|x - y|, 0) K_N(x - y) dy \right|. \quad (11.6.3)$$

Note that all integrals are well defined, since K_N is by (11.1.22) bounded by $2N + 1$. Using that

$$\max(|x - y|, 0) K_N(x - y) = e^{-iN(x-y)} \kappa(x - y) + \overline{e^{-iN(x-y)} \kappa(x - y)}, \quad (11.6.4)$$

Lemma 11.6.1 and (11.6.5), we bound (11.6.2) by

$$\frac{1}{2\pi} \int_{x-\pi}^{x+\pi} |g(y)| \left| K_N(x - y) - e^{-iN(x-y)} \kappa(x - y) + \overline{e^{-iN(x-y)} \kappa(x - y)} \right| dy \leq \pi\delta.$$

By dominated convergence and since $\kappa(x - y) = 0$ for $|x - y| > 1$, (11.6.3) equals

$$\frac{1}{2\pi} \lim_{r \rightarrow 0^+} \left| \int_{r < |x-y| < 1} g(y) \max(|x-y|, 0) K_N(x-y) dy \right|.$$

We bound the limit by a supremum and rewrite using (11.6.4),

$$\leq \frac{1}{2\pi} \sup_{r > 0} \left| \int_{r < |x-y| < 1} g(y) \left(e^{-iN(x-y)} \kappa(x-y) + \overline{e^{-iN(x-y)} \kappa(x-y)} \right) dy \right|$$

Using the triangle inequality, we further bound this by

$$\begin{aligned} &\leq \frac{1}{2\pi} \sup_{r > 0} \left| \int_{r < |x-y| < 1} g(y) e^{-iNy} \kappa(x-y) dy \right| \\ &\quad + \frac{1}{2\pi} \sup_{r > 0} \left| \int_{r < |x-y| < 1} \bar{g}(y) e^{-iNy} \kappa(x-y) dy \right|. \end{aligned}$$

By the definition (11.1.16) of T , this is

$$\leq \frac{1}{2\pi} (Tg(x) + T\bar{g}(x)).$$

□

Lemma 11.6.3 (real Carleson operator measurable). *Let f be a bounded measurable function on \mathbb{R} . Then Tf as defined in (11.1.16) is measurable.*

Proof. Since a countable supremum of measurable functions is measurable, it suffices to show that for every $n \in \mathbb{Z}$,

$$x \mapsto \sup_{r > 0} \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right|$$

is measurable. So let $n \in \mathbb{Z}$. Note that for each $x \in \mathbb{R}$, the function

$$r \mapsto \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right|$$

is continuous on $(0, \infty)$ since the integrand is locally bounded on the domain $0 < |x - y| < 1$ by the assumptions on f and Lemma 11.1.11. Thus, for each $x \in \mathbb{R}$,

$$\sup_{r > 0} \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right| = \sup_{r \in \mathbb{Q}_{>0}} \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right|$$

The right hand side is again a countable supremum so it remains to show that for every $r \in \mathbb{Q}_{>0}$,

$$x \mapsto \left| \int_{r < |x-y| < 1} f(y) \kappa(x-y) e^{iny} dy \right| = \left| \int \mathbf{1}_{\{r < |x-\cdot| < 1\}}(y) f(y) \kappa(x-y) e^{iny} dy \right|$$

is measurable, which follows from the fact that the integrand is measurable in (x, y) . □

Lemma 11.6.4 (partial Fourier sums of small). *Let $g : \mathbb{R} \rightarrow \mathbb{C}$ be a measurable 2π -periodic function such that for some $\delta > 0$ and every $x \in \mathbb{R}$,*

$$|g(x)| \leq \delta. \quad (11.6.5)$$

Then for every $\epsilon > 0$, there exists a measurable set $E \subset [0, 2\pi]$ with $|E| < \epsilon$ such that for every $x \in [0, 2\pi] \setminus E$ and $N > 0$,

$$|S_N g(x)| \leq C_\epsilon \delta, \quad (11.6.6)$$

where

$$C_\epsilon = \left(\frac{8}{\pi\epsilon}\right)^{\frac{1}{2}} C_{4,2} + \pi. \quad (11.6.7)$$

Proof. Define

$$E := \{x \in [0, 2\pi] : \sup_{N>0} |S_N g(x)| > C_\epsilon \delta\}.$$

Then (11.6.6) clearly holds, and it remains to show that $|E| \leq \epsilon$. With Lemma 11.6.2, we obtain

$$E \subset \{x \in [0, 2\pi] : C_\epsilon \delta < \frac{1}{2\pi}(Tg(x) + T\bar{g}(x)) + \pi\delta\} \subset E_1 \cup E_2,$$

where

$$\begin{aligned} E_1 &:= \{x \in [0, 2\pi] : \pi(C_\epsilon - \pi)\delta < Tg(x)\} \\ E_2 &:= \{x \in [0, 2\pi] : \pi(C_\epsilon - \pi)\delta < T\bar{g}(x)\}. \end{aligned}$$

By Lemma 11.6.3, E_1 and E_2 are measurable. Thus,

$$\pi(C_\epsilon - \pi)\delta|E_1| \leq \int_{E_1} Tg(x) dx = \delta \int_{E_1} T(\delta^{-1}g\mathbf{1}_{[-\pi, 3\pi]})(x) dx.$$

Applying Lemma 11.1.5 with $F = [-\pi, 3\pi]$ and $G = E'$, it follows that this is

$$\leq \delta \cdot C_{4,2} |F|^{\frac{1}{2}} |E_1|^{\frac{1}{2}} \leq (4\pi)^{\frac{1}{2}} C_{4,2} \delta \cdot |E'|^{\frac{1}{2}}.$$

Rearranging, we obtain

$$|E_1| \leq \left(\frac{(4\pi)^{\frac{1}{2}} C_{4,2}}{\pi(C_\epsilon - \pi)}\right)^2 = \frac{\epsilon}{2}.$$

Analogously, we get the same estimate for $|E_2|$. This completes the proof using $|E| \leq |E_1| + |E_2|$. \square

Proof of Lemma 11.1.3. Lemma 11.1.3 follows directly from Lemma 11.6.4 with $\delta := \epsilon'$. \square

11.7. Carleson on the real line. We prove Lemma 11.1.5.

Consider the standard distance function

$$\rho(x, y) = |x - y| \quad (11.7.1)$$

on the real line \mathbb{R} .

Lemma 11.7.1 (real line metric). *The space (\mathbb{R}, ρ) is a complete locally compact metric space.*

Proof. This is part of the Lean library. \square

Lemma 11.7.2 (real line ball). *For $x \in \mathbb{R}$ and $R > 0$, the ball $B(x, R)$ is the interval $(x - R, x + R)$*

Proof. Let $x' \in B(x, R)$. By definition of the ball, $|x' - x| < R$. It follows that $x' - x < R$ and $x - x' < R$. It follows $x' < x + R$ and $x' > x - R$. This implies $x' \in (x - R, x + R)$. Conversely, let $x' \in (x - R, x + R)$. Then $x' < x + R$ and $x' > x - R$. It follows that $x' - x < R$ and $x - x' < R$. It follows that $|x' - x| < R$, hence $x' \in B(x, R)$. This proves the lemma. \square

We consider the Lebesgue measure μ on \mathbb{R} .

Lemma 11.7.3 (real line measure). *The measure μ is a sigma-finite non-zero Radon-Borel measure on \mathbb{R} .*

Proof. This is part of the Lean library. \square

Lemma 11.7.4 (real line ball measure). *We have for every $x \in \mathbb{R}$ and $R > 0$*

$$\mu(B(x, R)) = 2R. \quad (11.7.2)$$

Proof. We have with Lemma 11.7.2

$$\mu(B(x, R)) = \mu((x - R, x + R)) = 2R. \quad (11.7.3)$$

\square

Lemma 11.7.5 (real line doubling). *We have for every $x \in \mathbb{R}$ and $R > 0$*

$$\mu(B(x, 2R)) = 2\mu(B(x, R)). \quad (11.7.4)$$

Proof. We have with Lemma 11.7.4

$$\mu(B(x, 2R)) = 4R = 2\mu(B(x, R)). \quad (11.7.5)$$

This proves the lemma. \square

The preceding four lemmas show that $(\mathbb{R}, \rho, \mu, 4)$ is a doubling metric measure space. Indeed, we even show that $(\mathbb{R}, \rho, \mu, 1)$ is a doubling metric measure space, but we may relax the estimate in Lemma 11.7.5 to conclude that $(\mathbb{R}, \rho, \mu, 4)$ is a doubling metric measure space.

For each $n \in \mathbb{Z}$ define $\vartheta_n : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\vartheta_n(x) = nx. \quad (11.7.6)$$

Let Θ be the collection $\{\vartheta_n, n \in \mathbb{Z}\}$. Note that for every $n \in \mathbb{Z}$ we have $\vartheta_n(0) = 0$. Define

$$d_{B(x, R)}(\vartheta_n, \vartheta_m) := 2R|n - m|. \quad (11.7.7)$$

Lemma 11.7.6 (frequency metric). *For every $R > 0$ and $x \in X$, the function $d_{B(x,R)}$ is a metric on Θ .*

Proof. This follows immediately from the fact that the standard metric on \mathbb{Z} is a metric. \square

Lemma 11.7.7 (oscillation control). *For every $R > 0$ and $x \in X$, and for all $n, m \in \mathbb{Z}$, we have*

$$\sup_{y, y' \in B(x, R)} |ny - ny' - my + my'| \leq 2|n - m|R. \quad (11.7.8)$$

Proof. The right hand side of (11.7.8) equals

$$\sup_{y, y' \in B(x, R)} |(n - m)(y - x) - (n - m)(y' - x)|.$$

The lemma then follows from the triangle inequality. \square

Lemma 11.7.8 (frequency monotone). *For any $x, x' \in X$ and $R, R' > 0$ with $B(x, R) \subset B(x, R')$, and for any $n, m \in \mathbb{Z}$*

$$d_{B(x, R)}(\vartheta_n, \vartheta_m) \leq d_{B(x', R')}(\vartheta_n, \vartheta_m).$$

Proof. This follows immediately from the definition (11.7.7) and $R \leq R'$. \square

Lemma 11.7.9 (frequency ball doubling). *For any $x, x' \in \mathbb{R}$ and $R > 0$ with $x \in B(x', 2R)$ and any $n, m \in \mathbb{Z}$, we have*

$$d_{B(x', 2R)}(\vartheta_n, \vartheta_m) \leq 2d_{B(x, R)}(\vartheta_n, \vartheta_m). \quad (11.7.9)$$

Proof. With (11.7.7), both sides of (11.7.9) are equal to $4R|n - m|$. This proves the lemma. \square

Lemma 11.7.10 (frequency ball growth). *For any $x, x' \in \mathbb{R}$ and $R > 0$ with $B(x, R) \subset B(x', 2R)$ and any $n, m \in \mathbb{Z}$, we have*

$$2d_{B(x, R)}(\vartheta_n, \vartheta_m) \leq d_{B(x', 2R)}(\vartheta_n, \vartheta_m). \quad (11.7.10)$$

Proof. With (11.7.7), both sides of (11.7.9) are equal to $4R|n - m|$. This proves the lemma. \square

Lemma 11.7.11 (integer ball cover). *For every $x \in \mathbb{R}$ and $R > 0$ and every $n \in \mathbb{Z}$ and $R' > 0$, there exist $m_1, m_2, m_3 \in \mathbb{Z}$ such that*

$$B' \subset B_1 \cup B_2 \cup B_3, \quad (11.7.11)$$

where

$$B' = \{\vartheta \in \Theta : d_{B(x, R)}(\vartheta, \vartheta_n) < 2R'\} \quad (11.7.12)$$

and for $j = 1, 2, 3$

$$B_j = \{\vartheta \in \Theta : d_{B(x, R)}(\vartheta, \vartheta_{m_j}) < R'\}. \quad (11.7.13)$$

Proof. Let m_1 be the largest integer smaller than or equal to $n - \frac{R'}{2R}$. Let $m_2 = n$. Let m_3 be the smallest integer larger than or equal to $n + \frac{R'}{2R}$.

Let $\vartheta_{n'} \in B'$, then with (11.7.7), we have

$$2R|n - n'| < 2R'. \quad (11.7.14)$$

Assume first $n' \leq m_1$. With (11.7.14) we have

$$\begin{aligned} R|m_1 - n'| &= R(m_1 - n') = R(m_1 - n) + R(n - n') \\ &< -\frac{R'}{2} + R' = -\frac{R'}{2}. \end{aligned} \quad (11.7.15)$$

We conclude $\vartheta_{n'} \in B_1$.

Assume next $m_1 < n' < m_3$. Then $\vartheta_{n'} \in B_2$.

Assume finally that $m_3 \leq n'$. With (11.7.14) we have

$$\begin{aligned} R|m_3 - n'| &= R(n' - m_3) = R(n' - n) + R(n - m_3) \\ &< R' - \frac{R'}{2} = -\frac{R'}{2}. \end{aligned} \quad (11.7.16)$$

We conclude $\vartheta_{n'} \in B_1$. This completes the proof of the lemma. \square

Lemma 11.7.12 (real van der Corput). *For any $x \in \mathbb{R}$ and $R > 0$ and any function $\varphi : X \rightarrow \mathbb{C}$ supported on $B' = B(x, R)$ such that*

$$\|\varphi\|_{\text{Lip}(B')} = \sup_{x \in B'} |\varphi(x)| + R \sup_{x, y \in B', x \neq y} \frac{|\varphi(x) - \varphi(y)|}{\rho(x, y)} \quad (11.7.17)$$

is finite and for any $n, m \in \mathbb{Z}$, we have

$$\left| \int_{B'} e(\vartheta_n(x) - \vartheta_m(x)) \varphi(x) d\mu(x) \right| \leq 2\pi\mu(B') \frac{\|\varphi\|_{\text{Lip}(B')}}{1 + d_{B'}(\vartheta_n, \vartheta_m)}. \quad (11.7.18)$$

Proof. Set $n' = n - m$. Then we have to prove

$$\left| \int_{x-R}^{x+R} e^{in'y} \varphi(y) dy \right| \leq 4\pi R \|\varphi\|_{\text{Lip}(B')} (1 + 2R|n'|)^{-1}. \quad (11.7.19)$$

This follows from Lemma 11.1.7 with $\alpha = x - R$ and $\beta = x + R$. \square

Proof of Lemma 11.1.5. The preceding chain of lemmas establishes that Θ is a cancellative, compatible collection of functions on $(\mathbb{R}, \rho, \mu, 4)$. Again, some of the statements in these lemmas are stronger than what is needed for $a = 4$, but can be relaxed to give the desired conclusion for $a = 4$.

With κ as near (11.1.13), define the function $K : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{C}$ as in Theorem 1.1.1 by

$$K(x, y) := \kappa(x - y). \quad (11.7.20)$$

The function K is continuous outside the diagonal $x = y$ and vanishes on the diagonal. Hence it is measurable.

By Lemmas 11.1.11 and 11.1.12, it follows that K is a two-sided Calderón–Zygmund kernel on $(\mathbb{R}, \rho, \mu, 4)$. Lemma 11.1.6 verifies (10.0.3). Thus the assumptions of Theorem 10.0.1 are all satisfied. Applying the Theorem, Lemma 11.1.5 follows. \square

REFERENCES

- [Bec24] L. Becker. Maximal polynomial modulations of singular Radon transforms. *J. Funct. Anal.*, 286(6):Paper No. 110299, 46, 2024 (cited on page 5).
- [Bil67] P. Billard. Sur la convergence presque partout des séries de Fourier-Walsh des fonctions de l'espace $L^2(0, 1)$. *Studia Math.*, 28:363–388, 1966/67 (cited on page 4).
- [BvDJ⁺24] L. Becker, F. van Doorn, A. Jamneshan, R. Srivastava, and C. Thiele. Carleson operators on doubling metric measure spaces, 2024. eprint: 2405.06423v2 (cited on pages 1, 5).
- [Car66] L. Carleson. On convergence and growth of partial sums of Fourier series. *Acta Math.*, 116:135–157, 1966 (cited on pages 1, 4).
- [Chr90] M. Christ. A T(b) theorem with remarks on analytic capacity and the Cauchy integral. *Colloquium Mathematicum*, 2(60-61):601–628, 1990 (cited on page 21).
- [Dem15] C. Demeter. A guide to Carleson's theorem. *Rocky Mountain J. Math.*, 45(1):169–212, 2015 (cited on page 5).
- [DLT⁺08] C. Demeter, M. T. Lacey, T. Tao, and C. Thiele. Breaking the duality in the return times theorem. *Duke Math. J.*, 143(2):281–355, 2008 (cited on page 4).
- [dMU21] L. de Moura and S. Ullrich. The Lean 4 Theorem Prover and Programming Language. In A. Platzer and G. Sutcliffe, editors, *Automated Deduction – CADE 28: 28th International Conference on Automated Deduction, Virtual Event, July 12–15, 2021, Proceedings*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635, Cham. Springer International Publishing, 2021 (cited on page 5).
- [Fef73] C. Fefferman. Pointwise convergence of Fourier series. *Ann. of Math. (2)*, 98:551–571, 1973 (cited on pages 4, 32).
- [Fou13] J. B. J. Fourier. *Œuvres de Fourier. Vol. 1*. J. G. Darboux, editor. Cambridge Library Collection. Cambridge University Press, Cambridge, 2013, pages xxviii+563. Reprint of the 1888 original (cited on page 2).
- [Fou22] J. B. J. Fourier. *Théorie Analytique de la Chaleur*. Institut de France, Paris, 1822 (cited on page 2).
- [Hun68] R. A. Hunt. On the convergence of Fourier series. In *Orthogonal Expansions and their Continuous Analogues (Proc. Conf., Edwardsville, Ill., 1967)*, pages 235–255. Southern Illinois Univ. Press, Carbondale, IL, 1968 (cited on page 4).
- [Kol23] A. Kolmogorov. Une série de Fourier–Lebesgue divergente presque partout. *Fund. Math.*, 4:324–328, 1923 (cited on page 3).
- [Lac04] M. T. Lacey. Carleson's theorem: proof, complements, variations. *Publ. Mat.*, 48(2):251–307, 2004 (cited on page 5).
- [Lie09] V. Lie. The (weak- L^2) boundedness of the quadratic Carleson operator. *Geom. Funct. Anal.*, 19(2):457–497, 2009 (cited on page 4).
- [Lie20] V. Lie. The polynomial Carleson operator. *Ann. of Math. (2)*, 192(1):47–163, 2020 (cited on page 4).

- [LT00] M. Lacey and C. Thiele. A proof of boundedness of the Carleson operator. *Math. Res. Lett.*, 7(4):361–370, 2000 (cited on page 4).
- [Luz13] N. Luzin. Sur la convergence des séries trigonométrique de Fourier. *C. R. Acad. Sci. Paris*, 156:1655–1658, 1913 (cited on page 3).
- [Luz53] N. Luzin. *Sobranie sochinenii. Tom 1 (Collected works, Vol. 1)*. USSR Academy of Sciences, Moscow, 1953 (cited on page 3).
- [mat20] T. mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs, CPP 2020*, pages 367–381, New York, NY, USA. Association for Computing Machinery, 2020 (cited on page 5).
- [Mna22] G. Mnatsakanyan. On almost everywhere convergence of Malmquist–Takenaka series. *J. Funct. Anal.*, 282(12):Paper No. 109461, 33, 2022 (cited on page 5).
- [OST⁺12] R. Oberlin, A. Seeger, T. Tao, C. Thiele, and J. Wright. A variation norm Carleson theorem. *J. Eur. Math. Soc. (JEMS)*, 14(2):421–464, 2012 (cited on page 4).
- [PT03] M. Pramanik and E. Terwilleger. A weak L^2 estimate for a maximal dyadic sum operator on \mathbb{R}^n . *Illinois J. Math.*, 47(3):775–813, 2003 (cited on page 4).
- [Ram21] J. P. G. Ramos. The Hilbert transform along the parabola, the polynomial Carleson theorem and oscillatory singular integrals. *Math. Ann.*, 379(1-2):159–185, 2021 (cited on page 5).
- [Roo19] J. Roos. Bounds for anisotropic Carleson operators. *J. Fourier Anal. Appl.*, 25(5):2324–2355, 2019 (cited on page 4).
- [Sjö71] P. Sjölin. Convergence almost everywhere of certain singular integrals and multiple Fourier series. *Ark. Mat.*, 9:65–90, 1971 (cited on page 4).
- [Ste93] E. M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993, pages xiv+695. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III (cited on pages 112, 113, 115).
- [SW01] E. M. Stein and S. Wainger. Oscillatory integrals related to Carleson’s theorem. *Math. Res. Lett.*, 8(5-6):789–800, 2001 (cited on page 4).
- [Zor21] P. Zorin-Kranich. Maximal polynomial modulations of singular integrals. *Adv. Math.*, 386:Paper No. 107832, 40, 2021 (cited on pages 5, 12, 21).

- LARS BECKER, UNIVERSITY OF BONN
Email address: becker@math.uni-bonn.de
- MARÍA INÉS DE FRUTOS-FERNÁNDEZ, UNIVERSITY OF BONN
Email address: midff@math.uni-bonn.de
- LEO DIEDERING, UNIVERSITY OF BONN
Email address: leo.diedering@uni-bonn.de
- FLORIS VAN DOORN, UNIVERSITY OF BONN
Email address: vdoorn@math.uni-bonn.de
- SÉBASTIEN GOUÉZEL, UNIVERSITY OF RENNES
Email address: sebastien.gouezel@univ-rennes1.fr
- ASGAR JAMNESHAN, UNIVERSITY OF BONN
Email address: ajamnesh@math.uni-bonn.de
- EVGENIA KARUNUS, UNIVERSITY OF BONN
Email address: lakesare@gmail.com
- EDWARD VAN DE MEENT, UNIVERSITY OF UTRECHT
Email address: edwardvdmeent@gmail.com
- PIETRO MONTICONE, UNIVERSITY OF TRENTO
Email address: pit.monticone@gmail.com
- JASPER MULDER-SOHN, THE HAGUE
Email address: jasper.mulder@planet.nl
- JIM PORTEGIES, EINDHOVEN UNIVERSITY OF TECHNOLOGY
Email address: j.w.portegies@tue.nl
- JORIS ROOS, UNIVERSITY OF MASSACHUSETTS LOWELL
Email address: joris_roos@uml.edu
- MICHAEL ROTHGANG, UNIVERSITY OF BONN
Email address: rothgang@math.uni-bonn.de
- RAJULA SRIVASTAVA, UNIVERSITY OF BONN
Email address: rajulas@math.uni-bonn.de
- JAMES SUNDSTROM, BARUCH COLLEGE
Email address: james.sundstrom@baruch.cuny.edu
- JEREMY TAN, NATIONAL UNIVERSITY OF SINGAPORE
Email address: jtjierui@comp.nus.edu.sg
- CHRISTOPH THIELE, UNIVERSITY OF BONN
Email address: thiele@math.uni-bonn.de